




INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶ : C12N 15/31, C07K 14/35, 16/12, C12Q 1/68, C12N 15/62, G01N 33/53		A2	(11) International Publication Number: WO 98/16645
			(43) International Publication Date: 23 April 1998 (23.04.98)
(21) International Application Number: PCT/US97/18214		(74) Agents: MAKI, David, J. et al.; Seed and Berry LLP, 6300 Columbia Center, 701 Fifth Avenue, Seattle, WA 98104-7092 (US).	
(22) International Filing Date: 7 October 1997 (07.10.97)			
(30) Priority Data: 08/729,622 11 October 1996 (11.10.96) US 08/818,111 13 March 1997 (13.03.97) US		(81) Designated States: AL, AM, AT, AU, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GE, GH, HU, ID, IL, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, UZ, VN, YU, ARIPO patent (GH, KE, LS, MW, SD, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG).	
(71) Applicant: CORIXA CORPORATION [US/US]; 1124 Columbia Street, Seattle, WA 98104 (US).			
(72) Inventors: REED, Steven, G.; 2843 - 122nd Place N.E., Bellevue, WA 98005 (US). SKEIKY, Yasir, A., W.; 8327 - 25th Avenue N.W., Seattle, WA 98107 (US). DILLON, Davin, C.; 21607 N.E. 24th Street, Redmond, WA 98053 (US). CAMPOS-NETO, Antonio; 9308 Midship Court N.E., Bainbridge Island, WA 98021 (US). HOUGHTON, Raymond; 2636 - 242nd Place S.E., Bothell, WA 98021 (US). VEDVICK, Thomas, S.; 124 South 300th Place, Federal Way, WA 98003 (US). TWARDZIK, Daniel, R.; 10195 South Beach Drive, Bainbridge Island, WA 98110 (US). LODES, Michael, J.; 9223 - 36th Avenue S.W., Seattle, WA 98126 (US).		Published Without international search report and to be republished upon receipt of that report.	

(54) Title: COMPOUNDS AND METHODS FOR DIAGNOSIS OF TUBERCULOSIS



(57) Abstract

Compounds and methods for diagnosing tuberculosis are disclosed. The compounds provided include polypeptides that contain at least one antigenic portion of one or more *M. tuberculosis* proteins, and DNA sequences encoding such polypeptides. Diagnostic kits containing such polypeptides or DNA sequences and a suitable detection reagent may be used for the detection of *M. tuberculosis* infection in patients and biological samples. Antibodies directed against such polypeptides are also provided.

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece			TR	Turkey
BG	Bulgaria	HU	Hungary	ML	Mali	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MN	Mongolia	UA	Ukraine
BR	Brazil	IL	Israel	MR	Mauritania	UG	Uganda
BY	Belarus	IS	Iceland	MW	Malawi	US	United States of America
CA	Canada	IT	Italy	MX	Mexico	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NE	Niger	VN	Viet Nam
CG	Congo	KE	Kenya	NL	Netherlands	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NO	Norway	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	NZ	New Zealand		
CM	Cameroon			PL	Poland		
CN	China	KR	Republic of Korea	PT	Portugal		
CU	Cuba	KZ	Kazakstan	RO	Romania		
CZ	Czech Republic	LC	Saint Lucia	RU	Russian Federation		
DE	Germany	LI	Liechtenstein	SD	Sudan		
DK	Denmark	LK	Sri Lanka	SE	Sweden		
EE	Estonia	LR	Liberia	SG	Singapore		

COMPOUNDS AND METHODS FOR DIAGNOSIS OF TUBERCULOSIS

TECHNICAL FIELD

The present invention relates generally to the detection of *Mycobacterium tuberculosis* infection. The invention is more particularly related to polypeptides comprising a *Mycobacterium tuberculosis* antigen, or a portion or other variant thereof, and the use of such polypeptides for the serodiagnosis of *Mycobacterium tuberculosis* infection.

BACKGROUND OF THE INVENTION

Tuberculosis is a chronic, infectious disease, that is generally caused by infection with *Mycobacterium tuberculosis*. It is a major disease in developing countries, as well as an increasing problem in developed areas of the world, with about 8 million new cases and 3 million deaths each year. Although the infection may be asymptomatic for a considerable period of time, the disease is most commonly manifested as an acute inflammation of the lungs, resulting in fever and a nonproductive cough. If left untreated, serious complications and death typically result.

Although tuberculosis can generally be controlled using extended antibiotic therapy, such treatment is not sufficient to prevent the spread of the disease. Infected individuals may be asymptomatic, but contagious, for some time. In addition, although compliance with the treatment regimen is critical, patient behavior is difficult to monitor. Some patients do not complete the course of treatment, which can lead to ineffective treatment and the development of drug resistance.

Inhibiting the spread of tuberculosis will require effective vaccination and accurate, early diagnosis of the disease. Currently, vaccination with live bacteria is the most efficient method for inducing protective immunity. The most common *Mycobacterium* for this purpose is Bacillus Calmette-Guerin (BCG), an avirulent strain of *Mycobacterium bovis*. However, the safety and efficacy of BCG is a source of controversy and some countries, such as the United States, do not vaccinate the general public. Diagnosis is commonly achieved using a skin test, which involves intradermal exposure to tuberculin PPD (protein-purified derivative). Antigen-specific T cell responses result in measurable incubation at the injection

site by 48-72 hours after injection, which indicates exposure to Mycobacterial antigens. Sensitivity and specificity have, however, been a problem with this test, and individuals vaccinated with BCG cannot be distinguished from infected individuals.

While macrophages have been shown to act as the principal effectors of
5 *M. tuberculosis* immunity, T cells are the predominant inducers of such immunity. The essential role of T cells in protection against *M. tuberculosis* infection is illustrated by the frequent occurrence of *M. tuberculosis* in AIDS patients, due to the depletion of CD4 T cells associated with human immunodeficiency virus (HIV) infection. Mycobacterium-reactive CD4 T cells have been shown to be potent producers of gamma-interferon (IFN- γ), which, in
10 turn, has been shown to trigger the anti-mycobacterial effects of macrophages in mice. While the role of IFN- γ in humans is less clear, studies have shown that 1,25-dihydroxy-vitamin D3, either alone or in combination with IFN- γ or tumor necrosis factor-alpha, activates human macrophages to inhibit *M. tuberculosis* infection. Furthermore, it is known that IFN- γ stimulates human macrophages to make 1,25-dihydroxy-vitamin D3. Similarly, IL-12 has
15 been shown to play a role in stimulating resistance to *M. tuberculosis* infection. For a review of the immunology of *M. tuberculosis* infection see Chan and Kaufmann, in *Tuberculosis: Pathogenesis, Protection and Control*, Bloom (ed.), ASM Press, Washington, DC, 1994.

Accordingly, there is a need in the art for improved diagnostic methods for detecting tuberculosis. The present invention fulfills this need and further provides other
20 related advantages.

SUMMARY OF THE INVENTION

Briefly stated, the present invention provides compositions and methods for diagnosing tuberculosis. In one aspect, polypeptides are provided comprising an antigenic
25 portion of a soluble *M. tuberculosis* antigen, or a variant of such an antigen that differs only in conservative substitutions and/or modifications. In one embodiment of this aspect, the soluble antigen has one of the following N-terminal sequences:

- (a) Asp-Pro-Val-Asp-Ala-Val-Ile-Asn-Thr-Thr-Cys-Asn-Tyr-Gly-Gln-Val-Val-Ala-Ala-Leu (SEQ ID NO: 115);

- (b) Ala-Val-Glu-Ser-Gly-Met-Leu-Ala-Leu-Gly-Thr-Pro-Ala-Pro-Ser
(SEQ ID NO: 116);
- (c) Ala-Ala-Met-Lys-Pro-Arg-Thr-Gly-Asp-Gly-Pro-Leu-Glu-Ala-Ala-
Lys-Glu-Gly-Arg (SEQ ID NO: 117);
- 5 (d) Tyr-Tyr-Trp-Cys-Pro-Gly-Gln-Pro-Phe-Asp-Pro-Ala-Trp-Gly-Pro
(SEQ ID NO: 118);
- (e) Asp-Ile-Gly-Ser-Glu-Ser-Thr-Glu-Asp-Gln-Gln-Xaa-Ala-Val (SEQ ID
NO: 119);
- (f) Ala-Glu-Glu-Ser-Ile-Ser-Thr-Xaa-Glu-Xaa-Ile-Val-Pro (SEQ ID
10 NO: 120);
- (g) Asp-Pro-Glu-Pro-Ala-Pro-Pro-Val-Pro-Thr-Thr-Ala-Ala-Ser-Pro-Pro-
Ser (SEQ ID NO: 121);
- (h) Ala-Pro-Lys-Thr-Tyr-Xaa-Glu-Glu-Leu-Lys-Gly-Thr-Asp-Thr-Gly
(SEQ ID NO: 122);
- 15 (i) Asp-Pro-Ala-Ser-Ala-Pro-Asp-Val-Pro-Thr-Ala-Ala-Gln-Leu-Thr-Ser-
Leu-Leu-Asn-Ser-Leu-Ala-Asp-Pro-Asn-Val-Ser-Phe-Ala-Asn (SEQ
ID NO: 123);
- (j) Xaa-Asp-Ser-Glu-Lys-Ser-Ala-Thr-Ile-Lys-Val-Thr-Asp-Ala-Ser;
(SEQ ID NO: 129)
- 20 (k) Ala-Gly-Asp-Thr-Xaa-Ile-Tyr-Ile-Val-Gly-Asn-Leu-Thr-Ala-Asp;
(SEQ ID NO: 130) or
- (l) Ala-Pro-Glu-Ser-Gly-Ala-Gly-Leu-Gly-Gly-Thr-Val-Gln-Ala-Gly;
(SEQ ID NO: 131)

25 wherein Xaa may be any amino acid.

In a related aspect, polypeptides are provided comprising an immunogenic portion of an *M. tuberculosis* antigen, or a variant of such an antigen that differs only in conservative substitutions and/or modifications, the antigen having one of the following N-terminal sequences:

- (m) Xaa-Tyr-Ile-Ala-Tyr-Xaa-Thr-Thr-Ala-Gly-Ile-Val-Pro-Gly-Lys-Ile-Asn-Val-His-Leu-Val; (SEQ ID NO: 132) or
- (n) Asp-Pro-Pro-Asp-Pro-His-Gln-Xaa-Asp-Met-Thr-Lys-Gly-Tyr-Tyr-Pro-Gly-Gly-Arg-Arg-Xaa-Phe; (SEQ ID NO: 124)

5 wherein Xaa may be any amino acid.

In another embodiment, the soluble *M. tuberculosis* antigen comprises an amino acid sequence encoded by a DNA sequence selected from the group consisting of the sequences recited in SEQ ID NOS: 1, 2, 4-10, 13-25, 52, 94 and 96, the complements of said sequences, and DNA sequences that hybridize to a sequence recited in SEQ ID NOS: 1, 2,
10 4-10, 13-25, 52, 94 and 96 or a complement thereof under moderately stringent conditions.

In a related aspect, the polypeptides comprise an antigenic portion of a *M. tuberculosis* antigen, or a variant of such an antigen that differs only in conservative substitutions and/or modifications, wherein the antigen comprises an amino acid sequence encoded by a DNA sequence selected from the group consisting of the sequences recited in
15 SEQ ID NOS: 26-51, 133, 134, 158-178 and 196, the complements of said sequences, and DNA sequences that hybridize to a sequence recited in SEQ ID NOS: 26-51, 133, 134, 158-178 and 196 or a complement thereof under moderately stringent conditions.

In related aspects, DNA sequences encoding the above polypeptides, recombinant expression vectors comprising these DNA sequences and host cells transformed
20 or transfected with such expression vectors are also provided.

In another aspect, the present invention provides fusion proteins comprising a first and a second inventive polypeptide or, alternatively, an inventive polypeptide and a known *M. tuberculosis* antigen.

In further aspects of the subject invention, methods and diagnostic kits are
25 provided for detecting tuberculosis in a patient. The methods comprise: (a) contacting a biological sample with at least one of the above polypeptides; and (b) detecting in the sample the presence of antibodies that bind to the polypeptide or polypeptides, thereby detecting *M. tuberculosis* infection in the biological sample. Suitable biological samples include whole blood, sputum, serum, plasma, saliva, cerebrospinal fluid and urine. The diagnostic kits
30 comprise one or more of the above polypeptides in combination with a detection reagent.

The present invention also provides methods for detecting *M. tuberculosis* infection comprising: (a) obtaining a biological sample from a patient; (b) contacting the sample with at least one oligonucleotide primer in a polymerase chain reaction, the oligonucleotide primer being specific for a DNA sequence encoding the above polypeptides; and (c) detecting in the sample a DNA sequence that amplifies in the presence of the first and second oligonucleotide primers. In one embodiment, the oligonucleotide primer comprises at least about 10 contiguous nucleotides of such a DNA sequence.

In a further aspect, the present invention provides a method for detecting *M. tuberculosis* infection in a patient comprising: (a) obtaining a biological sample from the patient; (b) contacting the sample with an oligonucleotide probe specific for a DNA sequence encoding the above polypeptides; and (c) detecting in the sample a DNA sequence that hybridizes to the oligonucleotide probe. In one embodiment, the oligonucleotide probe comprises at least about 15 contiguous nucleotides of such a DNA sequence.

In yet another aspect, the present invention provides antibodies, both polyclonal and monoclonal, that bind to the polypeptides described above, as well as methods for their use in the detection of *M. tuberculosis* infection.

These and other aspects of the present invention will become apparent upon reference to the following detailed description and attached drawings. All references disclosed herein are hereby incorporated by reference in their entirety as if each was incorporated individually.

BRIEF DESCRIPTION OF THE DRAWINGS AND SEQUENCE IDENTIFIERS

Figure 1A and B illustrate the stimulation of proliferation and interferon- γ production in T cells derived from a first and a second *M. tuberculosis*-immune donor, respectively, by the 14 Kd, 20 Kd and 26 Kd antigens described in Example 1.

Figures 2A-D illustrate the reactivity of antisera raised against secretory *M. tuberculosis* proteins, the known *M. tuberculosis* antigen 85b and the inventive antigens Tb38-1 and TbH-9, respectively, with *M. tuberculosis* lysate (lane 2), *M. tuberculosis* secretory proteins (lane 3), recombinant Tb38-1 (lane 4), recombinant TbH-9 (lane 5) and recombinant 85b (lane 5).

Figure 3A illustrates the stimulation of proliferation in a TbH-9-specific T cell clone by secretory *M. tuberculosis* proteins, recombinant TbH-9 and a control antigen, TbRa11.

Figure 3B illustrates the stimulation of interferon- γ production in a TbH-9-specific T cell clone by secretory *M. tuberculosis* proteins, PPD and recombinant TbH-9.

Figure 4 illustrates the reactivity of two representative polypeptides with sera from *M. tuberculosis*-infected and uninfected individuals, as compared to the reactivity of bacterial lysate.

Figure 5 shows the reactivity of four representative polypeptides with sera from *M. tuberculosis*-infected and uninfected individuals, as compared to the reactivity of the 38 kD antigen.

Figure 6 shows the reactivity of recombinant 38 kD and TbRa11 antigens with sera from *M. tuberculosis* patients, PPD positive donors and normal donors.

Figure 7 shows the reactivity of the antigen TbRa2A with 38 kD negative sera.

Figure 8 shows the reactivity of the antigen of SEQ ID NO: 60 with sera from *M. tuberculosis* patients and normal donors.

Figure 9 illustrates the reactivity of the recombinant antigen TbH-29 (SEQ ID NO: 137) with sera from *M. tuberculosis* patients, PPD positive donors and normal donors as determined by indirect ELISA.

Figure 10 illustrates the reactivity of the recombinant antigen TbH-33 (SEQ ID NO: 140) with sera from *M. tuberculosis* patients and from normal donors, and with a pool of sera from *M. tuberculosis* patients, as determined both by direct and indirect ELISA.

Figure 11 illustrates the reactivity of increasing concentrations of the recombinant antigen TbH-33 (SEQ ID NO: 140) with sera from *M. tuberculosis* patients and from normal donors as determined by ELISA.

SEQ. ID NO. 1 is the DNA sequence of TbRa1.

SEQ. ID NO. 2 is the DNA sequence of TbRa10.

SEQ. ID NO. 3 is the DNA sequence of TbRa11.

SEQ. ID NO. 4 is the DNA sequence of TbRa12.

- SEQ. ID NO. 5 is the DNA sequence of TbRa13.
SEQ. ID NO. 6 is the DNA sequence of TbRa16.
SEQ. ID NO. 7 is the DNA sequence of TbRa17.
SEQ. ID NO. 8 is the DNA sequence of TbRa18.
5 SEQ. ID NO. 9 is the DNA sequence of TbRa19.
SEQ. ID NO. 10 is the DNA sequence of TbRa24.
SEQ. ID NO. 11 is the DNA sequence of TbRa26.
SEQ. ID NO. 12 is the DNA sequence of TbRa28.
SEQ. ID NO. 13 is the DNA sequence of TbRa29.
10 SEQ. ID NO. 14 is the DNA sequence of TbRa2A.
SEQ. ID NO. 15 is the DNA sequence of TbRa3.
SEQ. ID NO. 16 is the DNA sequence of TbRa32.
SEQ. ID NO. 17 is the DNA sequence of TbRa35.
SEQ. ID NO. 18 is the DNA sequence of TbRa36.
15 SEQ. ID NO. 19 is the DNA sequence of TbRa4.
SEQ. ID NO. 20 is the DNA sequence of TbRa9.
SEQ. ID NO. 21 is the DNA sequence of TbRaB.
SEQ. ID NO. 22 is the DNA sequence of TbRaC.
SEQ. ID NO. 23 is the DNA sequence of TbRaD.
20 SEQ. ID NO. 24 is the DNA sequence of YYWCPG.
SEQ. ID NO. 25 is the DNA sequence of AAMK.
SEQ. ID NO. 26 is the DNA sequence of TbL-23.
SEQ. ID NO. 27 is the DNA sequence of TbL-24.
SEQ. ID NO. 28 is the DNA sequence of TbL-25.
25 SEQ. ID NO. 29 is the DNA sequence of TbL-28.
SEQ. ID NO. 30 is the DNA sequence of TbL-29.
SEQ. ID NO. 31 is the DNA sequence of TbH-5.
SEQ. ID NO. 32 is the DNA sequence of TbH-8.
SEQ. ID NO. 33 is the DNA sequence of TbH-9.
30 SEQ. ID NO. 34 is the DNA sequence of TbM-1.

- SEQ. ID NO. 35 is the DNA sequence of TbM-3.
SEQ. ID NO. 36 is the DNA sequence of TbM-6.
SEQ. ID NO. 37 is the DNA sequence of TbM-7.
SEQ. ID NO. 38 is the DNA sequence of TbM-9.
5 SEQ. ID NO. 39 is the DNA sequence of TbM-12.
SEQ. ID NO. 40 is the DNA sequence of TbM-13.
SEQ. ID NO. 41 is the DNA sequence of TbM-14.
SEQ. ID NO. 42 is the DNA sequence of TbM-15.
SEQ. ID NO. 43 is the DNA sequence of TbH-4.
10 SEQ. ID NO. 44 is the DNA sequence of TbH-4-FWD.
SEQ. ID NO. 45 is the DNA sequence of TbH-12.
SEQ. ID NO. 46 is the DNA sequence of Tb38-1.
SEQ. ID NO. 47 is the DNA sequence of Tb38-4.
SEQ. ID NO. 48 is the DNA sequence of TbL-17.
15 SEQ. ID NO. 49 is the DNA sequence of TbL-20.
SEQ. ID NO. 50 is the DNA sequence of TbL-21.
SEQ. ID NO. 51 is the DNA sequence of TbH-16.
SEQ. ID NO. 52 is the DNA sequence of DPEP.
SEQ. ID NO. 53 is the deduced amino acid sequence of DPEP.
20 SEQ. ID NO. 54 is the protein sequence of DPV N-terminal Antigen.
SEQ. ID NO. 55 is the protein sequence of AVGS N-terminal Antigen.
SEQ. ID NO. 56 is the protein sequence of AAMK N-terminal Antigen.
SEQ. ID NO. 57 is the protein sequence of YYWC N-terminal Antigen.
SEQ. ID NO. 58 is the protein sequence of DIGS N-terminal Antigen.
25 SEQ. ID NO. 59 is the protein sequence of AEES N-terminal Antigen.
SEQ. ID NO. 60 is the protein sequence of DPEP N-terminal Antigen.
SEQ. ID NO. 61 is the protein sequence of APKT N-terminal Antigen.
SEQ. ID NO. 62 is the protein sequence of DPAS N-terminal Antigen.
SEQ. ID NO. 63 is the deduced amino acid sequence of TbM-1 Peptide.
30 SEQ. ID NO. 64 is the deduced amino acid sequence of TbRa1.

- SEQ. ID NO. 65 is the deduced amino acid sequence of TbRa10.
SEQ. ID NO. 66 is the deduced amino acid sequence of TbRa11.
SEQ. ID NO. 67 is the deduced amino acid sequence of TbRa12.
SEQ. ID NO. 68 is the deduced amino acid sequence of TbRa13.
5 SEQ. ID NO. 69 is the deduced amino acid sequence of TbRa16.
SEQ. ID NO. 70 is the deduced amino acid sequence of TbRa17.
SEQ. ID NO. 71 is the deduced amino acid sequence of TbRa18.
SEQ. ID NO. 72 is the deduced amino acid sequence of TbRa19.
SEQ. ID NO. 73 is the deduced amino acid sequence of TbRa24.
10 SEQ. ID NO. 74 is the deduced amino acid sequence of TbRa26.
SEQ. ID NO. 75 is the deduced amino acid sequence of TbRa28.
SEQ. ID NO. 76 is the deduced amino acid sequence of TbRa29.
SEQ. ID NO. 77 is the deduced amino acid sequence of TbRa2A.
SEQ. ID NO. 78 is the deduced amino acid sequence of TbRa3.
15 SEQ. ID NO. 79 is the deduced amino acid sequence of TbRa32.
SEQ. ID NO. 80 is the deduced amino acid sequence of TbRa35.
SEQ. ID NO. 81 is the deduced amino acid sequence of TbRa36.
SEQ. ID NO. 82 is the deduced amino acid sequence of TbRa4.
SEQ. ID NO. 83 is the deduced amino acid sequence of TbRa9.
20 SEQ. ID NO. 84 is the deduced amino acid sequence of TbRaB.
SEQ. ID NO. 85 is the deduced amino acid sequence of TbRaC.
SEQ. ID NO. 86 is the deduced amino acid sequence of TbRaD.
SEQ. ID NO. 87 is the deduced amino acid sequence of YYWCPG.
SEQ. ID NO. 88 is the deduced amino acid sequence of TbAAMK.
25 SEQ. ID NO. 89 is the deduced amino acid sequence of Tb38-1.
SEQ. ID NO. 90 is the deduced amino acid sequence of TbH-4.
SEQ. ID NO. 91 is the deduced amino acid sequence of TbH-8.
SEQ. ID NO. 92 is the deduced amino acid sequence of TbH-9.
SEQ. ID NO. 93 is the deduced amino acid sequence of TbH-12.
30 SEQ. ID NO. 94 is the DNA sequence of DPAS.

- SEQ. ID NO. 95 is the deduced amino acid sequence of DPAS.
- SEQ. ID NO. 96 is the DNA sequence of DPV.
- SEQ. ID NO. 97 is the deduced amino acid sequence of DPV.
- SEQ. ID NO. 98 is the DNA sequence of ESAT-6.
- 5 SEQ. ID NO. 99 is the deduced amino acid sequence of ESAT-6.
- SEQ. ID NO. 100 is the DNA sequence of TbH-8-2.
- SEQ. ID NO. 101 is the DNA sequence of TbH-9FL.
- SEQ. ID NO. 102 is the deduced amino acid sequence of TbH-9FL.
- SEQ. ID NO. 103 is the DNA sequence of TbH-9-1.
- 10 SEQ. ID NO. 104 is the deduced amino acid sequence of TbH-9-1.
- SEQ. ID NO. 105 is the DNA sequence of TbH-9-4.
- SEQ. ID NO. 106 is the deduced amino acid sequence of TbH-9-4.
- SEQ. ID NO. 107 is the DNA sequence of Tb38-1F2 IN.
- SEQ. ID NO. 108 is the DNA sequence of Tb38-1F2 RP.
- 15 SEQ. ID NO. 109 is the deduced amino acid sequence of Tb37-FL.
- SEQ. ID NO. 110 is the deduced amino acid sequence of Tb38-IN.
- SEQ. ID NO. 111 is the DNA sequence of Tb38-1F3.
- SEQ. ID NO. 112 is the deduced amino acid sequence of Tb38-1F3.
- SEQ. ID NO. 113 is the DNA sequence of Tb38-1F5.
- 20 SEQ. ID NO. 114 is the DNA sequence of Tb38-1F6.
- SEQ. ID NO. 115 is the deduced N-terminal amino acid sequence of DPV.
- SEQ. ID NO. 116 is the deduced N-terminal amino acid sequence of AVGS.
- SEQ. ID NO. 117 is the deduced N-terminal amino acid sequence of AAMK.
- SEQ. ID NO. 118 is the deduced N-terminal amino acid sequence of YYWC.
- 25 SEQ. ID NO. 119 is the deduced N-terminal amino acid sequence of DIGS.
- SEQ. ID NO. 120 is the deduced N-terminal amino acid sequence of AAES.
- SEQ. ID NO. 121 is the deduced N-terminal amino acid sequence of DPEP.
- SEQ. ID NO. 122 is the deduced N-terminal amino acid sequence of APKT.
- SEQ. ID NO. 123 is the deduced N-terminal amino acid sequence of DPAS.
- 30 SEQ. ID NO. 124 is the protein sequence of DPPD N-terminal Antigen.

SEQ ID NO. 125-128 are the protein sequences of four DPPD cyanogen bromide fragments.

SEQ ID NO. 129 is the N-terminal protein sequence of XDS antigen.

SEQ ID NO. 130 is the N-terminal protein sequence of AGD antigen.

5 SEQ ID NO. 131 is the N-terminal protein sequence of APE antigen.

SEQ ID NO. 132 is the N-terminal protein sequence of XYI antigen.

SEQ ID NO. 133 is the DNA sequence of TbH-29.

SEQ ID NO. 134 is the DNA sequence of TbH-30.

SEQ ID NO. 135 is the DNA sequence of TbH-32.

10 SEQ ID NO. 136 is the DNA sequence of TbH-33.

SEQ ID NO. 137 is the predicted amino acid sequence of TbH-29.

SEQ ID NO. 138 is the predicted amino acid sequence of TbH-30.

SEQ ID NO. 139 is the predicted amino acid sequence of TbH-32.

SEQ ID NO. 140 is the predicted amino acid sequence of TbH-33.

15 SEQ ID NO: 141-146 are PCR primers used in the preparation of a fusion protein containing TbRa3, 38 kD and Tb38-1.

SEQ ID NO: 147 is the DNA sequence of the fusion protein containing TbRa3, 38 kD and Tb38-1.

20 SEQ ID NO: 148 is the amino acid sequence of the fusion protein containing TbRa3, 38 kD and Tb38-1.

SEQ ID NO: 149 is the DNA sequence of the M. tuberculosis antigen 38 kD.

SEQ ID NO: 150 is the amino acid sequence of the M. tuberculosis antigen 38 kD.

SEQ ID NO: 151 is the DNA sequence of XP14.

SEQ ID NO: 152 is the DNA sequence of XP24.

25 SEQ ID NO: 153 is the DNA sequence of XP31.

SEQ ID NO: 154 is the 5' DNA sequence of XP32.

SEQ ID NO: 155 is the 3' DNA sequence of XP32.

SEQ ID NO: 156 is the predicted amino acid sequence of XP14.

30 SEQ ID NO: 157 is the predicted amino acid sequence encoded by the reverse complement of XP14.

- SEQ ID NO: 158 is the DNA sequence of XP27.
- SEQ ID NO: 159 is the DNA sequence of XP36.
- SEQ ID NO: 160 is the 5' DNA sequence of XP4.
- SEQ ID NO: 161 is the 5' DNA sequence of XP5.
- 5 SEQ ID NO: 162 is the 5' DNA sequence of XP17.
- SEQ ID NO: 163 is the 5' DNA sequence of XP30.
- SEQ ID NO: 164 is the 5' DNA sequence of XP2.
- SEQ ID NO: 165 is the 3' DNA sequence of XP2.
- SEQ ID NO: 166 is the 5' DNA sequence of XP3.
- 10 SEQ ID NO: 167 is the 3' DNA sequence of XP3.
- SEQ ID NO: 168 is the 5' DNA sequence of XP6.
- SEQ ID NO: 169 is the 3' DNA sequence of XP6.
- SEQ ID NO: 170 is the 5' DNA sequence of XP18.
- SEQ ID NO: 171 is the 3' DNA sequence of XP18.
- 15 SEQ ID NO: 172 is the 5' DNA sequence of XP19.
- SEQ ID NO: 173 is the 3' DNA sequence of XP19.
- SEQ ID NO: 174 is the 5' DNA sequence of XP22.
- SEQ ID NO: 175 is the 3' DNA sequence of XP22.
- SEQ ID NO: 176 is the 5' DNA sequence of XP25.
- 20 SEQ ID NO: 177 is the 3' DNA sequence of XP25.
- SEQ ID NO: 178 is the full-length DNA sequence of TbH4-XP1.
- SEQ ID NO: 179 is the predicted amino acid sequence of TbH4-XP1.
- SEQ ID NO: 180 is the predicted amino acid sequence encoded by the reverse complement of TbH4-XP1.
- 25 SEQ ID NO: 181 is a first predicted amino acid sequence encoded by XP36.
- SEQ ID NO: 182 is a second predicted amino acid sequence encoded by XP36.
- SEQ ID NO: 183 is the predicted amino acid sequence encoded by the reverse complement of XP36.
- SEQ ID NO: 184 is the DNA sequence of RDIF2.
- 30 SEQ ID NO: 185 is the DNA sequence of RDIF5.

SEQ ID NO: 186 is the DNA sequence of RDIF8.

SEQ ID NO: 187 is the DNA sequence of RDIF10.

SEQ ID NO: 188 is the DNA sequence of RDIF11.

SEQ ID NO: 189 is the predicted amino acid sequence of RDIF2.

5 SEQ ID NO: 190 is the predicted amino acid sequence of RDIF5.

SEQ ID NO: 191 is the predicted amino acid sequence of RDIF8.

SEQ ID NO: 192 is the predicted amino acid sequence of RDIF10.

SEQ ID NO: 193 is the predicted amino acid sequence of RDIF11.

SEQ ID NO: 194 is the 5' DNA sequence of RDIF12.

10 SEQ ID NO: 195 is the 3' DNA sequence of RDIF12.

SEQ ID NO: 196 is the DNA sequence of RDIF7.

SEQ ID NO: 197 is the predicted amino acid sequence of RDIF7.

SEQ ID NO: 198 is the DNA sequence of DIF2-1.

SEQ ID NO: 199 is the predicted amino acid sequence of DIF2-1.

15 SEQ ID NO: 200-207 are PCR primers used in the preparation of a fusion protein containing TbRa3, 38 kD, Tb38-1 and DPEP (hereinafter referred to as TbF-2).

SEQ ID NO: 208 is the DNA sequence of the fusion protein TbF-2.

SEQ ID NO: 209 is the amino acid sequence of the fusion protein TbF-2.

20

DETAILED DESCRIPTION OF THE INVENTION

As noted above, the present invention is generally directed to compositions and methods for diagnosing tuberculosis. The compositions of the subject invention include polypeptides that comprise at least one antigenic portion of a *M. tuberculosis* antigen, or a
25 variant of such an antigen that differs only in conservative substitutions and/or modifications. Polypeptides within the scope of the present invention include, but are not limited to, soluble *M. tuberculosis* antigens. A "soluble *M. tuberculosis* antigen" is a protein of *M. tuberculosis* origin that is present in *M. tuberculosis* culture filtrate. As used herein, the term "polypeptide" encompasses amino acid chains of any length, including full length proteins
30 (*i.e.*, antigens), wherein the amino acid residues are linked by covalent peptide bonds. Thus,

a polypeptide comprising an antigenic portion of one of the above antigens may consist entirely of the antigenic portion, or may contain additional sequences. The additional sequences may be derived from the native *M. tuberculosis* antigen or may be heterologous, and such sequences may (but need not) be antigenic.

5 An "antigenic portion" of an antigen (which may or may not be soluble) is a portion that is capable of reacting with sera obtained from an *M. tuberculosis*-infected individual (*i.e.*, generates an absorbance reading with sera from infected individuals that is at least three standard deviations above the absorbance obtained with sera from uninfected individuals, in a representative ELISA assay described herein). An "*M. tuberculosis*-infected
10 individual" is a human who has been infected with *M. tuberculosis* (*e.g.*, has an intradermal skin test response to PPD that is at least 0.5 cm in diameter). Infected individuals may display symptoms of tuberculosis or may be free of disease symptoms. Polypeptides comprising at least an antigenic portion of one or more *M. tuberculosis* antigens as described herein may generally be used, alone or in combination, to detect tuberculosis in a patient.

15 The compositions and methods of this invention also encompass variants of the above polypeptides. A "variant," as used herein, is a polypeptide that differs from the native antigen only in conservative substitutions and/or modifications, such that the antigenic properties of the polypeptide are retained. Such variants may generally be identified by modifying one of the above polypeptide sequences, and evaluating the antigenic properties of
20 the modified polypeptide using, for example, the representative procedures described herein.

A "conservative substitution" is one in which an amino acid is substituted for another amino acid that has similar properties, such that one skilled in the art of peptide chemistry would expect the secondary structure and hydrophobic nature of the polypeptide to be substantially unchanged. In general, the following groups of amino acids represent
25 conservative changes: (1) ala, pro, gly, glu, asp, gln, asn, ser, thr; (2) cys, ser, tyr, thr; (3) val, ile, leu, met, ala, phe; (4) lys, arg, his; and (5) phe, tyr, trp, his.

Variants may also (or alternatively) be modified by, for example, the deletion or addition of amino acids that have minimal influence on the antigenic properties, secondary structure and hydrophobic nature of the polypeptide. For example, a polypeptide may be
30 conjugated to a signal (or leader) sequence at the N-terminal end of the protein which co-

translationally or post-translationally directs transfer of the protein. The polypeptide may also be conjugated to a linker or other sequence for ease of synthesis, purification or identification of the polypeptide (e.g., poly-His), or to enhance binding of the polypeptide to a solid support. For example, a polypeptide may be conjugated to an immunoglobulin Fc region.

In a related aspect, combination polypeptides are disclosed. A "combination polypeptide" is a polypeptide comprising at least one of the above antigenic portions and one or more additional antigenic *M. tuberculosis* sequences, which are joined via a peptide linkage into a single amino acid chain. The sequences may be joined directly (i.e., with no intervening amino acids) or may be joined by way of a linker sequence (e.g., Gly-Cys-Gly) that does not significantly diminish the antigenic properties of the component polypeptides.

In general, *M. tuberculosis* antigens, and DNA sequences encoding such antigens, may be prepared using any of a variety of procedures. For example, soluble antigens may be isolated from *M. tuberculosis* culture filtrate by procedures known to those of ordinary skill in the art, including anion-exchange and reverse phase chromatography. Purified antigens may then be evaluated for a desired property, such as the ability to react with sera obtained from an *M. tuberculosis*-infected individual. Such screens may be performed using the representative methods described herein. Antigens may then be partially sequenced using, for example, traditional Edman chemistry. See Edman and Berg, *Eur. J. Biochem.* 80:116-132, 1967.

Antigens may also be produced recombinantly using a DNA sequence that encodes the antigen, which has been inserted into an expression vector and expressed in an appropriate host. DNA molecules encoding soluble antigens may be isolated by screening an appropriate *M. tuberculosis* expression library with anti-sera (e.g., rabbit) raised specifically against soluble *M. tuberculosis* antigens. DNA sequences encoding antigens that may or may not be soluble may be identified by screening an appropriate *M. tuberculosis* genomic or cDNA expression library with sera obtained from patients infected with *M. tuberculosis*. Such screens may generally be performed using techniques well known in the art, such as those described in Sambrook et al., *Molecular Cloning: A Laboratory Manual*, Cold Spring Harbor Laboratories, Cold Spring Harbor, NY, 1989.

DNA sequences encoding soluble antigens may also be obtained by screening an appropriate *M. tuberculosis* cDNA or genomic DNA library for DNA sequences that hybridize to degenerate oligonucleotides derived from partial amino acid sequences of isolated soluble antigens. Degenerate oligonucleotide sequences for use in such a screen may
5 be designed and synthesized, and the screen may be performed, as described (for example) in Sambrook et al., *Molecular Cloning: A Laboratory Manual*, Cold Spring Harbor Laboratories, Cold Spring Harbor, NY (and references cited therein). Polymerase chain reaction (PCR) may also be employed, using the above oligonucleotides in methods well known in the art, to isolate a nucleic acid probe from a cDNA or genomic library. The library
10 screen may then be performed using the isolated probe.

Regardless of the method of preparation, the antigens described herein are "antigenic." More specifically, the antigens have the ability to react with sera obtained from an *M. tuberculosis*-infected individual. Reactivity may be evaluated using, for example, the representative ELISA assays described herein, where an absorbance reading with sera from
15 infected individuals that is at least three standard deviations above the absorbance obtained with sera from uninfected individuals is considered positive.

Antigenic portions of *M. tuberculosis* antigens may be prepared and identified using well known techniques, such as those summarized in Paul, *Fundamental Immunology*, 3d ed., Raven Press, 1993, pp. 243-247 and references cited therein. Such techniques include
20 screening polypeptide portions of the native antigen for antigenic properties. The representative ELISAs described herein may generally be employed in these screens. An antigenic portion of a polypeptide is a portion that, within such representative assays, generates a signal in such assays that is substantially similar to that generated by the full length antigen. In other words, an antigenic portion of a *M. tuberculosis* antigen generates at
25 least about 20%, and preferably about 100%, of the signal induced by the full length antigen in a model ELISA as described herein.

Portions and other variants of *M. tuberculosis* antigens may be generated by synthetic or recombinant means. Synthetic polypeptides having fewer than about 100 amino acids, and generally fewer than about 50 amino acids, may be generated using techniques
30 well known in the art. For example, such polypeptides may be synthesized using any of the

commercially available solid-phase techniques, such as the Merrifield solid-phase synthesis method, where amino acids are sequentially added to a growing amino acid chain. See Merrifield, *J. Am. Chem. Soc.* 85:2149-2146, 1963. Equipment for automated synthesis of polypeptides is commercially available from suppliers such as Applied BioSystems, Inc.,
5 Foster City, CA, and may be operated according to the manufacturer's instructions. Variants of a native antigen may generally be prepared using standard mutagenesis techniques, such as oligonucleotide-directed site-specific mutagenesis. Sections of the DNA sequence may also be removed using standard techniques to permit preparation of truncated polypeptides.

Recombinant polypeptides containing portions and/or variants of a native
10 antigen may be readily prepared from a DNA sequence encoding the polypeptide using a variety of techniques well known to those of ordinary skill in the art. For example, supernatants from suitable host/vector systems which secrete recombinant protein into culture media may be first concentrated using a commercially available filter. Following concentration, the concentrate may be applied to a suitable purification matrix such as an
15 affinity matrix or an ion exchange resin. Finally, one or more reverse phase HPLC steps can be employed to further purify a recombinant protein.

Any of a variety of expression vectors known to those of ordinary skill in the art may be employed to express recombinant polypeptides as described herein. Expression may be achieved in any appropriate host cell that has been transformed or transfected with an
20 expression vector containing a DNA molecule that encodes a recombinant polypeptide. Suitable host cells include prokaryotes, yeast and higher eukaryotic cells. Preferably, the host cells employed are *E. coli*, yeast or a mammalian cell line, such as COS or CHO. The DNA sequences expressed in this manner may encode naturally occurring antigens, portions of naturally occurring antigens, or other variants thereof.

25 In general, regardless of the method of preparation, the polypeptides disclosed herein are prepared in substantially pure form. Preferably, the polypeptides are at least about 80% pure, more preferably at least about 90% pure and most preferably at least about 99% pure. For use in the methods described herein, however, such substantially pure polypeptides may be combined.

In certain specific embodiments, the subject invention discloses polypeptides comprising at least an antigenic portion of a soluble *M. tuberculosis* antigen (or a variant of such an antigen), where the antigen has one of the following N-terminal sequences:

- 5 (a) Asp-Pro-Val-Asp-Ala-Val-Ile-Asn-Thr-Thr-Cys-Asn-Tyr-Gly-Gln-Val-Val-Ala-Ala-Leu (SEQ ID NO: 115);
- (b) Ala-Val-Glu-Ser-Gly-Met-Leu-Ala-Leu-Gly-Thr-Pro-Ala-Pro-Ser (SEQ ID NO: 116);
- (c) Ala-Ala-Met-Lys-Pro-Arg-Thr-Gly-Asp-Gly-Pro-Leu-Glu-Ala-Ala-Lys-Glu-Gly-Arg (SEQ ID NO: 117);
- 10 (d) Tyr-Tyr-Trp-Cys-Pro-Gly-Gln-Pro-Phe-Asp-Pro-Ala-Trp-Gly-Pro (SEQ ID NO: 118);
- (e) Asp-Ile-Gly-Ser-Glu-Ser-Thr-Glu-Asp-Gln-Gln-Xaa-Ala-Val (SEQ ID NO: 119);
- (f) Ala-Glu-Glu-Ser-Ile-Ser-Thr-Xaa-Glu-Xaa-Ile-Val-Pro (SEQ ID NO: 120);
- 15 (g) Asp-Pro-Glu-Pro-Ala-Pro-Pro-Val-Pro-Thr-Thr-Ala-Ala-Ser-Pro-Pro-Ser (SEQ ID NO: 121);
- (h) Ala-Pro-Lys-Thr-Tyr-Xaa-Glu-Glu-Leu-Lys-Gly-Thr-Asp-Thr-Gly (SEQ ID NO: 122);
- 20 (i) Asp-Pro-Ala-Ser-Ala-Pro-Asp-Val-Pro-Thr-Ala-Ala-Gln-Gln-Thr-Ser-Leu-Leu-Asn-Ser-Leu-Ala-Asp-Pro-Asn-Val-Ser-Phe-Ala-Asn (SEQ ID NO: 123);
- (j) Xaa-Asp-Ser-Glu-Lys-Ser-Ala-Thr-Ile-Lys-Val-Thr-Asp-Ala-Ser; (SEQ ID NO: 129)
- 25 (k) Ala-Gly-Asp-Thr-Xaa-Ile-Tyr-Ile-Val-Gly-Asn-Leu-Thr-Ala-Asp; (SEQ ID NO: 130) or
- (l) Ala-Pro-Glu-Ser-Gly-Ala-Gly-Leu-Gly-Gly-Thr-Val-Gln-Ala-Gly; (SEQ ID NO: 131)

wherein Xaa may be any amino acid, preferably a cysteine residue. A DNA sequence
30 encoding the antigen identified as (g) above is provided in SEQ ID NO: 52, the deduced

amino acid sequence of which is provided in SEQ ID NO: 53. A DNA sequence encoding the antigen identified as (a) above is provided in SEQ ID NO: 96; its deduced amino acid sequence is provided in SEQ ID NO: 97. A DNA sequence corresponding to antigen (d) above is provided in SEQ ID NO: 24, a DNA sequence corresponding to antigen (c) is provided in SEQ ID NO: 25 and a DNA sequence corresponding to antigen (I) is disclosed in SEQ ID NO: 94 and its deduced amino acid sequence is provided in SEQ ID NO: 95.

In a further specific embodiment, the subject invention discloses polypeptides comprising at least an immunogenic portion of an *M. tuberculosis* antigen having one of the following N-terminal sequences, or a variant thereof that differs only in conservative substitutions and/or modifications:

(m) Xaa-Tyr-Ile-Ala-Tyr-Xaa-Thr-Thr-Ala-Gly-Ile-Val-Pro-Gly-Lys-Ile-Asn-Val-His-Leu-Val; (SEQ ID NO: 132) or

(n) Asp-Pro-Pro-Asp-Pro-His-Gln-Xaa-Asp-Met-Thr-Lys-Gly-Tyr-Tyr-Pro-Gly-Gly-Arg-Arg-Xaa-Phe; (SEQ ID NO: 124)

wherein Xaa may be any amino acid, preferably a cysteine residue.

In other specific embodiments, the subject invention discloses polypeptides comprising at least an antigenic portion of a soluble *M. tuberculosis* antigen (or a variant of such an antigen) that comprises one or more of the amino acid sequences encoded by (a) the DNA sequences of SEQ ID NOS: 1, 2, 4-10, 13-25, 52, 94 and 96, (b) the complements of such DNA sequences, or (c) DNA sequences substantially homologous to a sequence in (a) or (b).

In further specific embodiments, the subject invention discloses polypeptides comprising at least an antigenic portion of a *M. tuberculosis* antigen (or a variant of such an antigen), which may or may not be soluble, that comprises one or more of the amino acid sequences encoded by (a) the DNA sequences of SEQ ID NOS: 26-51, 133, 134, 158-178 and 196, (b) the complements of such DNA sequences or (c) DNA sequences substantially homologous to a sequence in (a) or (b).

In the specific embodiments discussed above, the *M. tuberculosis* antigens include variants that are encoded DNA sequences which are substantially homologous to one

or more of DNA sequences specifically recited herein. "Substantial homology," as used herein, refers to DNA sequences that are capable of hybridizing under moderately stringent conditions. Suitable moderately stringent conditions include prewashing in a solution of 5X SSC, 0.5% SDS, 1.0 mM EDTA (pH 8.0); hybridizing at 50°C-65°C, 5X SSC, overnight or, in the event of cross-species homology, at 45°C with 0.5X SSC; followed by washing twice at 65°C for 20 minutes with each of 2X, 0.5X and 0.2X SSC containing 0.1% SDS). Such hybridizing DNA sequences are also within the scope of this invention, as are nucleotide sequences that, due to code degeneracy, encode an immunogenic polypeptide that is encoded by a hybridizing DNA sequence.

10 In a related aspect, the present invention provides fusion proteins comprising a first and a second inventive polypeptide or, alternatively, a polypeptide of the present invention and a known *M. tuberculosis* antigen, such as the 38 kD antigen described above or ESAT-6 (SEQ ID NOS: 98 and 99), together with variants of such fusion proteins. The fusion proteins of the present invention may also include a linker peptide between the first and second polypeptides.

15 A DNA sequence encoding a fusion protein of the present invention is constructed using known recombinant DNA techniques to assemble separate DNA sequences encoding the first and second polypeptides into an appropriate expression vector. The 3' end of a DNA sequence encoding the first polypeptide is ligated, with or without a peptide linker, to the 5' end of a DNA sequence encoding the second polypeptide so that the reading frames of the sequences are in phase to permit mRNA translation of the two DNA sequences into a single fusion protein that retains the biological activity of both the first and the second polypeptides.

25 A peptide linker sequence may be employed to separate the first and the second polypeptides by a distance sufficient to ensure that each polypeptide folds into its secondary and tertiary structures. Such a peptide linker sequence is incorporated into the fusion protein using standard techniques well known in the art. Suitable peptide linker sequences may be chosen based on the following factors: (1) their ability to adopt a flexible extended conformation; (2) their inability to adopt a secondary structure that could interact with functional epitopes on the first and second polypeptides; and (3) the lack of hydrophobic

30

or charged residues that might react with the polypeptide functional epitopes. Preferred peptide linker sequences contain Gly, Asn and Ser residues. Other near neutral amino acids, such as Thr and Ala may also be used in the linker sequence. Amino acid sequences which may be usefully employed as linkers include those disclosed in Maratea et al., *Gene* 40:39-46, 5 1985; Murphy et al., *Proc. Natl. Acad. Sci. USA* 83:8258-8562, 1986; U.S. Patent No. 4,935,233 and U.S. Patent No. 4,751,180. The linker sequence may be from 1 to about 50 amino acids in length. Peptide linker sequences are not required when the first and second polypeptides have non-essential N-terminal amino acid regions that can be used to separate the functional domains and prevent steric hindrance.

10 In another aspect, the present invention provides methods for using the polypeptides described above to diagnose tuberculosis. In this aspect, methods are provided for detecting *M. tuberculosis* infection in a biological sample, using one or more of the above polypeptides, alone or in combination. In embodiments in which multiple polypeptides are employed, polypeptides other than those specifically described herein, such as the 38 kD 15 antigen described in Andersen and Hansen, *Infect. Immun.* 57:2481-2488, 1989, may be included. As used herein, a "biological sample" is any antibody-containing sample obtained from a patient. Preferably, the sample is whole blood, sputum, serum, plasma, saliva, cerebrospinal fluid or urine. More preferably, the sample is a blood, serum or plasma sample obtained from a patient or a blood supply. The polypeptide(s) are used in an assay, as 20 described below, to determine the presence or absence of antibodies to the polypeptide(s) in the sample, relative to a predetermined cut-off value. The presence of such antibodies indicates previous sensitization to mycobacterial antigens which may be indicative of tuberculosis.

In embodiments in which more than one polypeptide is employed, the 25 polypeptides used are preferably complementary (*i.e.*, one component polypeptide will tend to detect infection in samples where the infection would not be detected by another component polypeptide). Complementary polypeptides may generally be identified by using each polypeptide individually to evaluate serum samples obtained from a series of patients known to be infected with *M. tuberculosis*. After determining which samples test positive (as 30 described below) with each polypeptide, combinations of two or more polypeptides may be

formulated that are capable of detecting infection in most, or all, of the samples tested. Such polypeptides are complementary. For example, approximately 25-30% of sera from tuberculosis-infected individuals are negative for antibodies to any single protein, such as the 38 kD antigen mentioned above. Complementary polypeptides may, therefore, be used in
5 combination with the 38 kD antigen to improve sensitivity of a diagnostic test.

There are a variety of assay formats known to those of ordinary skill in the art for using one or more polypeptides to detect antibodies in a sample. See, e.g., Harlow and Lane, *Antibodies: A Laboratory Manual*, Cold Spring Harbor Laboratory, 1988, which is incorporated herein by reference. In a preferred embodiment, the assay involves the use of
10 polypeptide immobilized on a solid support to bind to and remove the antibody from the sample. The bound antibody may then be detected using a detection reagent that contains a reporter group. Suitable detection reagents include antibodies that bind to the antibody/polypeptide complex and free polypeptide labeled with a reporter group (e.g., in a semi-competitive assay). Alternatively, a competitive assay may be utilized, in which an
15 antibody that binds to the polypeptide is labeled with a reporter group and allowed to bind to the immobilized antigen after incubation of the antigen with the sample. The extent to which components of the sample inhibit the binding of the labeled antibody to the polypeptide is indicative of the reactivity of the sample with the immobilized polypeptide.

The solid support may be any solid material known to those of ordinary skill
20 in the art to which the antigen may be attached. For example, the solid support may be a test well in a microtiter plate or a nitrocellulose or other suitable membrane. Alternatively, the support may be a bead or disc, such as glass, fiberglass, latex or a plastic material such as polystyrene or polyvinylchloride. The support may also be a magnetic particle or a fiber optic sensor, such as those disclosed, for example, in U.S. Patent No. 5,359,681.

25 The polypeptides may be bound to the solid support using a variety of techniques known to those of ordinary skill in the art, which are amply described in the patent and scientific literature. In the context of the present invention, the term "bound" refers to both noncovalent association, such as adsorption, and covalent attachment (which may be a direct linkage between the antigen and functional groups on the support or may be a linkage
30 by way of a cross-linking agent). Binding by adsorption to a well in a microtiter plate or to a

membrane is preferred. In such cases, adsorption may be achieved by contacting the polypeptide, in a suitable buffer, with the solid support for a suitable amount of time. The contact time varies with temperature, but is typically between about 1 hour and 1 day. In general, contacting a well of a plastic microtiter plate (such as polystyrene or polyvinylchloride) with an amount of polypeptide ranging from about 10 ng to about 1 μ g, and preferably about 100 ng, is sufficient to bind an adequate amount of antigen.

Covalent attachment of polypeptide to a solid support may generally be achieved by first reacting the support with a bifunctional reagent that will react with both the support and a functional group, such as a hydroxyl or amino group, on the polypeptide. For example, the polypeptide may be bound to supports having an appropriate polymer coating using benzoquinone or by condensation of an aldehyde group on the support with an amine and an active hydrogen on the polypeptide (*see, e.g.*, Pierce Immunotechnology Catalog and Handbook, 1991, at A12-A13).

In certain embodiments, the assay is an enzyme linked immunosorbent assay (ELISA). This assay may be performed by first contacting a polypeptide antigen that has been immobilized on a solid support, commonly the well of a microtiter plate, with the sample, such that antibodies to the polypeptide within the sample are allowed to bind to the immobilized polypeptide. Unbound sample is then removed from the immobilized polypeptide and a detection reagent capable of binding to the immobilized antibody-polypeptide complex is added. The amount of detection reagent that remains bound to the solid support is then determined using a method appropriate for the specific detection reagent.

More specifically, once the polypeptide is immobilized on the support as described above, the remaining protein binding sites on the support are typically blocked. Any suitable blocking agent known to those of ordinary skill in the art, such as bovine serum albumin or Tween 20TM (Sigma Chemical Co., St. Louis, MO) may be employed. The immobilized polypeptide is then incubated with the sample, and antibody is allowed to bind to the antigen. The sample may be diluted with a suitable diluent, such as phosphate-buffered saline (PBS) prior to incubation. In general, an appropriate contact time (*i.e.*, incubation time) is that period of time that is sufficient to detect the presence of antibody within a *M. tuberculosis*-infected sample. Preferably, the contact time is sufficient to achieve a level

of binding that is at least 95% of that achieved at equilibrium between bound and unbound antibody. Those of ordinary skill in the art will recognize that the time necessary to achieve equilibrium may be readily determined by assaying the level of binding that occurs over a period of time. At room temperature, an incubation time of about 30 minutes is generally
5 sufficient.

Unbound sample may then be removed by washing the solid support with an appropriate buffer, such as PBS containing 0.1% Tween 20™. Detection reagent may then be added to the solid support. An appropriate detection reagent is any compound that binds to the immobilized antibody-polypeptide complex and that can be detected by any of a variety
10 of means known to those in the art. Preferably, the detection reagent contains a binding agent (such as, for example, Protein A, Protein G, immunoglobulin, lectin or free antigen) conjugated to a reporter group. Preferred reporter groups include enzymes (such as horseradish peroxidase), substrates, cofactors, inhibitors, dyes, radionuclides, luminescent groups, fluorescent groups and biotin. The conjugation of binding agent to reporter group
15 may be achieved using standard methods known to those of ordinary skill in the art. Common binding agents may also be purchased conjugated to a variety of reporter groups from many commercial sources (e.g., Zymed Laboratories, San Francisco, CA, and Pierce, Rockford, IL).

The detection reagent is then incubated with the immobilized antibody-polypeptide complex for an amount of time sufficient to detect the bound antibody. An
20 appropriate amount of time may generally be determined from the manufacturer's instructions or by assaying the level of binding that occurs over a period of time. Unbound detection reagent is then removed and bound detection reagent is detected using the reporter group. The method employed for detecting the reporter group depends upon the nature of the
25 reporter group. For radioactive groups, scintillation counting or autoradiographic methods are generally appropriate. Spectroscopic methods may be used to detect dyes, luminescent groups and fluorescent groups. Biotin may be detected using avidin, coupled to a different reporter group (commonly a radioactive or fluorescent group or an enzyme). Enzyme reporter groups may generally be detected by the addition of substrate (generally for a
30 specific period of time), followed by spectroscopic or other analysis of the reaction products.

To determine the presence or absence of anti-*M. tuberculosis* antibodies in the sample, the signal detected from the reporter group that remains bound to the solid support is generally compared to a signal that corresponds to a predetermined cut-off value. In one preferred embodiment, the cut-off value is the average mean signal obtained when the immobilized antigen is incubated with samples from an uninfected patient. In general, a sample generating a signal that is three standard deviations above the predetermined cut-off value is considered positive for tuberculosis. In an alternate preferred embodiment, the cut-off value is determined using a Receiver Operator Curve, according to the method of Sackett et al., *Clinical Epidemiology: A Basic Science for Clinical Medicine*, Little Brown and Co., 1985, pp. 106-107. Briefly, in this embodiment, the cut-off value may be determined from a plot of pairs of true positive rates (*i.e.*, sensitivity) and false positive rates (100%-specificity) that correspond to each possible cut-off value for the diagnostic test result. The cut-off value on the plot that is the closest to the upper left-hand corner (*i.e.*, the value that encloses the largest area) is the most accurate cut-off value, and a sample generating a signal that is higher than the cut-off value determined by this method may be considered positive. Alternatively, the cut-off value may be shifted to the left along the plot, to minimize the false positive rate, or to the right, to minimize the false negative rate. In general, a sample generating a signal that is higher than the cut-off value determined by this method is considered positive for tuberculosis.

In a related embodiment, the assay is performed in a rapid flow-through or strip test format, wherein the antigen is immobilized on a membrane, such as nitrocellulose. In the flow-through test, antibodies within the sample bind to the immobilized polypeptide as the sample passes through the membrane. A detection reagent (*e.g.*, protein A-colloidal gold) then binds to the antibody-polypeptide complex as the solution containing the detection reagent flows through the membrane. The detection of bound detection reagent may then be performed as described above. In the strip test format, one end of the membrane to which polypeptide is bound is immersed in a solution containing the sample. The sample migrates along the membrane through a region containing detection reagent and to the area of immobilized polypeptide. Concentration of detection reagent at the polypeptide indicates the presence of anti-*M. tuberculosis* antibodies in the sample. Typically, the concentration of

detection reagent at that site generates a pattern, such as a line, that can be read visually. The absence of such a pattern indicates a negative result. In general, the amount of polypeptide immobilized on the membrane is selected to generate a visually discernible pattern when the biological sample contains a level of antibodies that would be sufficient to generate a positive
5 signal in an ELISA, as discussed above. Preferably, the amount of polypeptide immobilized on the membrane ranges from about 25 ng to about 1 μ g, and more preferably from about 50 ng to about 500 ng. Such tests can typically be performed with a very small amount (*e.g.*, one drop) of patient serum or blood.

Of course, numerous other assay protocols exist that are suitable for use with
10 the polypeptides of the present invention. The above descriptions are intended to be exemplary only.

In yet another aspect, the present invention provides antibodies to the inventive polypeptides. Antibodies may be prepared by any of a variety of techniques known to those of ordinary skill in the art. *See, e.g.*, Harlow and Lane, *Antibodies: A Laboratory*
15 *Manual*, Cold Spring Harbor Laboratory, 1988. In one such technique, an immunogen comprising the antigenic polypeptide is initially injected into any of a wide variety of mammals (*e.g.*, mice, rats, rabbits, sheep and goats). In this step, the polypeptides of this invention may serve as the immunogen without modification. Alternatively, particularly for relatively short polypeptides, a superior immune response may be elicited if the polypeptide
20 is joined to a carrier protein, such as bovine serum albumin or keyhole limpet hemocyanin. The immunogen is injected into the animal host, preferably according to a predetermined schedule incorporating one or more booster immunizations, and the animals are bled periodically. Polyclonal antibodies specific for the polypeptide may then be purified from such antisera by, for example, affinity chromatography using the polypeptide coupled to a
25 suitable solid support.

Monoclonal antibodies specific for the antigenic polypeptide of interest may be prepared, for example, using the technique of Kohler and Milstein, *Eur. J. Immunol.* 6:511-519, 1976, and improvements thereto. Briefly, these methods involve the preparation of immortal cell lines capable of producing antibodies having the desired specificity (*i.e.*,
30 reactivity with the polypeptide of interest). Such cell lines may be produced, for example,

from spleen cells obtained from an animal immunized as described above. The spleen cells are then immortalized by, for example, fusion with a myeloma cell fusion partner, preferably one that is syngeneic with the immunized animal. A variety of fusion techniques may be employed. For example, the spleen cells and myeloma cells may be combined with a nonionic detergent for a few minutes and then plated at low density on a selective medium that supports the growth of hybrid cells, but not myeloma cells. A preferred selection technique uses HAT (hypoxanthine, aminopterin, thymidine) selection. After a sufficient time, usually about 1 to 2 weeks, colonies of hybrids are observed. Single colonies are selected and tested for binding activity against the polypeptide. Hybridomas having high reactivity and specificity are preferred.

Monoclonal antibodies may be isolated from the supernatants of growing hybridoma colonies. In addition, various techniques may be employed to enhance the yield, such as injection of the hybridoma cell line into the peritoneal cavity of a suitable vertebrate host, such as a mouse. Monoclonal antibodies may then be harvested from the ascites fluid or the blood. Contaminants may be removed from the antibodies by conventional techniques, such as chromatography, gel filtration, precipitation, and extraction. The polypeptides of this invention may be used in the purification process in, for example, an affinity chromatography step.

Antibodies may be used in diagnostic tests to detect the presence of *M. tuberculosis* antigens using assays similar to those detailed above and other techniques well known to those of skill in the art, thereby providing a method for detecting *M. tuberculosis* infection in a patient.

Diagnostic reagents of the present invention may also comprise DNA sequences encoding one or more of the above polypeptides, or one or more portions thereof. For example, at least two oligonucleotide primers may be employed in a polymerase chain reaction (PCR) based assay to amplify *M. tuberculosis*-specific cDNA derived from a biological sample, wherein at least one of the oligonucleotide primers is specific for a DNA molecule encoding a polypeptide of the present invention. The presence of the amplified cDNA is then detected using techniques well known in the art, such as gel electrophoresis. Similarly, oligonucleotide probes specific for a DNA molecule encoding a polypeptide of the

present invention may be used in a hybridization assay to detect the presence of an inventive polypeptide in a biological sample.

As used herein, the term "oligonucleotide primer/probe specific for a DNA molecule" means an oligonucleotide sequence that has at least about 80%, preferably at least about 90% and more preferably at least about 95%, identity to the DNA molecule in question. Oligonucleotide primers and/or probes which may be usefully employed in the inventive diagnostic methods preferably have at least about 10-40 nucleotides. In a preferred embodiment, the oligonucleotide primers comprise at least about 10 contiguous nucleotides of a DNA molecule encoding one of the polypeptides disclosed herein. Preferably, oligonucleotide probes for use in the inventive diagnostic methods comprise at least about 15 contiguous oligonucleotides of a DNA molecule encoding one of the polypeptides disclosed herein. Techniques for both PCR based assays and hybridization assays are well known in the art (see, for example, Mullis *et al. Ibid*; Ehrlich, *Ibid*). Primers or probes may thus be used to detect *M. tuberculosis*-specific sequences in biological samples. DNA probes or primers comprising oligonucleotide sequences described above may be used alone, in combination with each other, or with previously identified sequences, such as the 38 kD antigen discussed above.

The following Examples are offered by way of illustration and not by way of limitation.

EXAMPLES

EXAMPLE 1

PURIFICATION AND CHARACTERIZATION OF POLYPEPTIDES FROM *M. TUBERCULOSIS* CULTURE FILTRATE

This example illustrates the preparation of *M. tuberculosis* soluble polypeptides from culture filtrate. Unless otherwise noted, all percentages in the following example are weight per volume.

M. tuberculosis (either H37Ra, ATCC No. 25177, or H37Rv, ATCC No. 25618) was cultured in sterile GAS media at 37°C for fourteen days. The media was then vacuum filtered (leaving the bulk of the cells) through a 0.45 μ filter into a sterile 2.5 L bottle. The media was then filtered through a 0.2 μ filter into a sterile 4 L bottle. NaN₃ was then added to the culture filtrate to a concentration of 0.04%. The bottles were then placed in a 4°C cold room.

The culture filtrate was concentrated by placing the filtrate in a 12 L reservoir that had been autoclaved and feeding the filtrate into a 400 ml Amicon stir cell which had been rinsed with ethanol and contained a 10,000 kDa MWCO membrane. The pressure was maintained at 60 psi using nitrogen gas. This procedure reduced the 12 L volume to approximately 50 ml.

The culture filtrate was then dialyzed into 0.1% ammonium bicarbonate using a 8,000 kDa MWCO cellulose ester membrane, with two changes of ammonium bicarbonate solution. Protein concentration was then determined by a commercially available BCA assay (Pierce, Rockford, IL).

The dialyzed culture filtrate was then lyophilized, and the polypeptides resuspended in distilled water. The polypeptides were then dialyzed against 0.01 mM 1,3 bis[tris(hydroxymethyl)-methylamino]propane, pH 7.5 (Bis-Tris propane buffer), the initial conditions for anion exchange chromatography. Fractionation was performed using gel perfusion chromatography on a POROS 146 II Q/M anion exchange column 4.6 mm x 100 mm (Perseptive BioSystems, Framingham, MA) equilibrated in 0.01 mM Bis-Tris propane buffer pH 7.5. Polypeptides were eluted with a linear 0-0.5 M NaCl gradient in the above buffer system. The column eluent was monitored at a wavelength of 220 nm.

The pools of polypeptides eluting from the ion exchange column were dialyzed against distilled water and lyophilized. The resulting material was dissolved in 0.1% trifluoroacetic acid (TFA) pH 1.9 in water, and the polypeptides were purified on a Delta-Pak C18 column (Waters, Milford, MA) 300 Angstrom pore size, 5 micron particle size (3.9 x 150 mm). The polypeptides were eluted from the column with a linear gradient from 0-60% dilution buffer (0.1% TFA in acetonitrile). The flow rate was 0.75 ml/minute and the HPLC eluent was monitored at 214 nm. Fractions containing the eluted polypeptides were collected

to maximize the purity of the individual samples. Approximately 200 purified polypeptides were obtained.

The purified polypeptides were then screened for the ability to induce T-cell proliferation in PBMC preparations. The PBMCs from donors known to be PPD skin test positive and whose T cells were shown to proliferate in response to PPD and crude soluble proteins from MTB were cultured in medium comprising RPMI 1640 supplemented with 10% pooled human serum and 50 µg/ml gentamicin. Purified polypeptides were added in duplicate at concentrations of 0.5 to 10 µg/mL. After six days of culture in 96-well round-bottom plates in a volume of 200 µl, 50 µl of medium was removed from each well for determination of IFN-γ levels, as described below. The plates were then pulsed with 1 µCi/well of tritiated thymidine for a further 18 hours, harvested and tritium uptake determined using a gas scintillation counter. Fractions that resulted in proliferation in both replicates three fold greater than the proliferation observed in cells cultured in medium alone were considered positive.

IFN-γ was measured using an enzyme-linked immunosorbent assay (ELISA). ELISA plates were coated with a mouse monoclonal antibody directed to human IFN-γ (Chemicon) in PBS for four hours at room temperature. Wells were then blocked with PBS containing 5% (W/V) non-fat dried milk for 1 hour at room temperature. The plates were then washed six times in PBS/0.2% TWEEN-20 and samples diluted 1:2 in culture medium in the ELISA plates were incubated overnight at room temperature. The plates were again washed and a polyclonal rabbit anti-human IFN-γ serum diluted 1:3000 in PBS/10% normal goat serum was added to each well. The plates were then incubated for two hours at room temperature, washed and horseradish peroxidase-coupled anti-rabbit IgG (Jackson Labs.) was added at a 1:2000 dilution in PBS/5% non-fat dried milk. After a further two hour incubation at room temperature, the plates were washed and TMB substrate added. The reaction was stopped after 20 min with 1 N sulfuric acid. Optical density was determined at 450 nm using 570 nm as a reference wavelength. Fractions that resulted in both replicates giving an OD two fold greater than the mean OD from cells cultured in medium alone, plus 3 standard deviations, were considered positive.

For sequencing, the polypeptides were individually dried onto Biobrene™ (Perkin Elmer/Applied BioSystems Division, Foster City, CA) treated glass fiber filters. The filters with polypeptide were loaded onto a Perkin Elmer/Applied BioSystems Division Procise 492 protein sequencer. The polypeptides were sequenced from the amino
5 terminal and using traditional Edman chemistry. The amino acid sequence was determined for each polypeptide by comparing the retention time of the PTH amino acid derivative to the appropriate PTH derivative standards.

Using the procedure described above, antigens having the following N-terminal sequences were isolated:

- 10 (a) Asp-Pro-Val-Asp-Ala-Val-Ile-Asn-Thr-Thr-Xaa-Asn-Tyr-Gly-Gln-Val-Val-Ala-Ala-Leu (SEQ ID NO: 54);
- (b) Ala-Val-Glu-Ser-Gly-Met-Leu-Ala-Leu-Gly-Thr-Pro-Ala-Pro-Ser (SEQ ID NO: 55);
- (c) Ala-Ala-Met-Lys-Pro-Arg-Thr-Gly-Asp-Gly-Pro-Leu-Glu-Ala-Ala-
15 Lys-Glu-Gly-Arg (SEQ ID NO: 56);
- (d) Tyr-Tyr-Trp-Cys-Pro-Gly-Gln-Pro-Phe-Asp-Pro-Ala-Trp-Gly-Pro (SEQ ID NO: 57);
- (e) Asp-Ile-Gly-Ser-Glu-Ser-Thr-Glu-Asp-Gln-Gln-Xaa-Ala-Val (SEQ ID NO: 58);
- 20 (f) Ala-Glu-Glu-Ser-Ile-Ser-Thr-Xaa-Glu-Xaa-Ile-Val-Pro (SEQ ID NO: 59);
- (g) Asp-Pro-Glu-Pro-Ala-Pro-Pro-Val-Pro-Thr-Ala-Ala-Ala-Ala-Pro-Pro-Ala (SEQ ID NO: 60); and
- (h) Ala-Pro-Lys-Thr-Tyr-Xaa-Glu-Glu-Leu-Lys-Gly-Thr-Asp-Thr-Gly
25 (SEQ ID NO: 61);

wherein Xaa may be any amino acid.

An additional antigen was isolated employing a microbore HPLC purification step in addition to the procedure described above. Specifically, 20 µl of a fraction comprising a mixture of antigens from the chromatographic purification step previously described, was
30 purified on an Aquapore C18 column (Perkin Elmer/Applied Biosystems Division, Foster

City, CA) with a 7 micron pore size, column size 1 mm x 100 mm, in a Perkin Elmer/Applied Biosystems Division Model 172 HPLC. Fractions were eluted from the column with a linear gradient of 1%/minute of acetonitrile (containing 0.05% TFA) in water (0.05% TFA) at a flow rate of 80 µl/minute. The eluent was monitored at 250 nm. The original fraction was
5 separated into 4 major peaks plus other smaller components and a polypeptide was obtained which was shown to have a molecular weight of 12.054 Kd (by mass spectrometry) and the following N-terminal sequence:

(i) Asp-Pro-Ala-Ser-Ala-Pro-Asp-Val-Pro-Thr-Ala-Ala-Gln-Gln-Thr-Ser-
Leu-Leu-Asn-Asn-Leu-Ala-Asp-Pro-Asp-Val-Ser-Phe-Ala-Asp (SEQ
10 ID NO: 62).

This polypeptide was shown to induce proliferation and IFN-γ production in PBMC preparations using the assays described above.

Additional soluble antigens were isolated from *M. tuberculosis* culture filtrate as follows. *M. tuberculosis* culture filtrate was prepared as described above. Following
15 dialysis against Bis-Tris propane buffer, at pH 5.5, fractionation was performed using anion exchange chromatography on a Poros QE column 4.6 x 100 mm (Perseptive Biosystems) equilibrated in Bis-Tris propane buffer pH 5.5. Polypeptides were eluted with a linear 0-1.5 M NaCl gradient in the above buffer system at a flow rate of 10 ml/min. The column eluent was monitored at a wavelength of 214 nm.

20 The fractions eluting from the ion exchange column were pooled and subjected to reverse phase chromatography using a Poros R2 column 4.6 x 100 mm (Perseptive Biosystems). Polypeptides were eluted from the column with a linear gradient from 0-100% acetonitrile (0.1% TFA) at a flow rate of 5 ml/min. The eluent was monitored at 214 nm.

25 Fractions containing the eluted polypeptides were lyophilized and resuspended in 80 µl of aqueous 0.1% TFA and further subjected to reverse phase chromatography on a Vydac C4 column 4.6 x 150 mm (Western Analytical, Temecula, CA) with a linear gradient of 0-100% acetonitrile (0.1% TFA) at a flow rate of 2 ml/min. Eluent was monitored at 214 nm.

The fraction with biological activity was separated into one major peak plus other smaller components. Western blot of this peak onto PVDF membrane revealed three major bands of molecular weights 14 Kd, 20 Kd and 26 Kd. These polypeptides were determined to have the following N-terminal sequences, respectively:

- 5 (j) Xaa-Asp-Ser-Glu-Lys-Ser-Ala-Thr-Ile-Lys-Val-Thr-Asp-Ala-Ser;
(SEQ ID NO: 129)
- (k) Ala-Gly-Asp-Thr-Xaa-Ile-Tyr-Ile-Val-Gly-Asn-Leu-Thr-Ala-Asp;
(SEQ ID NO: 130) and
- 10 (l) Ala-Pro-Glu-Ser-Gly-Ala-Gly-Leu-Gly-Gly-Thr-Val-Gln-Ala-Gly;
(SEQ ID NO: 131), wherein Xaa may be any amino acid.

Using the assays described above, these polypeptides were shown to induce proliferation and IFN- γ production in PBMC preparations. Figs. 1A and B show the results of such assays using PBMC preparations from a first and a second donor, respectively.

- DNA sequences that encode the antigens designated as (a), (c), (d) and (g)
- 15 above were obtained by screening a *M. tuberculosis* genomic library using ^{32}P end labeled degenerate oligonucleotides corresponding to the N-terminal sequence and containing *M. tuberculosis* codon bias. The screen performed using a probe corresponding to antigen (a) above identified a clone having the sequence provided in SEQ ID NO: 96. The polypeptide encoded by SEQ ID NO: 96 is provided in SEQ ID NO: 97. The screen performed using a
- 20 probe corresponding to antigen (g) above identified a clone having the sequence provided in SEQ ID NO: 52. The polypeptide encoded by SEQ ID NO: 52 is provided in SEQ ID NO: 53. The screen performed using a probe corresponding to antigen (d) above identified a clone having the sequence provided in SEQ ID NO: 24, and the screen performed with a probe corresponding to antigen (c) identified a clone having the sequence provided in SEQ ID
- 25 NO: 25.

- The above amino acid sequences were compared to known amino acid sequences in the gene bank using the DNA STAR system. The database searched contains some 173,000 proteins and is a combination of the Swiss, PIR databases along with translated protein sequences (Version 87). No significant homologies to the amino acid sequences for
- 30 antigens (a)-(h) and (l) were detected.

The amino acid sequence for antigen (i) was found to be homologous to a sequence from *M. leprae*. The full length *M. leprae* sequence was amplified from genomic DNA using the sequence obtained from GENBANK. This sequence was then used to screen an *M. tuberculosis* library and a full length copy of the *M. tuberculosis* homologue was
5 obtained (SEQ ID NO: 94).

The amino acid sequence for antigen (j) was found to be homologous to a known *M. tuberculosis* protein translated from a DNA sequence. To the best of the inventors' knowledge, this protein has not been previously shown to possess T-cell stimulatory activity. The amino acid sequence for antigen (k) was found to be related to a
10 sequence from *M. leprae*.

In the proliferation and IFN- γ assays described above, using three PPD positive donors, the results for representative antigens provided above are presented in Table
1:

15

TABLE 1RESULTS OF PBMC PROLIFERATION AND IFN- γ ASSAYS

Sequence	Proliferation	IFN- γ
(a)	+	-
(c)	+++	+++
(d)	++	++
(g)	+++	+++
(h)	+++	+++

In Table 1, responses that gave a stimulation index (SI) of between 2 and 4
20 (compared to cells cultured in medium alone) were scored as +, as SI of 4-8 or 2-4 at a concentration of 1 μ g or less was scored as ++ and an SI of greater than 8 was scored as +++. The antigen of sequence (i) was found to have a high SI (+++) for one donor and lower SI (++ and +) for the two other donors in both proliferation and IFN- γ assays. These results

indicate that these antigens are capable of inducing proliferation and/or interferon- γ production.

EXAMPLE 2

5 USE OF PATIENT SERA TO ISOLATE *M. TUBERCULOSIS* ANTIGENS

This example illustrates the isolation of antigens from *M. tuberculosis* lysate by screening with serum from *M. tuberculosis*-infected individuals.

Dessicated *M. tuberculosis* H37Ra (Difco Laboratories) was added to a 2% NP40 solution, and alternately homogenized and sonicated three times. The resulting suspension was centrifuged at 13,000 rpm in microfuge tubes and the supernatant put through a 0.2 micron syringe filter. The filtrate was bound to Macro Prep DEAE beads (BioRad, Hercules, CA). The beads were extensively washed with 20 mM Tris pH 7.5 and bound proteins eluted with 1M NaCl. The NaCl elute was dialyzed overnight against 10 mM Tris, pH 7.5. Dialyzed solution was treated with DNase and RNase at 0.05 mg/ml for 30 min. at room temperature and then with α -D-mannosidase, 0.5 U/mg at pH 4.5 for 3-4 hours at room temperature. After returning to pH 7.5, the material was fractionated via FPLC over a Bio Scale-Q-20 column (BioRad). Fractions were combined into nine pools, concentrated in a Centriprep 10 (Amicon, Beverley, MA) and screened by Western blot for serological activity using a serum pool from *M. tuberculosis*-infected patients which was not immunoreactive with other antigens of the present invention.

The most reactive fraction was run in SDS-PAGE and transferred to PVDF. A band at approximately 85 Kd was cut out yielding the sequence:

(m) Xaa-Tyr-Ile-Ala-Tyr-Xaa-Thr-Thr-Ala-Gly-Ile-Val-Pro-Gly-Lys-Ile-Asn-Val-His-Leu-Val; (SEQ ID NO: 132), wherein Xaa may be any amino acid.

Comparison of this sequence with those in the gene bank as described above, revealed no significant homologies to known sequences.

A DNA sequence that encodes the antigen designated as (m) above was obtained by screening a genomic *M. tuberculosis* Erdman strain library using labeled

degenerate oligonucleotides corresponding to the N-terminal sequence of SEQ ID NO:137. A clone was identified having the DNA sequence provided in SEQ ID NO: 198. This sequence was found to encode the amino acid sequence provided in SEQ ID NO: 199. Comparison of these sequences with those in the genebank revealed some similarity to sequences previously identified in *M. tuberculosis* and *M. bovis*.

EXAMPLE 3

PREPARATION OF DNA SEQUENCES ENCODING *M. TUBERCULOSIS* ANTIGENS

This example illustrates the preparation of DNA sequences encoding *M. tuberculosis* antigens by screening a *M. tuberculosis* expression library with sera obtained from patients infected with *M. tuberculosis*, or with anti-sera raised against *M. tuberculosis* antigens.

A. PREPARATION OF *M. TUBERCULOSIS* SOLUBLE ANTIGENS USING RABBIT ANTI-SERA RAISED AGAINST *M. TUBERCULOSIS* SUPERNATANT

Genomic DNA was isolated from the *M. tuberculosis* strain H37Ra. The DNA was randomly sheared and used to construct an expression library using the Lambda ZAP expression system (Stratagene, La Jolla, CA). Rabbit anti-sera was generated against secretory proteins of the *M. tuberculosis* strains H37Ra, H37Rv and Erdman by immunizing a rabbit with concentrated supernatant of the *M. tuberculosis* cultures. Specifically, the rabbit was first immunized subcutaneously with 200 µg of protein antigen in a total volume of 2 ml containing 100 µg muramyl dipeptide (Calbiochem, La Jolla, CA) and 1 ml of incomplete Freund's adjuvant. Four weeks later the rabbit was boosted subcutaneously with 100 µg antigen in incomplete Freund's adjuvant. Finally, the rabbit was immunized intravenously four weeks later with 50 µg protein antigen. The anti-sera were used to screen the expression library as described in Sambrook et al., *Molecular Cloning: A Laboratory Manual*, Cold Spring Harbor Laboratories, Cold Spring Harbor, NY, 1989. Bacteriophage plaques expressing immunoreactive antigens were purified. Phagemid from the plaques was rescued and the nucleotide sequences of the *M. tuberculosis* clones deduced.

Thirty two clones were purified. Of these, 25 represent sequences that have not been previously identified in *M. tuberculosis*. Proteins were induced by IPTG and purified by gel elution, as described in Skeiky et al., *J. Exp. Med.* 181:1527-1537, 1995. Representative partial sequences of DNA molecules identified in this screen are provided in
5 SEQ ID NOS: 1-25. The corresponding predicted amino acid sequences are shown in SEQ ID NOS: 64-88.

On comparison of these sequences with known sequences in the gene bank using the databases described above, it was found that the clones referred to hereinafter as TbRA2A, TbRA16, TbRA18, and TbRA29 (SEQ ID NOS: 77, 69, 71, 76) show some
10 homology to sequences previously identified in *Mycobacterium leprae* but not in *M. tuberculosis*. TbRA11, TbRA26, TbRA28 and TbDPEP (SEQ ID NOS: 66, 74, 75, 53) have been previously identified in *M. tuberculosis*. No significant homologies were found to TbRA1, TbRA3, TbRA4, TbRA9, TbRA10, TbRA13, TbRA17, TbRA19, TbRA29, TbRA32, TbRA36 and the overlapping clones TbRA35 and TbRA12 (SEQ ID NOS: 64, 78,
15 82, 83, 65, 68, 76, 72, 76, 79, 81, 80, 67, respectively). The clone TbRa24 is overlapping with clone TbRa29.

B. USE OF SERA FROM PATIENTS HAVING PULMONARY OR PLEURAL TUBERCULOSIS TO IDENTIFY DNA SEQUENCES ENCODING *M. TUBERCULOSIS* ANTIGENS

20 The genomic DNA library described above, and an additional H37Rv library, were screened using pools of sera obtained from patients with active tuberculosis. To prepare the H37Rv library, *M. tuberculosis* strain H37Rv genomic DNA was isolated, subjected to partial Sau3A digestion and used to construct an expression library using the Lambda Zap expression system (Stratagene, La Jolla, Ca). Three different pools of sera, each containing
25 sera obtained from three individuals with active pulmonary or pleural disease, were used in the expression screening. The pools were designated TbL, TbM and TbH, referring to relative reactivity with H37Ra lysate (i.e., TbL = low reactivity, TbM = medium reactivity and TbH = high reactivity) in both ELISA and immunoblot format. A fourth pool of sera from seven patients with active pulmonary tuberculosis was also employed. All of the sera

lacked increased reactivity with the recombinant 38 kD *M. tuberculosis* H37Ra phosphate-binding protein.

All pools were pre-adsorbed with *E. coli* lysate and used to screen the H37Ra and H37Rv expression libraries, as described in Sambrook et al., *Molecular Cloning: A Laboratory Manual*, Cold Spring Harbor Laboratories, Cold Spring Harbor, NY, 1989. Bacteriophage plaques expressing immunoreactive antigens were purified. Phagemid from the plaques was rescued and the nucleotide sequences of the *M. tuberculosis* clones deduced.

Thirty two clones were purified. Of these, 31 represented sequences that had not been previously identified in human *M. tuberculosis*. Representative sequences of the DNA molecules identified are provided in SEQ ID NOS.: 26-51 and 100. Of these, TbH-8-2 (SEQ. ID NO. 100) is a partial clone of TbH-8, and TbH-4 (SEQ. ID NO. 43) and TbH-4-FWD (SEQ. ID NO. 44) are non-contiguous sequences from the same clone. Amino acid sequences for the antigens hereinafter identified as Tb38-1, TbH-4, TbH-8, TbH-9, and TbH-12 are shown in SEQ ID NOS.: 89-93. Comparison of these sequences with known sequences in the gene bank using the databases identified above revealed no significant homologies to TbH-4, TbH-8, TbH-9 and TbM-3, although weak homologies were found to TbH-9. TbH-12 was found to be homologous to a 34 kD antigenic protein previously identified in *M. paratuberculosis* (Acc. No. S28515). Tb38-1 was found to be located 34 base pairs upstream of the open reading frame for the antigen ESAT-6 previously identified in *M. bovis* (Acc. No. U34848) and in *M. tuberculosis* (Sorensen et al., *Infect. Immun.* 63:1710-1717, 1995).

Probes derived from Tb38-1 and TbH-9, both isolated from an H37Ra library, were used to identify clones in an H37Rv library. Tb38-1 hybridized to Tb38-1F2, Tb38-1F3, Tb38-1F5 and Tb38-1F6 (SEQ. ID NOS: 107, 108, 111, 113, and 114). (SEQ ID NOS: 107 and 108 are non-contiguous sequences from clone Tb38-1F2.) Two open reading frames were deduced in Tb38-1F2; one corresponds to Tb37FL (SEQ. ID. NO. 109), the second, a partial sequence, may be the homologue of Tb38-1 and is called Tb38-IN (SEQ. ID NO. 110). The deduced amino acid sequence of Tb38-1F3 is presented in SEQ. ID. NO. 112. A TbH-9 probe identified three clones in the H37Rv library: TbH-9-FL (SEQ. ID NO. 101), which may be the homologue of TbH-9 (R37Ra), TbH-9-1 (SEQ. ID NO. 103), and TbH-8-2 (SEQ.

ID NO. 105) is a partial clone of TbH-8. The deduced amino acid sequences for these three clones are presented in SEQ ID NOS: 102, 104 and 106.

Further screening of the *M. tuberculosis* genomic DNA library, as described above, resulted in the recovery of ten additional reactive clones, representing seven different
5 genes. One of these genes was identified as the 38 Kd antigen discussed above, one was determined to be identical to the 14Kd alpha crystallin heat shock protein previously shown to be present in *M. tuberculosis*, and a third was determined to be identical to the antigen TbH-8 described above. The determined DNA sequences for the remaining five clones (hereinafter referred to as TbH-29, TbH-30, TbH-32 and TbH-33) are provided in SEQ ID
10 NO: 133-136, respectively, with the corresponding predicted amino acid sequences being provided in SEQ ID NO: 137-140, respectively. The DNA and amino acid sequences for these antigens were compared with those in the gene bank as described above. No homologies were found to the 5' end of TbH-29 (which contains the reactive open reading frame), although the 3' end of TbH-29 was found to be identical to the *M. tuberculosis*
15 cosmid Y227. TbH-32 and TbH-33 were found to be identical to the previously identified *M. tuberculosis* insertion element IS6110 and to the *M. tuberculosis* cosmid Y50, respectively. No significant homologies to TbH-30 were found.

Positive phagemid from this additional screening were used to infect *E. coli* XL-1 Blue MRF', as described in Sambrook et al., *supra*. Induction of recombinant protein
20 was accomplished by the addition of IPTG. Induced and uninduced lysates were run in duplicate on SDS-PAGE and transferred to nitrocellulose filters. Filters were reacted with human *M. tuberculosis* sera (1:200 dilution) reactive with TbH and a rabbit sera (1:200 or 1:250 dilution) reactive with the N-terminal 4 Kd portion of lacZ. Sera incubations were performed for 2 hours at room temperature. Bound antibody was detected by addition of ¹²⁵I-
25 labeled Protein A and subsequent exposure to film for variable times ranging from 16 hours to 11 days. The results of the immunoblots are summarized in Table 2.

TABLE 2

<u>Antigen</u>	Human <i>M. tb</i> <u>Sera</u>	Anti-lacZ <u>Sera</u>
TbH-29	45 Kd	45 Kd
TbH-30	No reactivity	29 Kd
TbH-32	12 Kd	12 Kd
TbH-33	16 Kd	16 Kd

Positive reaction of the recombinant human *M. tuberculosis* antigens with both the human *M. tuberculosis* sera and anti-lacZ sera indicate that reactivity of the human *M. tuberculosis* sera is directed towards the fusion protein. Antigens reactive with the anti-lacZ sera but not with the human *M. tuberculosis* sera may be the result of the human *M. tuberculosis* sera recognizing conformational epitopes, or the antigen-antibody binding kinetics may be such that the 2 hour sera exposure in the immunoblot is not sufficient.

Studies were undertaken to determine whether the antigens TbH-9 and Tb38-1 represent cellular proteins or are secreted into *M. tuberculosis* culture media. In the first study, rabbit sera were raised against A) secretory proteins of *M. tuberculosis*, B) the known secretory recombinant *M. tuberculosis* antigen 85b, C) recombinant Tb38-1 and D) recombinant TbH-9, using protocols substantially as described in Example 3A. Total *M. tuberculosis* lysate, concentrated supernatant of *M. tuberculosis* cultures and the recombinant antigens 85b, TbH-9 and Tb38-1 were resolved on denaturing gels, immobilized on nitrocellulose membranes and duplicate blots were probed using the rabbit sera described above.

The results of this analysis using control sera (panel I) and antisera (panel II) against secretory proteins, recombinant 85b, recombinant Tb38-1 and recombinant TbH-9 are shown in Figures 2A-D, respectively, wherein the lane designations are as follows: 1) molecular weight protein standards; 2) 5 µg of *M. tuberculosis* lysate; 3) 5 µg secretory proteins; 4) 50 ng recombinant Tb38-1; 5) 50 ng recombinant TbH-9; and 6) 50 ng recombinant 85b. The recombinant antigens were engineered with six terminal histidine

residues and would therefore be expected to migrate with a mobility approximately 1 kD larger than the native protein. In Figure 2D, recombinant TbH-9 is lacking approximately 10 kD of the full-length 42 kD antigen, hence the significant difference in the size of the immunoreactive native TbH-9 antigen in the lysate lane (indicated by an arrow). These results demonstrate that Tb38-1 and TbH-9 are intracellular antigens and are not actively secreted by *M. tuberculosis*.

The finding that TbH-9 is an intracellular antigen was confirmed by determining the reactivity of TbH-9-specific human T cell clones to recombinant TbH-9, secretory *M. tuberculosis* proteins and PPD. A TbH-9-specific T cell clone (designated 131TbH-9) was generated from PBMC of a healthy PPD-positive donor. The proliferative response of 131TbH-9 to secretory proteins, recombinant TbH-9 and a control *M. tuberculosis* antigen, TbRa11, was determined by measuring uptake of tritiated thymidine, as described in Example 1. As shown in Figure 3A, the clone 131TbH-9 responds specifically to TbH-9, showing that TbH-9 is not a significant component of *M. tuberculosis* secretory proteins. Figure 3B shows the production of IFN- γ by a second TbH-9-specific T cell clone (designated PPD 800-10) prepared from PBMC from a healthy PPD-positive donor, following stimulation of the T cell clone with secretory proteins, PPD or recombinant TbH-9. These results further confirm that TbH-9 is not secreted by *M. tuberculosis*.

C. USE OF SERA FROM PATIENTS HAVING EXTRAPULMONARY TUBERCULOSIS TO IDENTIFY DNA SEQUENCES ENCODING *M. TUBERCULOSIS* ANTIGENS

Genomic DNA was isolated from *M. tuberculosis* Erdman strain, randomly sheared and used to construct an expression library employing the Lambda ZAP expression system (Stratagene, La Jolla, CA). The resulting library was screened using pools of sera obtained from individuals with extrapulmonary tuberculosis, as described above in Example 3B, with the secondary antibody being goat anti-human IgG + A + M (H+L) conjugated with alkaline phosphatase.

Eighteen clones were purified. Of these, 4 clones (hereinafter referred to as XP14, XP24, XP31 and XP32) were found to bear some similarity to known sequences. The determined DNA sequences for XP14, XP24 and XP31 are provided in SEQ ID NOS: 151-

153, respectively, with the 5' and 3' DNA sequences for XP32 being provided in SEQ ID NOS: 154 and 155, respectively. The predicted amino acid sequence for XP14 is provided in SEQ ID NO: 156. The reverse complement of XP14 was found to encode the amino acid sequence provided in SEQ ID NO: 157.

5 Comparison of the sequences for the remaining 14 clones (hereinafter referred to as XP1-XP6, XP17-XP19, XP22, XP25, XP27, XP30 and XP36) with those in the genebank as described above, revealed no homologies with the exception of the 3' ends of XP2 and XP6 which were found to bear some homology to known *M. tuberculosis* cosmids. The DNA sequences for XP27 and XP36 are shown in SEQ ID NOS: 158 and 159, 10 respectively, with the 5' sequences for XP4, XP5, XP17 and XP30 being shown in SEQ ID NOS: 160-163, respectively, and the 5' and 3' sequences for XP2, XP3, XP6, XP18, XP19, XP22 and XP25 being shown in SEQ ID NOS: 164 and 165; 166 and 167; 168 and 169; 170 and 171; 172 and 173; 174 and 175; and 176 and 177, respectively. XP1 was found to overlap with the DNA sequences for TbH4, disclosed above. The full-length DNA sequence 15 for TbH4-XP1 is provided in SEQ ID NO: 178. This DNA sequence was found to contain an open reading frame encoding the amino acid sequence shown in SEQ ID NO: 179. The reverse complement of TbH4-XP1 was found to contain an open reading frame encoding the amino acid sequence shown in SEQ ID NO: 180. The DNA sequence for XP36 was found to contain two open reading frames encoding the amino acid sequence shown in SEQ ID NOS: 20 181 and 182, with the reverse complement containing an open reading frame encoding the amino acid sequence shown in SEQ ID NO: 183.

Recombinant XP1 protein was prepared as described above in Example 3B, with a metal ion affinity chromatography column being employed for purification. Recombinant XP1 was found to stimulate cell proliferation and IFN- γ production in T cells 25 isolated from an *M. tuberculosis*-immune donors.

D. PREPARATION OF *M. TUBERCULOSIS* SOLUBLE ANTIGENS USING RABBIT ANTI-SERA
RAISED AGAINST *M. TUBERCULOSIS* FRACTIONATED PROTEINS

M. tuberculosis lysate was prepared as described above in Example 2. The 30 resulting material was fractionated by HPLC and the fractions screened by Western blot for

serological activity with a serum pool from *M. tuberculosis*-infected patients which showed little or no immunoreactivity with other antigens of the present invention. Rabbit anti-sera was generated against the most reactive fraction using the method described in Example 3A. The anti-sera was used to screen an *M. tuberculosis* Erdman strain genomic DNA expression library prepared as described above. Bacteriophage plaques expressing immunoreactive antigens were purified. Phagemid from the plaques was rescued and the nucleotide sequences of the *M. tuberculosis* clones determined.

Ten different clones were purified. Of these, one was found to be TbRa35, described above, and one was found to be the previously identified *M. tuberculosis* antigen, HSP60. Of the remaining eight clones, six (hereinafter referred to as RDIF2, RDIF5, RDIF8, RDIF10, RDIF11 and RDIF12) were found to bear some similarity to previously identified *M. tuberculosis* sequences. The determined DNA sequences for RDIF2, RDIF5, RDIF8, RDIF10 and RDIF11 are provided in SEQ ID NOS: 184-188, respectively, with the corresponding predicted amino acid sequences being provided in SEQ ID NOS: 189-193, respectively. The 5' and 3' DNA sequences for RDIF12 are provided in SEQ ID NOS: 194 and 195, respectively. No significant homologies were found to the antigen RDIF-7. The determined DNA and predicted amino acid sequences for RDIF7 are provided in SEQ ID NOS: 196 and 197, respectively. One additional clone, referred to as RDIF6 was isolated, however, this was found to be identical to RDIF5.

Recombinant RDIF6, RDIF8, RDIF10 and RDIF11 were prepared as described above. These antigens were found to stimulate cell proliferation and IFN- γ production in T cells isolated from *M. tuberculosis*-immune donors.

EXAMPLE 4

PURIFICATION AND CHARACTERIZATION OF A POLYPEPTIDE FROM TUBERCULIN PURIFIED PROTEIN DERIVATIVE

An *M. tuberculosis* polypeptide was isolated from tuberculin purified protein derivative (PPD) as follows.

PPD was prepared as published with some modification (Seibert, F. et al.,
Tuberculin purified protein derivative. Preparation and analyses of a large quantity for
standard. The American Review of Tuberculosis 44:9-25, 1941). *M. tuberculosis* Rv strain
was grown for 6 weeks in synthetic medium in roller bottles at 37°C. Bottles containing the
5 bacterial growth were then heated to 100°C in water vapor for 3 hours. Cultures were sterile
filtered using a 0.22 µ filter and the liquid phase was concentrated 20 times using a 3 kD cut-
off membrane. Proteins were precipitated once with 50% ammonium sulfate solution and
eight times with 25% ammonium sulfate solution. The resulting proteins (PPD) were
fractionated by reverse phase liquid chromatography (RP-HPLC) using a C18 column (7.8 x
10 300 mM; Waters, Milford, MA) in a Biocad HPLC system (Perceptive Biosystems,
Framingham, MA). Fractions were eluted from the column with a linear gradient from 0-
100% buffer (0.1% TFA in acetonitrile). The flow rate was 10 ml/minute and eluent was
monitored at 214 nm and 280 nm.

Six fractions were collected, dried, suspended in PBS and tested individually
15 in *M. tuberculosis*-infected guinea pigs for induction of delayed type hypersensitivity (DTH)
reaction. One fraction was found to induce a strong DTH reaction and was subsequently
fractionated further by RP-HPLC on a microbore Vydac C18 column (Cat. No. 218TP5115)
in a Perkin Elmer/Applied Biosystems Division Model 172 HPLC. Fractions were eluted
with a linear gradient from 5-100% buffer (0.05% TFA in acetonitrile) with a flow rate of 80
20 µl/minute. Eluent was monitored at 215 nm. Eight fractions were collected and tested for
induction of DTH in *M. tuberculosis*-infected guinea pigs. One fraction was found to induce
strong DTH of about 16 mm induration. The other fractions did not induce detectable DTH.
The positive fraction was submitted to SDS-PAGE gel electrophoresis and found to contain a
single protein band of approximately 12 kD molecular weight.

25 This polypeptide, herein after referred to as DPPD, was sequenced from the
amino terminal using a Perkin Elmer/Applied Biosystems Division Procise 492 protein
sequencer as described above and found to have the N-terminal sequence shown in SEQ ID
NO.: 124. Comparison of this sequence with known sequences in the gene bank as described
above revealed no known homologies. Four cyanogen bromide fragments of DPPD were
30 isolated and found to have the sequences shown in SEQ ID NOS: 125-128.

EXAMPLE 5

SYNTHESIS OF SYNTHETIC POLYPEPTIDES

5 Polypeptides may be synthesized on a Millipore 9050 peptide synthesizer using Fmoc chemistry with HPTU (O-Benzotriazole-N,N,N',N'-tetramethyluronium hexafluorophosphate) activation. A Gly-Cys-Gly sequence may be attached to the amino terminus of the peptide to provide a method of conjugation or labeling of the peptide. Cleavage of the peptides from the solid support may be carried out using the following
10 cleavage mixture: trifluoroacetic acid:ethanedithiol:thioanisole:water:phenol (40:1:2:2:3). After cleaving for 2 hours, the peptides may be precipitated in cold methyl-t-butyl-ether. The peptide pellets may then be dissolved in water containing 0.1% trifluoroacetic acid (TFA) and lyophilized prior to purification by C18 reverse phase HPLC. A gradient of 0-60% acetonitrile (containing 0.1% TFA) in water (containing 0.1% TFA) may be used to elute the
15 peptides. Following lyophilization of the pure fractions, the peptides may be characterized using electrospray mass spectrometry and by amino acid analysis.

This procedure was used to synthesize a TbM-1 peptide that contains one and a half repeats of a TbM-1 sequence. The TbM-1 peptide has the sequence GCGDRSGGNLDQIRLRDRSGGNL (SEQ ID NO: 63).

20

EXAMPLE 6

USE OF REPRESENTATIVE ANTIGENS FOR SERODIAGNOSIS OF TUBERCULOSIS

25 This Example illustrates the diagnostic properties of several representative antigens.

Assays were performed in 96-well plates were coated with 200 ng antigen diluted to 50 μ L in carbonate coating buffer, pH 9.6. The wells were coated overnight at 4°C (or 2 hours at 37°C). The plate contents were then removed and the wells were blocked for 2
30 hours with 200 μ L of PBS/1% BSA. After the blocking step, the wells were washed five

times with PBS/0.1% Tween 20™. 50 µL sera, diluted 1:100 in PBS/0.1% Tween 20™/0.1% BSA, was then added to each well and incubated for 30 minutes at room temperature. The plates were then washed again five times with PBS/0.1% Tween 20™.

The enzyme conjugate (horseradish peroxidase - Protein A, Zymed, San Francisco, CA) was then diluted 1:10,000 in PBS/0.1% Tween 20™/0.1% BSA, and 50 µL of the diluted conjugate was added to each well and incubated for 30 minutes at room temperature. Following incubation, the wells were washed five times with PBS/0.1% Tween 20™. 100 µL of tetramethylbenzidine peroxidase (TMB) substrate (Kirkegaard and Perry Laboratories, Gaithersburg, MD) was added, undiluted, and incubated for about 15 minutes. The reaction was stopped with the addition of 100 µL of 1 N H₂SO₄ to each well, and the plates were read at 450 nm.

Figure 4 shows the ELISA reactivity of two recombinant antigens isolated using method A in Example 3 (TbRa3 and TbRa9) with sera from *M. tuberculosis* positive and negative patients. The reactivity of these antigens is compared to that of bacterial lysate isolated from *M. tuberculosis* strain H37Ra (Difco, Detroit, MI). In both cases, the recombinant antigens differentiated positive from negative sera. Based on cut-off values obtained from receiver-operator curves, TbRa3 detected 56 out of 87 positive sera, and TbRa9 detected 111 out of 165 positive sera.

Figure 5 illustrates the ELISA reactivity of representative antigens isolated using method B of Example 3. The reactivity of the recombinant antigens TbH4, TbH12, Tb38-1 and the peptide TbM-1 (as described in Example 4) is compared to that of the 38 kD antigen described by Andersen and Hansen, *Infect. Immun.* 57:2481-2488, 1989. Again, all of the polypeptides tested differentiated positive from negative sera. Based on cut-off values obtained from receiver-operator curves, TbH4 detected 67 out of 126 positive sera, TbH12 detected 50 out of 125 positive sera, 38-1 detected 61 out of 101 positive sera and the TbM-1 peptide detected 25 out of 30 positive sera.

The reactivity of four antigens (TbRa3, TbRa9, TbH4 and TbH12) with sera from a group of *M. tuberculosis* infected patients with differing reactivity in the acid fast stain of sputum (Smithwick and David, *Tubercle* 52:226, 1971) was also examined, and compared

to the reactivity of *M. tuberculosis* lysate and the 38 kD antigen. The results are presented in Table 3, below:

TABLE 3

5 REACTIVITY OF ANTIGENS WITH SERA FROM *M. TUBERCULOSIS* PATIENTS

Patient	Acid Fast Sputum	ELISA Values					
		Lysate	38kD	TbRa9	TbH12	TbH4	TbRa3
Tb01B93I-2	++++	1.853	0.634	0.998	1.022	1.030	1.314
Tb01B93I-19	++++	2.657	2.322	0.608	0.837	1.857	2.335
Tb01B93I-8	+++	2.703	0.527	0.492	0.281	0.501	2.002
Tb01B93I-10	+++	1.665	1.301	0.685	0.216	0.448	0.458
Tb01B93I-11	+++	2.817	0.697	0.509	0.301	0.173	2.608
Tb01B93I-15	+++	1.28	0.283	0.808	0.218	1.537	0.811
Tb01B93I-16	+++	2.908	>3	0.899	0.441	0.593	1.080
Tb01B93I-25	+++	0.395	0.131	0.335	0.211	0.107	0.948
Tb01B93I-87	+++	2.653	2.432	2.282	0.977	1.221	0.857
Tb01B93I-89	+++	1.912	2.370	2.436	0.876	0.520	0.952
Tb01B94I-108	+++	1.639	0.341	0.797	0.368	0.654	0.798
Tb01B94I-201	+++	1.721	0.419	0.661	0.137	0.064	0.692
Tb01B93I-88	++	1.939	1.269	2.519	1.381	0.214	0.530
Tb01B93I-92	++	2.355	2.329	2.78	0.685	0.997	2.527
Tb01B94I-109	++	0.993	0.620	0.574	0.441	0.5	2.558
Tb01B94I-210	++	2.777	>3	0.393	0.367	1.004	1.315
Tb01B94I-224	++	2.913	0.476	0.251	1.297	1.990	0.256

Patient	Acid Fast Sputum	ELISA Values					
		Lysate	38kD	TbRa9	TbH12	TbH4	TbRa3
Tb01B93I-9	+	2.649	0.278	0.210	0.140	0.181	1.586
Tb01B93I-14	+	>3	1.538	0.282	0.291	0.549	2.880
Tb01B93I-21	+	2.645	0.739	2.499	0.783	0.536	1.770
Tb01B93I-22	+	0.714	0.451	2.082	0.285	0.269	1.159
Tb01B93I-31	+	0.956	0.490	1.019	0.812	0.176	1.293
Tb01B93I-32	—	2.261	0.786	0.668	0.273	0.535	0.405
Tb01B93I-52	—	0.658	0.114	0.434	0.330	0.273	1.140
Tb01B93I-99	—	2.118	0.584	1.62	0.119	0.977	0.729
Tb01B94I-130	—	1.349	0.224	0.86	0.282	0.383	2.146
Tb01B94I-131	—	0.685	0.324	1.173	0.059	0.118	1.431
AT4-0070	Normal	0.072	0.043	0.092	0.071	0.040	0.039
AT4-0105	Normal	0.397	0.121	0.118	0.103	0.078	0.390
3/15/94-1	Normal	0.227	0.064	0.098	0.026	0.001	0.228
4/15/93-2	Normal	0.114	0.240	0.071	0.034	0.041	0.264
5/26/94-4	Normal	0.089	0.259	0.096	0.046	0.008	0.053
5/26/94-3	Normal	0.139	0.093	0.085	0.019	0.067	0.01

Based on cut-off values obtained from receiver-operator curves, TbRa3 detected 23 out of 27 positive sera, TbRa9 detected 22 out of 27, TbH4 detected 18 out of 27 and TbH12 detected 15 out of 27. If used in combination, these four antigens would have a theoretical sensitivity of 27 out of 27, indicating that these antigens should complement each other in the serological detection of *M. tuberculosis* infection. In addition, several of the recombinant antigens detected positive sera that were not detected using the 38 kD antigen, indicating that these antigens may be complementary to the 38 kD antigen.

The reactivity of the recombinant antigen TbRa11 with sera from *M. tuberculosis* patients shown to be negative for the 38 kD antigen, as well as with sera from PPD positive and normal donors, was determined by ELISA as described above. The results are shown in Figure 6 which indicates that TbRa11, while being negative with sera from PPD positive and normal donors, detected sera that were negative with the 38 kD antigen. Of the thirteen 38 kD negative sera tested, nine were positive with TbRa11, indicating that this antigen may be reacting with a sub-group of 38 kD antigen negative sera. In contrast, in a group of 38 kD positive sera where TbRa11 was reactive, the mean OD 450 for TbRa11 was lower than that for the 38 kD antigen. The data indicate an inverse relationship between the presence of TbRa11 activity and 38 kD positivity.

The antigen TbRa2A was tested in an indirect ELISA using initially 50 µl of serum at 1:100 dilution for 30 minutes at room temperature followed by washing in PBS Tween and incubating for 30 minutes with biotinylated Protein A (Zymed, San Francisco, CA) at a 1:10,000 dilution. Following washing, 50 µl of streptavidin-horseradish peroxidase (Zymed) at 1:10,000 dilution was added and the mixture incubated for 30 minutes. After washing, the assay was developed with TMB substrate as described above. The reactivity of TbRa2A with sera from *M. tuberculosis* patients and normal donors is shown in Table 4. The mean value for reactivity of TbRa2A with sera from *M. tuberculosis* patients was 0.444 with a standard deviation of 0.309. The mean for reactivity with sera from normal donors was 0.109 with a standard deviation of 0.029. Testing of 38 kD negative sera (Figure 7) also indicated that the TbRa2A antigen was capable of detecting sera in this category.

TABLE 4

REACTIVITY OF TBRA2A WITH SERA FROM *M. TUBERCULOSIS* PATIENTS AND FROM NORMAL

DONORS

Serum ID	Status	OD 450
Tb85	TB	0.680
Tb86	TB	0.450
Tb87	TB	0.263
Tb88	TB	0.275
Tb89	TB	0.403

Tb91	TB	0.393
Tb92	TB	0.401
Tb93	TB	0.232
Tb94	TB	0.333
Tb95	TB	0.435
Tb96	TB	0.284
Tb97	TB	0.320
Tb99	TB	0.328
Tb100	TB	0.817
Tb101	TB	0.607
Tb102	TB	0.191
Tb103	TB	0.228
Tb107	TB	0.324
Tb109	TB	1.572
Tb112	TB	0.338
DL4-0176	Normal	0.036
AT4-0043	Normal	0.126
AT4-0044	Normal	0.130
AT4-0052	Normal	0.135
AT4-0053	Normal	0.133
AT4-0062	Normal	0.128
AT4-0070	Normal	0.088
AT4-0091	Normal	0.108
AT4-0100	Normal	0.106
AT4-0105	Normal	0.108
AT4-0109	Normal	0.105

The reactivity of the recombinant antigen (g) (SEQ ID NO: 60) with sera from *M. tuberculosis* patients and normal donors was determined by ELISA as described above. Figure 8 shows the results of the titration of antigen (g) with four *M. tuberculosis* positive sera that were all reactive with the 38 kD antigen and with four donor sera. All four positive sera were reactive with antigen (g).

The reactivity of the recombinant antigen TbH-29 (SEQ ID NO: 137) with sera from *M. tuberculosis* patients, PPD positive donors and normal donors was determined by indirect ELISA as described above. The results are shown in Figure 9. TbH-29 detected 30 out of 60 *M. tuberculosis* sera, 2 out of 8 PPD positive sera and 2 out of 27 normal sera.

Figure 10 shows the results of ELISA tests (both direct and indirect) of the antigen TbH-33 (SEQ ID NO: 140) with sera from *M. tuberculosis* patients and from normal

donors and with a pool of sera from *M. tuberculosis* patients. The mean OD 450 was demonstrated to be higher with sera from *M. tuberculosis* patients than from normal donors, with the mean OD 450 being significantly higher in the indirect ELISA than in the direct ELISA. Figure 11 is a titration curve for the reactivity of recombinant TbH-33 with sera from *M. tuberculosis* patients and from normal donors showing an increase in OD 450 with increasing concentration of antigen.

The reactivity of the recombinant antigens RDIF6, RDIF8 and RDIF10 (SEQ ID NOS: 184-187, respectively) with sera from *M. tuberculosis* patients and normal donors was determined by ELISA as described above. RDIF6 detected 6 out of 32 *M. tuberculosis* sera and 0 out of 15 normal sera; RDIF8 detected 14 out of 32 *M. tuberculosis* sera and 0 out of 15 normal sera; and RDIF10 detected 4 out of 27 *M. tuberculosis* sera and 1 out of 15 normal sera. In addition, RDIF10 was found to detect 0 out of 5 sera from PPD-positive donors.

EXAMPLE 7

PREPARATION AND CHARACTERIZATION OF *M. TUBERCULOSIS* FUSION PROTEINS

A fusion protein containing TbRa3, the 38 kD antigen and Tb38-1 was prepared as follows.

Each of the DNA constructs TbRa3, 38 kD and Tb38-1 were modified by PCR in order to facilitate their fusion and the subsequent expression of the fusion protein TbRa3-38 kD-Tb38-1. TbRa3, 38 kD and Tb38-1 DNA was used to perform PCR using the primers PDM-64 and PDM-65 (SEQ ID NO: 141 and 142), PDM-57 and PDM-58 (SEQ ID NO: 143 and 144), and PDM-69 and PDM-60 (SEQ ID NO: 145-146), respectively. In each case, the DNA amplification was performed using 10 μ l 10X Pfu buffer, 2 μ l 10 mM dNTPs, 2 μ l each of the PCR primers at 10 μ M concentration, 81.5 μ l water, 1.5 μ l Pfu DNA polymerase (Stratagene, La Jolla, CA) and 1 μ l DNA at either 70 ng/ μ l (for TbRa3) or 50 ng/ μ l (for 38 kD and Tb38-1). For TbRa3, denaturation at 94°C was performed for 2 min, followed by 40 cycles of 96°C for 15 sec and 72°C for 1 min, and lastly by 72°C for 4 min. For 38 kD, denaturation at 96°C was performed for 2 min, followed by 40 cycles of 96°C for 30 sec,

68°C for 15 sec and 72°C for 3 min, and finally by 72°C for 4 min. For Tb38-1 denaturation at 94°C for 2 min was followed by 10 cycles of 96°C for 15 sec, 68°C for 15 sec and 72°C for 1.5 min, 30 cycles of 96°C for 15 sec, 64°C for 15 sec and 72°C for 1.5, and finally by 72°C for 4 min.

5 The TbRa3 PCR fragment was digested with NdeI and EcoRI and cloned directly into pT7⁺L2 IL 1 vector using NdeI and EcoRI sites. The 38 kD PCR fragment was digested with Sse8387I, treated with T4 DNA polymerase to make blunt ends and then digested with EcoRI for direct cloning into the pT7⁺L2Ra3-1 vector which was digested with StuI and EcoRI. The 38-1 PCR fragment was digested with Eco47III and EcoRI and directly
10 subcloned into pT7⁺L2Ra3/38kD-17 digested with the same enzymes. The whole fusion was then transferred to pET28b using NdeI and EcoRI sites. The fusion construct was confirmed by DNA sequencing.

 The expression construct was transformed to BLR pLys S *E. coli* (Novagen, Madison, WI) and grown overnight in LB broth with kanamycin (30 µg/ml) and
15 chloramphenicol (34 µg/ml). This culture (12 ml) was used to inoculate 500 ml 2XYT with the same antibiotics and the culture was induced with IPTG at an OD₅₆₀ of 0.44 to a final concentration of 1.2 mM. Four hours post-induction, the bacteria were harvested and sonicated in 20 mM Tris (8.0), 100 mM NaCl, 0.1% DOC, 20 µg/ml Leupeptin, 20 mM PMSF followed by centrifugation at 26,000 X g. The resulting pellet was resuspended in 8 M
20 urea, 20 mM Tris (8.0), 100 mM NaCl and bound to Pro-bond nickel resin (Invitrogen, Carlsbad, CA). The column was washed several times with the above buffer then eluted with an imidazole gradient (50 mM, 100 mM, 500 mM imidazole was added to 8 M urea, 20 mM Tris (8.0), 100 mM NaCl). The eluates containing the protein of interest were then dialyzed against 10 mM Tris (8.0).

25 The DNA and amino acid sequences for the resulting fusion protein (hereinafter referred to as TbRa3-38 kD-Tb38-1) are provided in SEQ ID NO: 147 and 148, respectively.

 A fusion protein containing the two antigens TbH-9 and Tb38-1 (hereinafter referred to as TbH9-Tb38-1) without a hinge sequence, was prepared using a similar

procedure to that described above. The DNA sequence for the TbH9-Tb38-1 fusion protein is provided in SEQ ID NO: 151.

A fusion protein containing TbRa3, the antigen 38kD, Tb38-1 and DPEP was prepared as follows.

5 Each of the DNA constructs TbRa3, 38 kD and Tb38-1 were modified by PCR and cloned into vectors essentially as described above, with the primers PDM-69 (SEQ ID NO:145 and PDM-83 (SEQ ID NO: 200) being used for amplification of the Tb38-1A fragment. Tb38-1A differs from Tb38-1 by a DraI site at the 3' end of the coding region that keeps the final amino acid intact while creating a blunt restriction site that is in frame. The
10 TbRa3/38kD/Tb38-1A fusion was then transferred to pET28b using NdeI and EcoRI sites.

DPEP DNA was used to perform PCR using the primers PDM-84 and PDM-85 (SEQ ID NO: 201 and 202, respectively) and 1 µl DNA at 50 ng/µl. Denaturation at 94 °C was performed for 2 min, followed by 10 cycles of 96 °C for 15 sec, 68 °C for 15 sec and 72 °C for 1.5 min; 30 cycles of 96 °C for 15 sec, 64 °C for 15 sec and 72 °C for 1.5 min; and
15 finally by 72 °C for 4 min. The DPEP PCR fragment was digested with EcoRI and Eco72I and clones directly into the pET28Ra3/38kD/38-1A construct which was digested with DraI and EcoRI. The fusion construct was confirmed to be correct by DNA sequencing. Recombinant protein was prepared as described above. The DNA and amino acid sequences for the resulting fusion protein (hereinafter referred to as TbF-2) are provided in SEQ ID NO:
20 203 and 204, respectively.

EXAMPLE 8

USE OF *M. TUBERCULOSIS* FUSION PROTEINS FOR SERODIAGNOSIS OF TUBERCULOSIS

25

The effectiveness of the fusion protein TbRa3-38 kD-Tb38-1, prepared as described above, in the serodiagnosis of tuberculosis infection was examined by ELISA.

The ELISA protocol was as described above in Example 6, with the fusion protein being coated at 200 ng/well. A panel of sera was chosen from a group of tuberculosis
30 patients previously shown, either by ELISA or by western blot analysis, to react with each of

the three antigens individually or in combination. Such a panel enabled the dissection of the serological reactivity of the fusion protein to determine if all three epitopes functioned with the fusion protein. As shown in Table 5, all four sera that reacted with TbRa3 only were detectable with the fusion protein. Three sera that reacted only with Tb38-1 were also detectable, as were two sera that reacted with 38 kD alone. The remaining 15 sera were all positive with the fusion protein based on a cut-off in the assay of mean negatives +3 standard deviations. This data demonstrates the functional activity of all three epitopes in the fusion protein.

10

TABLE 5
REACTIVITY OF TRI-PEPTIDE FUSION PROTEIN WITH SERA FROM *M. TUBERCULOSIS* PATIENTS

Serum ID	Status	ELISA and/or Western Blot Reactivity with Individual proteins			Fusion recombinant OD 450	Fusion Recombinant Status
		38kd	Tb38-1	TbRa3		
01B93I-40	TB	-	-	+	0.413	+
01B93I-41	TB	-	+	+	0.392	+
01B93I-29	TB	+	-	+	2.217	+
01B93I-109	TB	+	±	+	0.522	+
01B93I-132	TB	+	+	+	0.937	+
5004	TB	±	+	±	1.098	+
15004	TB	+	+	+	2.077	+
39004	TB	+	+	+	1.675	+
68004	TB	+	+	+	2.388	+
99004	TB	-	+	±	0.607	+
107004	TB	-	+	±	0.667	+
92004	TB	+	±	±	1.070	+
97004	TB	+	-	±	1.152	+
118004	TB	+	-	±	2.694	+
173004	TB	+	+	+	3.258	+
175004	TB	+	-	+	2.514	+
274004	TB	-	-	+	3.220	+
276004	TB	-	+	-	2.991	+
282004	TB	+	-	-	0.824	+

289004	TB	-	-	+	0.848	+
308004	TB	-	+	-	3.338	+
314004	TB	-	+	-	1.362	+
317004	TB	+	-	-	0.763	+
312004	TB	-	-	+	1.079	+
D176	PPD	-	-	-	0.145	-
D162	PPD	-	-	-	0.073	-
D161	PPD	-	-	-	0.097	-
D27	PPD	-	-	-	0.082	-
A6-124	NORMAL	-	-	-	0.053	-
A6-125	NORMAL	-	-	-	0.087	-
A6-126	NORMAL	-	-	-	0.346	±
A6-127	NORMAL	-	-	-	0.064	-
A6-128	NORMAL	-	-	-	0.034	-
A6-129	NORMAL	-	-	-	0.037	-
A6-130	NORMAL	-	-	-	0.057	-
A6-131	NORMAL	-	-	-	0.054	-
A6-132	NORMAL	-	-	-	0.022	-
A6-133	NORMAL	-	-	-	0.147	-
A6-134	NORMAL	-	-	-	0.101	-
A6-135	NORMAL	-	-	-	0.066	-
A6-136	NORMAL	-	-	-	0.054	-
A6-137	NORMAL	-	-	-	0.065	-
A6-138	NORMAL	-	-	-	0.041	-
A6-139	NORMAL	-	-	-	0.103	-
A6-140	NORMAL	-	-	-	0.212	-
A6-141	NORMAL	-	-	-	0.056	-
A6-142	NORMAL	-	-	-	0.051	-

The reactivity of the fusion protein TbF-2 with sera from *M. tuberculosis*-infected patients was examined by ELISA using the protocol described above. The results of these studies (Table 6) demonstrate that all four antigens function independently in the fusion

5 protein.

TABLE 6
REACTIVITY OF TbF-2 FUSION PROTEIN WITH TB AND NORMAL SERA

Serum ID	Status	TbF OD450	Status	TbF-2 OD450	Status	ELISA Reactivity			
						38 kD	TbRa3	Tb38-1	DPEP
B931-40	TB	0.57	+	0.321	+	-	+	-	+
B931-41	TB	0.601	+	0.396	+	+	+	+	-
B931-109	TB	0.494	+	0.404	+	+	+	±	-
B931-132	TB	1.502	+	1.292	+	+	+	+	±
5004	TB	1.806	+	1.666	+	±	±	+	-
15004	TB	2.862	+	2.468	+	+	+	+	-
39004	TB	2.443	+	1.722	+	+	+	+	-
68004	TB	2.871	+	2.575	+	+	+	+	-
99004	TB	0.691	+	0.971	+	-	±	+	-
107004	TB	0.875	+	0.732	+	-	±	+	-
92004	TB	1.632	+	1.394	+	+	±	±	-
97004	TB	1.491	+	1.979	+	+	±	-	+
118004	TB	3.182	+	3.045	+	+	±	-	-
173004	TB	3.644	+	3.578	+	+	+	+	-
175004	TB	3.332	+	2.916	+	+	+	-	-
274004	TB	3.696	+	3.716	+	-	+	-	+
276004	TB	3.243	+	2.56	+	-	-	+	-
282004	TB	1.249	+	1.234	+	+	-	-	-
289004	TB	1.373	+	1.17	+	-	+	-	-
308004	TB	3.708	+	3.355	+	-	-	+	-
314004	TB	1.663	+	1.399	+	-	-	+	-
317004	TB	1.163	+	0.92	+	+	-	-	-
312004	TB	1.709	+	1.453	+	-	+	-	-
380004	TB	0.238	-	0.461	+	-	±	-	+
451004	TB	0.18	-	0.2	-	-	-	-	±
478004	TB	0.188	-	0.469	+	-	-	-	±
410004	TB	0.384	+	2.392	+	±	-	-	+
411004	TB	0.306	+	0.874	+	-	+	-	+
421004	TB	0.357	+	1.456	+	-	+	-	+
528004	TB	0.047	-	0.196	-	-	-	-	+
A6-87	Normal	0.094	-	0.063	-	-	-	-	-
A6-88	Normal	0.214	-	0.19	-	-	-	-	-
A6-89	Normal	0.248	-	0.125	-	-	-	-	-
A6-90	Normal	0.179	-	0.206	-	-	-	-	-
A6-91	Normal	0.135	-	0.151	-	-	-	-	-
A6-92	Normal	0.064	-	0.097	-	-	-	-	-
A6-93	Normal	0.072	-	0.098	-	-	-	-	-
A6-94	Normal	0.072	-	0.064	-	-	-	-	-
A6-95	Normal	0.125	-	0.159	-	-	-	-	-
A6-96	Normal	0.121	-	0.12	-	-	-	-	-
Cut-off		0.284		0.266					

One of skill in the art will appreciate that the order of the individual antigens within the fusion protein may be changed and that comparable activity would be expected provided each of the epitopes is still functionally available. In addition, truncated forms of the proteins containing active epitopes may be used in the construction of fusion proteins.

5

From the foregoing, it will be appreciated that, although specific embodiments of the invention have been described herein for the purpose of illustration, various modifications may be made without deviating from the spirit and scope of the invention.

SEQUENCE LISTING

(1) GENERAL INFORMATION:

- (i) APPLICANTS: Reed, Steven G.
Skeiky, Yasir A.W.
Dillon, Davin C.
Campos-Neto, Antonia
Houghton, Raymond
Vedvick, Thomas S.
Twardzik, Daniel R.
Lodes, Michael J.
- (ii) TITLE OF INVENTION: COMPOUNDS AND METHODS FOR DIAGNOSIS OF
TUBERCULOSIS
- (iii) NUMBER OF SEQUENCES: 209
- (iv) CORRESPONDENCE ADDRESS:
 - (A) ADDRESSEE: SEED and BERRY LLP
 - (B) STREET: 6300 Columbia Center, 701 Fifth Avenue
 - (C) CITY: Seattle
 - (D) STATE: Washington
 - (E) COUNTRY: USA
 - (F) ZIP: 98104-7092
- (v) COMPUTER READABLE FORM:
 - (A) MEDIUM TYPE: Floppy disk
 - (B) COMPUTER: IBM PC compatible
 - (C) OPERATING SYSTEM: PC-DOS/MS-DOS
 - (D) SOFTWARE: PatentIn Release #1.0, Version #1.30
- (vi) CURRENT APPLICATION DATA:
 - (A) APPLICATION NUMBER:
 - (B) FILING DATE: 01-OCT-1997
 - (C) CLASSIFICATION:
- (viii) ATTORNEY/AGENT INFORMATION:
 - (A) NAME: Maki, David J.
 - (B) REGISTRATION NUMBER: 31,392
 - (C) REFERENCE/DOCKET NUMBER: 210121.417C7
- (ix) TELECOMMUNICATION INFORMATION:
 - (A) TELEPHONE: (206) 622-4900
 - (B) TELEFAX: (206) 682-6031

(2) INFORMATION FOR SEQ ID NO:1:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 766 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:1:

CGAGGCACCG GTAGTTTGAA CCAAACGCAC AATCGACGGG CAAACGAACG GAAGAACACA	60
ACCATGAAGA TGGTGAAATC GATCGCCGCA GGTCTGACCG CCGCGGCTGC AATCGGCGCC	120
GCTGCGGCCG GTGTGACTTC GATCATGGCT GGCGGCCCGG TCGTATACCA GATGCAGCCG	180
GTCGTCTTCG GCGCGCCACT GCCGTTGGAC CCGGCATCCG CCCCTGACGT CCCGACCGCC	240
GCCCAGTTGA CCAGCCTGCT CAACAGCCTC GCCGATCCCA ACGTGTCTGTT TGCGAACAAG	300
GGCAGTCTGG TCGAGGGCGG CATCGGGGGC ACCGAGGCGC GCATCGCCGA CCACAAGCTG	360
AAGAAGGCCG CCGAGCACGG GGATCTGCCG CTGTCTGTTCA GCGTGACGAA CATCCAGCCG	420
GCGGCCGCCG GTTCGGCCAC CGCCGACGTT TCCGTCTCGG GTCCGAAGCT CTCGTGCGCG	480
GTCACGCAGA ACGTCACGTT CGTGAATCAA GGCGGCTGGA TGCTGTCACG CGCATCGGCG	540
ATGGAGTTGC TGCAGGCCGC AGGGNAACTG ATTGGCGGGC CGGNTTCAGC CCGCTGTTCA	600
GCTACGCCGC CCGCCTGGTG ACGCGTCCAT GTCGAACACT CGCGCGTGTA GCACGGTGCG	660
GTNTGCGCAG GGNCGCACGC ACCGCCCGGT GCAAGCCGTC CTCGAGATAG GTGGTGNCTC	720
GNCACCAGNG ANCACCCCN NNTCGNCNNT TCTCGNTGNT GNATGA	766

(2) INFORMATION FOR SEQ ID NO:2:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 752 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:2:

ATGCATCACC ATCACCATCA CGATGAAGTC ACGGTAGAGA CGACCTCCGT CTTCCGCGCA	60
GACTTCCTCA GCGAGCTGGA CGCTCCTGCG CAAGCGGGTA CGGAGAGCGC GGTCTCCGGG	120
GTGGAAGGGC TCCCGCCGGG CTCGGCGTTG CTGGTAGTCA AACGAGGCCC CAACGCCGGG	180
TCCCGGTTCC TACTCGACCA AGCCATCACG TCGGCTGGTC GGCATCCCGA CAGCGACATA	240

TTTCTCGACG ACGTGACCGT GAGCCGTCGC CATGCTGAAT TCCGGTTGGA AAACAACGAA	300
TTCAATGTCTG TCGATGTCTGG GAGTCTCAAC GGCACCTACG TCAACCGCGA GCCCGTGGAT	360
TCGGCGGTGC TGGCGAACGG CGACGAGGTC CAGATCGGCA AGCTCCGGTT GGTGTTCTTG	420
ACCGGACCCA AGCAAGGCGA GGATGACGGG AGTACCGGGG GCCCGTGAGC GCACCCGATA	480
GCCCCGCGCT GGCCGGGATG TCGATCGGGG CGGTCTCCG ACCTGCTACG ACCGGATTTT	540
CCCTGATGTC CACCATCTCC AAGATTCTGAT TCTTGGGAGG CTTGAGGGTC NGGGTGACCC	600
CCCCGCGGGC CTCATTCTGG GGTNTCGGCN GGTTCACCC CNTACCNACT GCCNCCCGN	660
TTGCNAATTC NTTCTTCNCT GCCCNAAAG GGACNNTAN CTTGCCGCTN GAAANGGTNA	720
TCCNGGGCCC NTCCTNGAAN CCCNTCCCC CT	752

(2) INFORMATION FOR SEQ ID NO:3:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 813 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:3:

CATATGCATC ACCATCACCA TCACACTTCT AACCGCCCAG CGCGTCGGGG GCGTCGAGCA	60
CCACGCGACA CCGGGCCCGA TCGATCTGCT AGCTTGAGTC TGGTCAGGCA TCGTCGTCAG	120
CAGCGCGATG CCCTATGTTT GTCGTCGACT CAGATATCGC GGCAATCCAA TCTCCCGCCT	180
GCGGCCGGCG GTGCTGCAAA CTACTCCCGG AGGAATTTCTG ACGTGCGCAT CAAGATCTTC	240
ATGCTGGTCA CGGCTGTCGT TTTGCTCTGT TGTTGCGGTG TGGCCACGGC CGCGCCCAAG	300
ACCTACTGCG AGGAGTTGAA AGGCACCGAT ACCGGCCAGG CGTGCCAGAT TCAAATGTCC	360
GACCCGGCCT ACAACATCAA CATCAGCCTG CCCAGTTACT ACCCCGACCA GAAGTCGCTG	420
GAAAATTACA TCGCCCAGAC GCGCGACAAG TTCCTCAGCG CGGCCACATC GTCCACTCCA	480
CGCGAAGCCC CCTACGAATT GAATATCACC TCGGCCACAT ACCAGTCCGC GATACCGCCG	540
CGTGGTACGC AGGCCGTGGT GCTCAMGGTC TACCACAACG CCGGCGGCAC GCACCCAACG	600
ACCACGTACA AGGCCTTCGA TTGGGACCAG GCCTATCGCA AGCCAATCAC CTATGACACG	660
CTGTGGCAGG CTGACACCGA TCCGCTGCCA GTCGTCTTCC CCATTGTTGC AAGGTGAACT	720

GAGCAACGCA GACCGGGACA ACWGGTATCG ATAGCCGCCN AATGCCGGCT TGGAACCCNG 780
TGAAATTATC ACAACTTCGC AGTCACNAAA NAA 813

(2) INFORMATION FOR SEQ ID NO:4:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 447 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:4:

CGGTATGAAC ACGGCCGCGT CCGATAACTT CCAGCTGTCC CAGGGTGGGC AGGGATTTCGC 60
CATTCCGATC GGGCAGGCGA TGGCGATCGC GGGCCAGATC CGATCGGGTG GGGGGTCACC 120
CACCGTTCAT ATCGGGCCTA CCGCCTTCCT CGGCTTGGGT GTTGTCGACA ACAACGGCAA 180
CGGCGCACGA GTCCAACGCG TGGTCGGGAG CGCTCCGGCG GCAAGTCTCG GCATCTCCAC 240
CGGCGACGTG ATCACC GCGG TCGACGGCGC TCCGATCAAC TCGGCCACCG CGATGGCGGA 300
CGCGCTTAAC GGGCATCATC CCGGTGACGT CATCTCGGTG AACTGGCAAA CCAAGTCGGG 360
CGGCACGCGT ACAGGGAACG TGACATTGGC CGAGGGACCC CCGGCCTGAT TTCGTCGYGG 420
ATACCACCCG CCGGCCGGCC AATTGGA 447

(2) INFORMATION FOR SEQ ID NO:5:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 604 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:5:

GTCCCACTGC GGTCCGCGAG TATGTCGCCC AGCAAATGTC TGGCAGCCGC CCAACGGAAT 60
CCGGTGATCC GACGTCGAG GTTGTCGAAC CCGCCGCCGC GGAAGTATCG GTCCATGCCT 120
AGCCCGGCGA CGGCGAGCGC CGGAATGGCG CGAGTGAGGA GGCGGGCAAT TTGGCGGGGC 180

CCGGCGACGG NGAGCGCCGG AATGGCGCGA GTGAGGAGGT GGNCAGTCAT GCCCAGNGTG 240
 ATCCAATCAA CCTGNATTCG GNCTGNNGGN CCATTTGACA ATCGAGGTAG TGAGCGCAAA 300
 TGAATGATGG AAAACGGGNG GNGACGTCCG NTGTTCTGGT GGTGNTAGGT GNCTGNCTGG 360
 NGTNGNGGNT ATCAGGATGT TCTTCGNCGA AANCTGATGN CGAGGAACAG GGTGTNCCCG 420
 NNANCCNAN GGNGTCCNAN CCCNNNTCC TCGNCGANAT CANANAGNCG NTTGATGNGA 480
 NAAAAGGGTG GANCAGNNNN AANTNGNGGN CCNAANAANC NNNANNGNNG NNAGNTNGNT 540
 NNTNTTNNC ANNNNNNTG NNGNNGNCCN NNNCAANCNN NTNNNGNAA NNGGNTTNTT 600
 NAAT 604

(2) INFORMATION FOR SEQ ID NO:6:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 633 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:6:

TTGCANGTCG AACCACCTCA CTAAAGGGAA CAAAAGCTNG AGCTCCACCG CGGTGGCGGC 60
 CGCTCTAGAA CTAGTGKATM YYYCKGGCTG CAGSAATYCG GYACGAGCAT TAGGACAGTC 120
 TAACGGTCCT GTTACGGTGA TCGAATGACC GACGACATCC TGCTGATCGA CACCGACGAA 180
 CGGGTGCGAA CCCTCACCT CAACCGGCCG CAGTCCCGYA ACGCGCTCTC GGCGGCGCTA 240
 CGGGATCGGT TTTTCGCGGY GTTGGYCGAC GCCGAGGYCG ACGACGACAT CGACGTCGTC 300
 ATCCTCACCG GYGCCGATCC GGTGTTCTGC GCCGACTGG ACCTCAAGGT AGCTGGCCGG 360
 GCAGACCGCG CTGCCGACA TCTACCGCG GTGGGCGGCC ATGACCAAGC CGGTGATCGG 420
 CGCGATCAAC GGCGCCGCG TCACCGGCG GCTCGAACTG GCGCTGTACT GCGACATCCT 480
 GATCGCCTCC GAGCAGCCC GCTTCGNCGA CACCCACGCC CGGGTGGGGC TGCTGCCCAC 540
 CTGGGGACTC AGTGTGTGCT TGCCGCAAAA GGTCCGCATC GGNCTGGGCC GGTGGATGAG 600
 CCTGACCGGC GACTACCTGT CCGTGACCGA CGC 633

(2) INFORMATION FOR SEQ ID NO:7:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 1362 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:7:

CGACGACGAC GGCGCCGGAG AGCGGGCGCG AACGGCGATC GACGCGGCCC TGGCCAGAGT	60
CGGCACCACC CAGGAGGGAG TCGAATCATG AAATTTGTCA ACCATATTGA GCCCCTCGCG	120
CCCCGCCGAG CCGGCGGCGC GGTGCGCGAG GTCTATGCCG AGGCCC GCCG CGAGTTCGGC	180
CGGCTGCCCC AGCCGCTCGC CATGCTGTCC CCGGACGAGG GACTGCTCAC CGCCGGCTGG	240
GCGACGTTGC GCGAGAACT GCTGGTGGGC CAGGTGCCGC GTGGCCGCAA GGAAGCCGTC	300
GCCGCCGCCG TCGCGGCCAG CCTGCGCTGC CCCTGGTGCG TCGACGCACA CACCACCATG	360
CTGTACGCGG CAGGCCAAAC CGACACCGCC GCGGCGATCT TGGCCGGCAC AGCACCTGCC	420
GCCGGTGACC CGAACGCGCC GTATGTGGCG TGGGCGGCAG GAACCGGGAC ACCGGCGGGA	480
CCGCCGGCAC CGTTCGGCCC GGATGTCGCC GCCGAATACC TGGGCACCGC GGTGCAATTC	540
CACTTCATCG CACGCCTGGT CCTGGTGCTG CTGGACGAAA CCTTCCTGCC GGGGGGCCCG	600
CGCGCCCAAC AGCTCATGCG CCGCGCCGGT GGACTGGTGT TCGCCCGCAA GGTGCGCGCG	660
GAGCATCGGC CGGGCCGCTC CACCCGCCGG CTCGAGCCGC GAACGCTGCC CGACGATCTG	720
GCATGGGCAA CACCGTCCGA GCCCATAGCA ACCGCGTTCG CCGCGCTCAG CCACCACCTG	780
GACACCGCGC CGCACCTGCC GCCACCGACT CGTCAGGTGG TCAGGCGGGT CGTGGGGTCG	840
TGGCACGGCG AGCCAATGCC GATGAGCAGT CGCTGGACGA ACGAGCACAC CGCCGAGCTG	900
CCCGCCGACC TGCACGCGCC CACCCGTCTT GCCCTGCTGA CCGGCCTGGC CCCGCATCAG	960
GTGACCGACG ACGACGTCGC CGCGGCCCGA TCCCTGCTCG ACACCGATGC GCGCTGGTT	1020
GGCGCCCTGG CCTGGGCCGC CTTACCGCC GCGCGGCGCA TCGGCACCTG GATCGGCGCC	1080
GCCGCCGAGG GCCAGGTGTC GCGGCAAAAC CCGACTGGGT GAGTGTGCGC GCCCTGTCGG	1140
TAGGGTGTCA TCGCTGGCCC GAGGGATCTC GCGGCGGCGA ACGGAGGTGG CGACACAGGT	1200
GGAAGCTGCG CCCACTGGCT TCGCCCCAA CGCCGTCGTG GGC GTTCGGT TGGCCGCACT	1260
GGCCGATCAG GTCGGCGCCG GCCCTTGCC GAAGGTCCAG CTCAACGTGC CGTCACCGAA	1320

GGACCGGACG GTCACCGGGG GTCACCCTGC GCGCCCAAGG AA

1362

(2) INFORMATION FOR SEQ ID NO:8:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 1458 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:8:

GCGACGACCC CGATATGCCG GGCACCGTAG CGAAAGCCGT CGCCGACGCA CTCGGGCGCG	60
GTATCGCTCC CGTTGAGGAC ATTCAAGACT GCGTGGAGGC CCGGCTGGGG GAAGCCGCTC	120
TGGATGACGT GGCCCGTGTT TACATCATCT ACCGGCAGCG GCGCGCCGAG CTGCGGACGG	180
CTAAGGCCTT GCTCGGCGTG CGGGACGAGT TAAAGCTGAG CTTGGCGGCC GTGACGGTAC	240
TGCGCGAGCG CTATCTGCTG CACGACGAGC AGGGCCGGCC GGCCGAGTCG ACCGGCGAGC	300
TGATGGACCG ATCGGCGCGC TGTGTCGCGG CGGCCGAGGA CCAGTATGAG CCGGGCTCGT	360
CGAGGCGGTG GGCCGAGCGG TTCGCCACGC TATTACGCAA CCTGGAATTC CTGCCGAATT	420
CGCCACGTT GATGAACTCT GGCACCGACC TGGGACTGCT CGCCGGCTGT TTTGTTCTGC	480
CGATTGAGGA TTCGCTGCAA TCGATCTTTG CGACGCTGGG ACAGGCCGCC GAGCTGCAGC	540
GGGCTGGAGG CGGCACCGGA TATGCGTTCA GCCACCTGCG ACCCGCCGGG GATCGGGTGG	600
CCTCCACGGG CGGCACGGCC AGCGGACCGG TGTCGTTTCT ACGGCTGTAT GACAGTGCCG	660
CGGGTGTGGT CTCCATGGGC GGTGCGCCGC GTGGCGCCTG TATGGCTGTG CTTGATGTGT	720
CGCACCCGGA TATCTGTGAT TTCGTCACCG CCAAGGCCGA ATCCCCAGC GAGCTCCCGC	780
ATTTCAACCT ATCGGTTGGT GTGACCGACG CGTTCCTGCG GGCCGTCGAA CGCAACGGCC	840
TACACCGGCT GGTCAATCCG CGAACC GGCA AGATCGTCGC GCGGATGCCC GCCGCCGAGC	900
TGTTGACGC CATCTGCAA GCGCGCACG CCGGTGGCGA TCCCGGGCTG GTGTTTCTCG	960
ACACGATCAA TAGGGCAAAC CCGGTGCCGG GGAGAGGCCG CATCGAGGCG ACCAACCCGT	1020
GCGGGGAGGT CCCACTGCTG CCTTACGAGT CATGTAATCT CGGCTCGATC AACCTCGCCC	1080
GGATGCTCGC CGACGGTCGC GTCGACTGGG ACCGGCTCGA GGAGGTCGCC GGTGTGGCGG	1140
TGCGGTTTCT TGATGACGTC ATCGATGTCA GCCGCTACCC CTTCCCCGAA CTGGGTGAGG	1200

CGGCCCGCGC CACCCGCAAG ATCGGGCTGG GAGTCATGGG TTTGGCGGAA CTGCTTGCCG	1260
CACTGGGTAT TCCGTACGAC AGTGAAGAAG CCGTGCGGTT AGCCACCCGG CTCATGCGTC	1320
GCATACAGCA GGCGGCGCAC ACGGCATCGC GGAGGCTGGC CGAAGAGCGG GGCGCATTCC	1380
CGGCGTTTAC CGATAGCCGG TTCGCGCGGT CGGGCCCGAG GCGCAACGCA CAGGTCACCT	1440
CCGTCGCTCC GACGGGCA	1458

(2) INFORMATION FOR SEQ ID NO:9:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 862 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:9:

ACGGTGTAAT CGTGCTGGAT CTGGAACCGC GTGGCCCGCT ACCTACCGAG ATCTACTGGC	60
GGCGCAGGGG GCTGGCCCTG GGCATCGCGG TCGTCGTAGT CGGGATCGCG GTGGCCATCG	120
TCATCGCCTT CGTCGACAGC AGCGCCGGTG CCAAACCGGT CAGCGCCGAC AAGCCGGCCT	180
CCGCCCAGAG CCATCCGGGC TCGCCGGCAC CCAAGCACC CCAGCCGGCC GGGCAAACCG	240
AAGGTAACGC CGCCGCGGCC CCGCCGCAGG GCCAAAACCC CGAGACACCC ACGCCCACCG	300
CCGCGGTGCA GCCGCCGCCG GTGCTCAAGG AAGGGGACGA TTGCCCCGAT TCGACGCTGG	360
CCGTCAAAGG TTTGACCAAC GCGCCGCACT ACTACGTCGG CGACCAGCCG AAGTTCACCA	420
TGGTGGTCAC CAACATCGGC CTGGTGTCTT GTAAACGCGA CGTTGGGGCC GCGGTGTTGG	480
CCGCCTACGT TTA CTGCTG GACAACAAGC GGTGTGGTCA AACCTGGAC TGCGCGCCCT	540
CGAATGAGAC GCTGGTCAAG ACGTTTTCCC CCGGTGAGCA GGTAACGACC GCGGTGACCT	600
GGACCGGGAT GGGATCGGCG CCGCGCTGCC CATTGCCGCG GCCGGCGATC GGGCCGGGCA	660
CCTACAATCT CGTGGTACAA CTGGGCAATC TCGCTCGCT GCCGGTTCCG TTCATCCTGA	720
ATCAGCCGCC GCCGCCGCC GGGCCGGTAC CCGCTCCGGG TCCAGCGCAG GCGCCTCCGC	780
CGGAGTCTCC CGCGCAAGGC GGATAATTAT TGATCGCTGA TGGTCGATTC CGCCAGCTGT	840
GACAACCCCT CGCCTCGTGC CG	862

(2) INFORMATION FOR SEQ ID NO:10:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 622 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:10:

TTGATCAGCA CCGGCAAGGC GTCACATGCC TCCCTGGGTG TGCAGGTGAC CAATGACAAA	60
GACACCCCGG GCGCCAAGAT CGTCGAAGTA GTGGCCGGTG GTGCTGCCGC GAACGCTGGA	120
GTGCCGAAGG GCGTCGTTGT CACCAAGGTC GACGACCGCC CGATCAACAG CGCGGACGCG	180
TTGGTTGCCG CCGTGCGGTC CAAAGCGCCG GGCGCCACGG TGGCGCTAAC CTTTCAGGAT	240
CCCTCGGGCG GTAGCCGCAC AGTGCAAGTC ACCCTCGGCA AGGCGGAGCA GTGATGAAGG	300
TCGCCGCGCA GTGTTCAAAG CTCGGATATA CGGTGGCACC CATGGAACAG CGTGCGGAGT	360
TGGTGGTTGG CCGGGCACTT GTCGTCGTCG TTGACGATCG CACGGCGCAC GGCGATGAAG	420
ACCACAGCGG GCCGCTTGTC ACCGAGCTGC TCACCGAGGC CGGGTTTGTT GTCGACGGCG	480
TGGTGGCGGT GTCGGCCGAC GAGGTCGAGA TCCGAAATGC GCTGAACACA GCGGTGATCG	540
GCGGGGTGGA CCTGGTGGTG TCGGTCGGCG GGACCGNGT GACGNCTCGC GATGTCACCC	600
CGGAAGCCAC CCGNGACATT CT	622

(2) INFORMATION FOR SEQ ID NO:11:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 1200 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:11:

GGCGCAGCGG TAAGCCTGTT GGCCGCCGGC AACTGGTGT TGACAGCATG CGGCGGTGGC	60
ACCAACAGCT CGTCGTCAGG CGCAGGCGGA ACGTCTGGGT CGGTGCACTG CGGCGGCAAG	120
AAGGAGCTCC ACTCCAGCGG CTCGACCGCA CAAGAAAATG CCATGGAGCA GTTCGTCTAT	180

GCCTACGTGC GATCGTGCCC GGGCTACACG TTGGACTACA ACGCCAACGG GTCCGGTGCC	240
GGGGTGACCC AGTTTCTCAA CAACGAAACC GATTTCGCCG GCTCGGATGT CCCGTTGAAT	300
CCGTCGACCG GTCAACCTGA CCGGTCGGCG GAGCGGTGCG GTTCCCCGGC ATGGGACCTG	360
CCGACGGTGT TCGGCCCGAT CGCGATCACC TACAATATCA AGGGCGTGAG CACGCTGAAT	420
CTTGACGGAC CCACTACCGC CAAGATTTTC AACGGCACCA TCACCGTGTG GAATGATCCA	480
CAGATCCAAG CCCTCAACTC CGGCACCGAC CTGCCGCCAA CACCGATTAG CGTTATCTTC	540
CGCAGCGACA AGTCCGGTAC GTCGGACAAC TTCCAGAAAT ACCTCGACGG TGTATCCAAC	600
GGGGCGTGGG GCAAAGGCGC CAGCGAAACG TTCAGCGGGG GCGTCGGCGT CGGCGCCAGC	660
GGGAACAACG GAACGTCGGC CCTACTGCAG ACGACCGACG GGTGATCAC CTACAACGAG	720
TGGTCGTTTG CGGTGGGTAA GCAGTTGAAC ATGGCCCAGA TCATCACGTC GGCGGGTCCG	780
GATCCAGTGG CGATCACCAC CGAGTCGGTC GGTAAGACAA TCGCCGGGGC CAAGATCATG	840
GGACAAGGCA ACGACCTGGT ATTGGACACG TCGTCGTTCT ACAGACCCAC CCAGCCTGGC	900
TCTTACCCGA TCGTGCTGGC GACCTATGAG ATCGTCTGCT CGAAATACCC GGATGCGACG	960
ACCGGTACTG CGGTAAGGGC GTTTATGCAA GCCGCGATTG GTCCAGGCCA AGAAGGCCTG	1020
GACCAATACG GCTCCATTCC GTTGCCCAA TCGTTCCAAG CAAAATTGGC GGCCGCGGTG	1080
AATGCTATTT CTGACCTAG TGAAGGGAAT TCGACGGTGA GCGATGCCGT TCCGCAGGTA	1140
GGGTCGCAAT TTGGGCCGTA TCAGCTATTG CGGCTGCTGG GCCGAGGCGG GATGGGCGAG	1200

(2) INFORMATION FOR SEQ ID NO:12:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 1155 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:12:

GCAAGCAGCT GCAGGTCGTG CTGTTGACG AACTGGGCAT GCCGAAGACC AAACGCACCA	60
AGACCGGCTA CACCACGGAT GCCGACGCGC TGCAGTCGTT GTTCGACAAG ACCGGGCATC	120
CGTTTCTGCA ACATCTGCTC GCCCACC GCG ACGTCACCCG GCTCAAGGTC ACCGTCGACG	180

GGTTGCTCCA AGCGGTGGCC GCCGACGGCC GCATCCACAC CACGTTCAAC CAGACGATCG	240
CCGCGACCGG CCGGCTCTCC TCGACCGAAC CCAACCTGCA GAACATCCCG ATCCGCACCG	300
ACGCGGGCCG GCGGATCCGG GACGCGTTCG TGGTCGGGGA CGGTTACGCC GAGTTGATGA	360
CGGCCGACTA CAGCCAGATC GAGATGCGGA TCATGGGGCA CCTGTCCGGG GACGAGGGCC	420
TCATCGAGGC GTTCAACACC GGGGAGGACC TGTATTCGTT CGTCGCGTCC CGGGTGTTCG	480
GTGTGCCCAT CGACGAGGTC ACCGGCGAGT TGCGGCGCCG GGTCAAGGCG ATGTCCTACG	540
GGCTGGTTTA CGGGTTGAGC GCCTACGGCC TGTCGCAGCA GTTGAAAATC TCCACCGAGG	600
AAGCCAACGA GCAGATGGAC GCGTATTTTCG CCCGATTCGG CGGGGTGCGC GACTACCTGC	660
GCGCCGTAGT CGAGCGGGCC CGCAAGGACG GCTACACCTC GACGGTGCTG GGCCGTCGCC	720
GCTACCTGCC CGAGCTGGAC AGCAGCAACC GTCAAGTGCG GGAGGCCGCC GAGCGGGCGG	780
CGCTGAACGC GCCGATCCAG GGCAGCGCGG CCGACATCAT CAAGGTGGCC ATGATCCAGG	840
TCGACAAGGC GCTCAACGAG GCACAGCTGG CGTCGCGCAT GCTGCTGCAG GTCCACGACG	900
AGCTGCTGTT CGAAATCGCC CCCGGTGAAC GCGAGCGGGT CGAGGCCCTG GTGCGCGACA	960
AGATGGGCGG CGCTTACCCG CTCGACGTCC CGCTGGAGGT GTCGGTGGGC TACGGCCGCA	1020
GCTGGGACGC GGCGGCGCAC TGAGTGCCGA GCGTGCATCT GGGGCGGGAA TTCGGCGATT	1080
TTTCCGCCCT GAGTTCACGC TCGGCGCAAT CGGGACCGAG TTTGTCCAGC GTGTACCCGT	1140
CGAGTAGCCT CGTCA	1155

(2) INFORMATION FOR SEQ ID NO:13:

- (i) SEQUENCE CHARACTERISTICS:
- (A) LENGTH: 1771 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:13:

GAGCGCCGTC TGGTGTGTTGA ACGGTTTTAC CGGTCGGCAT CGGCACGGGC GTTGCCGGGT	60
TCGGGCCTCG GGTGCGCAT CGTCAAACAG GTGGTGCTCA ACCACGGCGG ATTGCTGCGC	120
ATCGAAGACA CCGACCCAGG CGGCCAGCCC CCTGGAACGT CGATTTACGT GCTGCTCCCC	180

GGCCGTCGGA	TGCCGATTCC	GCAGCTTCCC	GGTGCGACGG	CTGGCGCTCG	GAGCACGGAC	240
ATCGAGAACT	CTCGGGGTTC	GGCGAACGTT	ATCTCAGTGG	AATCTCAGTC	CACGCGCGCA	300
ACCTAGTTGT	GCAGTTACTG	TTGAAAAGCCA	CACCCATGCC	AGTCCACGCA	TGGCCAAGTT	360
GGCCCGAGTA	GTGGGCCTAG	TACAGGAAGA	GCAACCTAGC	GACATGACGA	ATCACCCACG	420
GTATTGCGCA	CCGCCGCAGC	AGCCGGGAAC	CCCAGGTTAT	GCTCAGGGGC	AGCAGCAAAC	480
GTACAGCCAG	CAGTTCGACT	GGCGTTACCC	ACCGTCCCCG	CCCCCGCAGC	CAACCCAGTA	540
CCGTCAACCC	TACGAGGCGT	TGGGTGGTAC	CCGGCCGGGT	CTGATACCTG	GCGTGATTCC	600
GACCATGACG	CCCCCTCCTG	GGATGGTTCG	CCAACGCCCT	CGTGCAGGCA	TGTTGGCCAT	660
CGGCGCGGTG	ACGATAGCGG	TGGTGTCCGC	CGGCATCGGC	GGCGCGGCCG	CATCCCTGGT	720
CGGGTTCAAC	CGGGCACCCG	CCGGCCCCAG	CGGCGGCCCA	GTGGCTGCCA	GCGCGGCGCC	780
AAGCATCCCC	GCAGCAAACA	TGCCGCCGGG	GTCGGTCGAA	CAGGTGGCGG	CCAAGGTGGT	840
GCCCAGTGTC	GTCATGTTGG	AAACCGATCT	GGGCCGCCAG	TCGGAGGAGG	GCTCCGGCAT	900
CATTCTGTCT	GCCGAGGGGC	TGATCTTGAC	CAACAACCAC	GTGATCGCGG	CGGCCGCCAA	960
GCCTCCCCTG	GGCAGTCCGC	CGCCGAAAAC	GACGGTAACC	TTCTCTGACG	GGCGGACCGC	1020
ACCC TTCACG	GTGGTGGGGG	CTGACCCAC	CAGTGATATC	GCCGTCGTCC	GTG TTCAGGG	1080
CGTCTCCGGG	CTCACCCCGA	TCTCCCTGGG	TTCCTCCTCG	GACCTGAGGG	TCGGTCAGCC	1140
GGTGCTGGCG	ATCGGGTCGC	CGCTCGGTTT	GGAGGGCACC	GTGACCACGG	GGATCGTCAG	1200
CGCTCTCAAC	CGTCCAGTGT	CGACGACCGG	CGAGGCCGGC	AACCAGAACA	CCGTGCTGGA	1260
CGCCATT CAG	ACCGACGCCG	CGATCAACCC	CGGTA ACTCC	GGGGGCGCGC	TGGTGAACAT	1320
GAACGCTCAA	CTCGTCGGAG	TCAACTCGGC	CATTGCCACG	CTGGGCGCGG	ACTCAGCCGA	1380
TGCGCAGAGC	GGCTCGATCG	GTCTCGGTTT	TGCGATTCCA	GTCGACCAGG	CCAAGCGCAT	1440
CGCCGACGAG	TTGATCAGCA	CCGGCAAGGC	GTCACATGCC	TCCCTGGGTG	TGCAGGTGAC	1500
CAATGACAAA	GACACCCCGG	GCGCCAAGAT	CGTCGAAGTA	GTGGCCGGTG	GTGCTGCCGC	1560
GAACGCTGGA	GTGCCGAAGG	GCGTCGTTGT	CACCAAGGTC	GACGACCGCC	CGATCAACAG	1620
CGCGGACGCG	TTGGTTGCCG	CCGTGCGGTC	CAAAGCGCCG	GGCGCCACGG	TGGCGCTAAC	1680
CTTTCAGGAT	CCCTCGGGCG	GTAGCCGCAC	AGTGCAAGTC	ACCCTCGGCA	AGGCGGAGCA	1740
GTGATGAAGG	TCGCCGCGCA	GTGTTCAAAG	C			1771

(2) INFORMATION FOR SEQ ID NO:14:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 1058 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:14:

CTCCACCGCG GTGGCGGCCG CTCTAGAACT AGTGGATCCC CCGGGCTGCA GGAATTCGGC	60
ACGAGGATCC GACGTCGCAG GTTGTGCAAC CCGCCGCCGC GGAAGTATCG GTCCATGCCT	120
AGCCCCGGCGA CGGCGAGCGC CGGAATGGCG CGAGTGAGGA GGCGGGCAAT TTGGCGGGGC	180
CCGGCGACGG CGAGCGCCGG AATGGCGCGA GTGAGGAGGC GGGCAGTCAT GCCCAGCGTG	240
ATCCAATCAA CCTGCATTCG GCCTGCGGGC CCATTTGACA ATCGAGGTAG TGAGCGCAAA	300
TGAATGATGG AAAACGGGCG GTGACGTCCG CTGTTCTGGT GGTGCTAGGT GCCTGCCTGG	360
CGTTGTGGCT ATCAGGATGT TCTTCGCCGA AACCTGATGC CGAGGAACAG GGTGTTCCCG	420
TGAGCCCGAC GGCGTCCGAC CCCGCGCTCC TCGCCGAGAT CAGGCAGTCG CTTGATGCGA	480
CAAAAGGGTT GACCAGCGTG CACGTAGCGG TCCGAACAAC CGGGAAAGTC GACAGCTTGC	540
TGGGTATTAC CAGTGCCGAT GTCGACGTCC GGGCCAATCC GCTCGCGGCA AAGGGCGTAT	600
GCACCTACAA CGACGAGCAG GGTGTCCCGT TTCGGGTACA AGGCGACAAC ATCTCGGTGA	660
AACTGTTCTGA CGACTGGAGC AATCTCGGCT CGATTTCTGA ACTGTCAACT TCACGCGTGC	720
TCGATCCTGC CGCTGGGGTG ACGCAGCTGC TGTCCGGTGT CACGAACCTC CAAGCGCAAG	780
GTACCGAAGT GATAGACGGA ATTTGACCA CCAAATCAC CGGGACCATC CCCGCGAGCT	840
CTGTCAAGAT GCTTGATCCT GGCGCCAAGA GTGCAAGGCC GGCGACCGTG TGGATTGCCC	900
AGGACGGCTC GCACCACCTC GTCCGAGCGA GCATCGACCT CGGATCCGGG TCGATTGAGC	960
TCACGCAGTC GAAATGGAAC GAACCCGTCA ACGTCGACTA GGCCGAAGTT GCGTCGACGC	1020
GTTGNTCGAA ACGCCCTTGT GAACGGTGTC AACGGNAC	1058

(2) INFORMATION FOR SEQ ID NO:15:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 542 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single

(D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:15:

GAATTCGGCA CGAGAGGTGA TCGACATCAT CGGGACCAGC CCCACATCCT GGGAACAGGC	60
GGCGGCGGAG GCGGTCCAGC GGGCGCGGGA TAGCGTCGAT GACATCCGCG TCGCTCGGGT	120
CATTGAGCAG GACATGGCCG TGGACAGCGC CGGCAAGATC ACCTACCGCA TCAAGCTCGA	180
AGTGTCGTTC AAGATGAGGC CGGCGCAACC GCGCTAGCAC GGGCCGGCGA GCAAGACGCA	240
AAATCGCACG GTTTGCGGTT GATTCGTGCG ATTTTGTGTC TGCTCGCCGA GGCCTACCAG	300
GCGCGGCCCCA GGTCCGCGTG CTGCCGTATC CAGGCGTGCA TCGCGATTCC GGCGGCCACG	360
CCGGAGTTAA TGCTTCGCGT CGACCCGAAC TGGGCGATCC GCCGGNGAGC TGATCGATGA	420
CCGTGGCCAG CCCGTCGATG CCCGAGTTGC CCGAGGAAAC GTGCTGCCAG GCCGGTAGGA	480
AGCGTCCGTA GGCGGCGGTG CTGACCGGCT CTGCCTGCGC CCTCAGTGCG GCCAGCGAGC	540
GG	542

(2) INFORMATION FOR SEQ ID NO:16:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 913 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:16:

CGGTGCCGCC CGCGCCTCCG TTGCCCCCAT TGCCGCCGTC GCCGATCAGC TGCGCATCGC	60
CACCATCACC GCCTTTGCCG CCGGCACCGC CGGTGGCGCC GGGGCCGCCG ATGCCACCGC	120
TTGACCCTGG CCGCCGGCGC CGCCATTGCC ATACAGCACC CCGCCGGGGG CACCGTTACC	180
GCCGTCGCCA CCGTCGCCG CGCTGCCGTT TCAGGCCGGG GAGGCCGAAT GAACCGCCGC	240
CAAGCCCGCC GCCGGCACCG TTGCCGCCTT TTCCGCCCGC CCCGCCGGCG CCGCCAATTG	300
CCGAACAGCC AMGCACCGTT GCCGCCAGCC CCGCCGCCGT TAACGGCGCT GCCGGGCGCC	360
GCCGCCGGAC CCGCCATTAC CGCCGTTCCC GTTCGGTGCC CCGCCGTTAC CGGCGCCGCC	420

GTTTGCCGCC AATATTCGGC GGGCACCGCC AGACCCGCCG GGGCCACCAT TGCCGCCGGG	480
CACCGAAACA ACAGCCCAAC GGTGCCGCCG GCCCGCCGT TTGCCGCCAT CACCGGCCAT	540
TCACCGCCAG CACCGCCGTT AATGTTTATG AACCCGGTAC CGCCAGCGCG GCCCTATTG	600
CCGGGCGCCG GAGNGCGTGC CCGCCGGCGC CGCCAACGCC CAAAAGCCCG GGGTTGCCAC	660
CGGCCCCGCC GGACCCACCG GTCCCGCCGA TCCCCCGTT GCCGCCGGTG CCGCCGCCAT	720
TGGTGCTGCT GAAGCCGTTA GCGCCGGTTC CGCSGGTTCC GGCGGTGGCG CCNTGGCCGC	780
CGGCCCCGCC GTTGCCGTAC AGCCACCCCC CGGTGGCGCC GTTGCCGCCA TTGCCGCCAT	840
TGCCGCCGTT GCCGCCATTG CCGCCGTTCC CGCCGCCACC GCCGNTTGG CCGCCGGCGC	900
CGCCGGCGGC CGC	913

(2) INFORMATION FOR SEQ ID NO:17:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 1872 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:17:

GACTACGTTG GTGTAGAAAA ATCTGCCGC CCGGACCCTT AAGGCTGGGA CAATTTCTGA	60
TAGCTACCCC GACACAGGAG GTTACGGGAT GAGCAATTCG CGCCGCCGCT CACTCAGGTG	120
GTCATGGTTG CTGAGCGTGC TGGCTGCCGT CGGGCTGGGC CTGGCCACGG CGCCGGCCCA	180
GGCGGCCCCG CCGGCCTTGT CGCAGGACCG GTTCGCCGAC TTCCCCGCGC TGCCCCTCGA	240
CCCGTCCGCG ATGGTCGCCC AAGTGGCGCC ACAGGTGGTC AACATCAACA CCAAACCTGGG	300
CTACAACAAC GCCGTGGGCG CCGGGACCGG CATCGTCATC GATCCCAACG GTGTCGTGCT	360
GACCAACAAC CACGTGATCG CGGGCGCCAC CGACATCAAT GCGTTCAGCG TCGGCTCCGG	420
CCAAACCTAC GGCGTCGATG TGGTCGGGTA TGACCGCACC CAGGATGTCG CGGTGCTGCA	480
GCTGCGCGGT GCCGGTGGCC TGCCGTCGGC GGCATCGGT GGCGGCGTCG CGGTTGGTGA	540
GCCCGTCGTC GCGATGGGCA ACAGCGGTGG GCAGGGCGGA ACGCCCCGTG CGGTGCCCTGG	600
CAGGGTGGTC GCGCTCGGCC AAACCGTGCA GCGTCGGAT TCGCTGACCG GTGCCGAAGA	660

GACATTGAAC GGGTTGATCC AGTTCGATGC CGCAATCCAG CCCGGTGATT CGGGCGGGCC	720
CGTCGTCAAC GGCCTAGGAC AGGTGGTCGG TATGAACACG GCCGCGTCCG ATAACTTCCA	780
GCTGTCCCAG GGTGGGCAGG GATTGCCCAT TCCGATCGGG CAGGCGATGG CGATCGCGGG	840
CCAAATCCGA TCGGGTGGGG GGTCACCCAC CGTTCATATC GGGCCTACCG CCTTCCTCGG	900
CTTGGGTGTT GTCGACAACA ACGGCAACGG CGCACGAGTC CAACGCGTGG TCGGAAGCGC	960
TCCGGCGGCA AGTCTCGGCA TCTCCACCGG CGACGTGATC ACCGCGGTCG ACGGCGCTCC	1020
GATCAACTCG GCCACCGCGA TGGCGGACGC GCTTAACGGG CATCATCCCG GTGACGTCAT	1080
CTCGGTGAAC TGGCAAACCA AGTCGGGCGG CACGCGTACA GGGAACTGA CATTGGCCGA	1140
GGGACCCCCG GCCTGATTG TCGCGGATAC CACCCGCCGG CCGGCCAATT GGATTGGCGC	1200
CAGCCGTGAT TGCCGCGTGA GCCCCGAGT TCCGTCTCCC GTGCGCGTGG CATTGTGGAA	1260
GCAATGAACG AGGCAGAACA CAGCGTTGAG CACCCTCCCG TGCAGGGCAG TTACGTCGAA	1320
GGCGGTGTGG TCGAGCATCC GGATGCCAAG GACTTCGGCA GCGCCGCCGC CCTGCCCGCC	1380
GATCCGACCT GGTTTAAGCA CGCCGTCTTC TACGAGGTGC TGGTCCGGGC GTTCTTCGAC	1440
GCCAGCGCGG ACGGTTCCGN CGATCTGCGT GGACTCATCG ATCGCCTCGA CTACCTGCAG	1500
TGGCTTGGA TCGACTGCAT CTGTTGCCGC CGTTCCTACG ACTCACCGCT GCGCGACGGC	1560
GGTTACGACA TTCGCGACTT CTACAAGGTG CTGCCCCAAT TCGGCACCGT CGACGATTTT	1620
GTCGCCCTGG TCGACACCGC TCACCGGCGA GGTATCCGCA TCATCACCGA CCTGGTGATG	1680
AATCACACCT CGGAGTCGCA CCCCTGGTTT CAGGAGTCCC GCCGCGACCC AGACGGACCG	1740
TACGGTGAAT ATTACGTGTG GAGCGACACC AGCGAGCGCT ACACCGACGC CCGGATCATC	1800
TTCGTCGACA CCGAAGAGTC GAACTGGTCA TTCGATCCTG TCCGCCGACA GTTNCTACTG	1860
GCACCGATTC TT	1872

(2) INFORMATION FOR SEQ ID NO:18:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 1482 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:18:

CTTCGCCGAA ACCTGATGCC GAGGAACAGG GTGTTCCCGT GAGCCCGACG GCGTCCGACC	60
CCGCGCTCCT CGCCGAGATC AGGCAGTCGC TTGATGCGAC AAAAGGGTTG ACCAGCGTGC	120
ACGTAGCGGT CCGAACAAACC GGGAAAGTCG ACAGCTTGCT GGGTATTACC AGTGCCGATG	180
TCGACGTCCG GGCCAATCCG CTCGCGGCAA AGGGCGTATG CACCTACAAC GACGAGCAGG	240
GTGTCCCGTT TCGGGTACAA GGCGACAACA TCTCGGTGAA ACTGTTGAC GACTGGAGCA	300
ATCTCGGCTC GATTTCTGAA CTGTCAACTT CACGCGTGCT CGATCCTGCC GCTGGGGTGA	360
CGCAGCTGCT GTCCGGTGTC ACGAACCTCC AAGCGCAAGG TACCGAAGTG ATAGACGGAA	420
TTTCGACCAC CAAAATCACC GGGACCATCC CCGCGAGCTC TGTCAAGATG CTTGATCCTG	480
GCGCCAAGAG TGCAAGGCCG GCGACCGTGT GGATTGCCCA GGACGGCTCG CACCACCTCG	540
TCCGAGCGAG CATCGACCTC GGATCCGGGT CGATTAGCT CACGCAGTCG AAATGGAACG	600
AACCCGTCAA CGTCGACTAG GCCGAAGTTG CGTCGACGCG TTGCTCGAAA CGCCCTTGTTG	660
AACGGTGTCA ACGGCACCCG AAAACTGACC CCCTGACGGC ATCTGAAAAT TGACCCCCTA	720
GACCGGGCGG TTGGTGTTA TTCTTCGGTG GTTCCGGCTG GTGGGACGCG GCCGAGGTCG	780
CGGTCTTTGA GCCGGTAGCT GTCGCCTTTG AGGGCGACGA CTTCAGCATG GTGGACGAGG	840
CGGTCGATCA TGGCGGCAGC AACGACGTCG TCGCCGCCGA AAACCTCGCC CCACCGGCCG	900
AAGGCCTTAT TGGACGTGAC GATCAAGCTG GCCCGCTCAT ACCGGGAGGA CACCAGCTGG	960
AAGAAGAGGT TGGCGGCCTC GGGCTCAAAC GGAATGTAAC CGACTTCGTC AACCACCAGG	1020
AGCGGATAGC GGCCAAACCG GGTGAGTTCG GCGTAGATGC GCCCGGCGTG GTGAGCCTCG	1080
GCGAACCGTG CTACCCATTC GGCGGCGGTG GCGAACAGCA CCCGATGACC GGCCTGACAC	1140
GCGCGTATCG CCAGGCCGAC CGCAAGATGA GTCTTCCCGG TGCCAGGCGG GGCCCCAAAAA	1200
CACGACGTTA TCGCGGGCGG TGATGAAATC CAGGGTGCCC AGATGTGCGA TGGTGTCGCG	1260
TTTGAGGCCA CGAGCATGCT CAAAGTCGAA CTCTTCCAAC GACTTCCGAA CCGGGAAGCG	1320
GGCGGCGCGG ATGCGGCCCT CACCACCATG GGAATCCCGG GCTGACACTT CCCGCTGCAG	1380
GCAGGCGGCC AGGTATTCTT CGTGGCTCCA GTTCTCGGCG CGGGCGCGAT CGGCCAGCCG	1440
GGACACTGAC TCACGCAGG TGGGAGCTTT CAATGCTCTT GT	1482

(2) INFORMATION FOR SEQ ID NO:19:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 876 base pairs

- (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:19:

GAATTCGGCA CGAGCCGGCG ATAGCTTCTG GGCCGCGGCC GACCAGATGG CTCGAGGGTT	60
CGTGCTCGGG GCCACCGCCG GGCGCACCAC CCTGACCGGT GAGGGCCTGC AACACGCCGA	120
CGGTCACCTCG TTGCTGCTGG ACGCCACCAA CCCGGCGGTG GTTGCCTACG ACCCGGCCCTT	180
CGCCTACGAA ATCGGCTACA TCGNGGAAAG CGGACTGGCC AGGATGTGCG GGGAGAACCC	240
GGAGAACATC TTCTTCTACA TCACCGTCTA CAACGAGCCG TACGTGCAGC CGCCGGAGCC	300
GGAGAACTTC GATCCCGAGG GCGTGCTGGG GGGTATCTAC CGNTATCACG CGGCCACCGA	360
GCAACGCACC AACAAGNGC AGATCCTGGC CTCCGGGGTA GCGATGCCCC CGGCGCTGCG	420
GGCAGCACAG ATGCTGGCCG CCGAGTGGGA TGTCGCCGCC GACGTGTGGT CGGTGACCAG	480
TTGGGGCGAG CTAAACCGCG ACGGGGTGGT CATCGAGACC GAGAAGCTCC GCCACCCCGA	540
TCGGCCGGCG GGCGTGCCCT ACGTGACGAG AGCGCTGGAG AATGCTCGGG GCCCGGTGAT	600
CGCGGTGTCG GACTGGATGC GCGCGGTCCC CGAGCAGATC CGACCGTGGG TGCCGGGCAC	660
ATACCTCACG TTGGGCACCG ACGGGTTCGG TTTTCCGAC ACTCGGCCCC CCGGTCGTCTG	720
TTACTTCAAC ACCGACGCCG AATCCCAGGT TGGTCGCGGT TTTGGGAGGG GTTGGCCGGG	780
TCGACGGGTG AATATCGACC CATTCGGTGC CGGTCGTGGG CCGCCCGCCC AGTTACCCGG	840
ATTCGACGAA GGTGGGGGGT TGCGCCCGAN TAAGTT	876

(2) INFORMATION FOR SEQ ID NO:20:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 1021 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:20:

ATCCCCCGG GCTGCAGGAA TTCGGCACGA GAGACAAAAT TCCACGCGTT AATGCAGGAA	60
--	----

CAGATTCATA ACGAATTCAC AGCGGCACAA CAATATGTCG CGATCGCGGT TTATTTTCGAC 120
AGCGAAGACC TGCCGCAGTT GGCGAAGCAT TTTTACAGCC AAGCGGTCGA GGAACGAAAC 180
CATGCAATGA TGCTCGTGCA ACACCTGCTC GACCGCGACC TTCGTGTCGA AATTTCCGGC 240
GTAGACACGG TGCGAAACCA GTTCGACAGA CCCC GCGAGG CACTGGCGCT GCGGCTCGAT 300
CAGGAACGCA CAGTCACCGA CCAGGTCGGT CGGCTGACAG CGGTGGCCCG CGACGAGGGC 360
GATTTCCCTCG GCGAGCAGTT CATGCAGTGG TTCTTGACAG AACAGATCGA AGAGGTGGCC 420
TTGATGGCAA CCCTGGTGCG GGTGCGGAT CGGGCCGGG CCAACCTGTT CGAGCTAGAG 480
AACTTCGTCG CACGTGAAGT GGATGTGGCG CCGGCCGCAT CAGGCGCCCC GCACGCTGCC 540
GGGGGCCGCC TCTAGATCCC TGGGGGGGAT CAGCGAGTGG TCCCGTTCGC CCGCCCGTCT 600
TCCAGCCAGG CCTTGGTGCG GCCGGGGTGG TGAGTACCAA TCCAGGCCAC CCGACCTCC 660
CGGNAAAAGT CGATGTCTC GTACTCATCG ACGTTCCAGG AGTACACCGC CCGGCCCTGA 720
GCTGCCGAGC GGTCAACGAG TTGCGGATAT TCCTTTAACG CAGGCAGTGA GGGTCCCACG 780
GCGGTTGGCC CGACCGCCGT GGCCGCACTG CTGGTCAGGT ATCGGGGGGT CTTGGCGAGC 840
AACAACGTCG GCAGGAGGGG TGGAGCCCGC CGGATCCGCA GACCGGGGGG GCGAAAACGA 900
CATCAACACC GCACGGGATC GATCTGCGGA GGGGGGTGCG GGAATACCGA ACCGGTGTAG 960
GAGCGCCAGC AGTTGTTTTT CCACCAGCGA AGCGTTTTTCG GGTCAATCGG GGCNNTTAAG 1020
T 1021

(2) INFORMATION FOR SEQ ID NO:21:

- (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 321 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:21:

CGTGCCGACG AACGGAAGAA CACAACCATG AAGATGGTGA AATCGATCGC CGCAGGTCTG 60
ACCGCCGCGG CTGCAATCGG CGCCGCTGCG GCCGGTGTGA CTTCGATCAT GGCTGGCGGN 120
CCGGTCGTAT ACCAGATGCA GCCGGTCGTC TTCGGCGCGC CACTGCCGTT GGACCCGGNA 180

TCCGCCCCTG ANGTCCTGAC CGCCGCCAG TGGACCAGNC TGCTCAACAG NCTCGNCGAT	240
CCCAACGTGT CGTTTGNAA CAAGGGNAGT CTGGTCGAGG GNGGNATCGG NGGNANCGAG	300
GGNGNGNATC GNCGANACA A	321

(2) INFORMATION FOR SEQ ID NO:22:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 373 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:22:

TCTTATCGGT TCCGGTTGGC GACGGGTTTT GGGNGCGGGT GGTTAACCCG CTCGGCCAGC	60
CGATCGACGG GCGCGGAGAC GTCGACTCCG ATACTCGGCG CGCGCTGGAG CTCCAGGCGC	120
CCTCGGTGGT GNACCGGCAA GGCGTGAAGG AGCCGTTGNA GACCGGGATC AAGGCGATTG	180
ACGCGATGAC CCCGATCGGC CGCGGGCAGC GCCAGCTGAT CATCGGGGAC CGCAAGACCG	240
GCAAAAACCG CCGTCTGTGT CGGACACCAT CCTCAAACCA GCGGGAAGAA CTGGGAGTCC	300
GGTGGATCCC AAGAAGCAGG TGCGCTTGTG TATACGTTGG CCATCGGGCA AGAAGGGGAA	360
CTTACCATCG CCG	373

(2) INFORMATION FOR SEQ ID NO:23:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 352 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:23:

GTGACGCCGT GATGGGATTC CTGGGCGGGG CCGGTCCGCT GGCGGTGGTG GATCAGCAAC	60
TGGTTACCCG GGTGCCGCAA GGCTGGTCGT TTGCTCAGGC AGCCGCTGTG CCGGTGGTGT	120
TCTTGACGGC CTGGTACGGG TTGGCCGATT TAGCCGAGAT CAAGGCGGGC GAATCGGTGC	180
TGATCCATGC CGGTACCGGC GGTGTGGGCA TGGCGGCTGT GCAGCTGGCT CGCCAGTGGG	240

GCGTGGAGGT TTTCGTCACC GCCAGCCGTG GNAAGTGGGA CACGCTGCGC GCCATNGNGT 300
TTGACGACGA NCCATATCGG NGATTCCCNC ACATNCGAAG TTCCGANGGA GA 352

(2) INFORMATION FOR SEQ ID NO:24:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 726 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:24:

GAAATCCGCG TTCATTCCGT TCGACCAGCG GCTGGCGATA ATCGACGAAG TGATCAAGCC 60
GCGGTTTCGCG GCGCTCATGG GTCACAGCGA GTAATCAGCA AGTTCTCTGG TATATCGCAC 120
CTAGCGTCCA GTTGCTTGCC AGATCGCTTT CGTACCGTCA TCGCATGTAC CGGTTTCGCGT 180
GCCGCACGCT CATGCTGGCG GCGTGCATCC TGGCCACGGG TGTGGCGGGT CTCGGGGTTCG 240
GCGCGCAGTC CGCAGCCCAA ACCGCGCCGG TGCCCCACTA CTACTGGTGC CCGGGGCAGC 300
CTTTTCGACCC CGCATGGGGG CCCAACTGGG ATCCCTACAC CTGCCATGAC GACTTCCACC 360
GCGACAGCGA CGGCCCCGAC CACAGCCGCG ACTACCCCGG ACCCATCCTC GAAGGTCCCG 420
TGCTTGACGA TCCCGGTGCT GCGCCGCCGC CCCC GGCTGC CGGTGGCGGC GCATAGCGCT 480
CGTTGACCGG GCCGCATCAG CGAATACGCG TATAAACCCG GCGGTGCCCC CGGCAAGCTA 540
CGACCCCGCG CGGGGCAGAT TTACGCTCCC GTGCCGATGG ATCGCGCCGT CCGATGACAG 600
AAAATAGGCG ACGGTTTTGG CAACCGCTTG GAGGACGCTT GAAGGGAACC TGTCATGAAC 660
GGCGACAGCG CCTCCACCAT CGACATCGAC AAGGTTGTTA CCCGCACACC CGTTCGCCGG 720
ATCGTG 726

(2) INFORMATION FOR SEQ ID NO:25:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 580 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:25:

CGCGACGACG	ACGAACGTCG	GGCCCACCAC	CGCCTATGCG	TTGATGCAGG	CGACCGGGAT	60
GGTCGCCGAC	CATATCCAAG	CATGCTGGGT	GCCCACTGAG	CGACCTTTTG	ACCAGCCGGG	120
CTGCCCCGATG	GCGGCCCGGT	GAAGTCATTG	CGCCGGGGCT	TGTGCACCTG	ATGAACCCGA	180
ATAGGGAACA	ATAGGGGGGT	GATTTGGCAG	TTCAATGTCG	GGTATGGCTG	GAAATCCAAT	240
GGCGGGGCAT	GCTCGGCGCC	GACCAGGCTC	GCGCAGGCGG	GCCAGCCCGA	ATCTGGAGGG	300
AGCACTCAAT	GGCGGCGATG	AAGCCCCGGA	CCGGCGACGG	TCCTTTGGAA	GCAACTAAGG	360
AGGGGCGCGG	CATTGTGATG	CGAGTACCAC	TTGAGGGTGG	CGGTCGCCTG	GTCGTCGAGC	420
TGACACCCGA	CGAAGCCGCC	GCACTGGGTG	ACGAACTCAA	AGGCGTTACT	AGCTAAGACC	480
AGCCCAACGG	CGAATGGTCG	GCGTTACGCG	CACACCTTCC	GGTAGATGTC	CAGTGTCTGC	540
TCGGCGATGT	ATGCCCAGGA	GAACTCTTGG	ATACAGCGCT			580

(2) INFORMATION FOR SEQ ID NO:26:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 160 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:26:

AACGGAGGCG	CCGGGGGTTT	TGGCGGGGCC	GGGGCGGTCG	GCGGCAACGG	CGGGGCCGGC	60
GGTACCGCCG	GGTTGTTCGG	TGTCGGCGGG	GCCGGTGGGG	CCGGAGGCAA	CGGCATCGCC	120
GGTGTACGCG	GTACGTCGGC	CAGCACACCG	GGTGGATCCG			160

(2) INFORMATION FOR SEQ ID NO:27:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 272 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:27:

GACACCGATA CGATGGTGAT GTACGCCAAC GTTGTGACA CGCTCGAGGC GTTCACGATC	60
CAGCGCACAC CCGACGGCGT GACCATCGGC GATGCGGCCC CGTTCGCGGA GGCGGCTGCC	120
AAGGCGATGG GAATCGACAA GCTGCGGGTA ATTCATACCG GAATGGACCC CGTCGTCGCT	180
GAACGCGAAC AGTGGGACGA CGGCAACAAC ACGTTGGCGT TGGCGCCCGG TGTCGTTGTC	240
GCCTACGAGC GCAACGTACA GACCAACGCC CG	272

(2) INFORMATION FOR SEQ ID NO:28:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 317 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:28:

GCAGCCGGTG GTTCTCGGAC TATCTGCGCA CGGTGACGCA GCGCGACGTG CGCGAGCTGA	60
AGCGGATCGA GCAGACGGAT CGCCTGCCGC GGTTCATGCG CTACCTGGCC GCTATCACCG	120
CGCAGGAGCT GAACGTGGCC GAAGCGGCGC GGGTCATCGG GGTCGACGCG GGGACGATCC	180
GTTCGGATCT GGCGTGGTTC GAGACGGTCT ATCTGGTACA TCGCCTGCCC GCCTGGTCGC	240
GGAATCTGAC CGCGAAGATC AAGAAGCGGT CAAAGATCCA CGTCGTCGAC AGTGGCTTCG	300
CGGCCTGGTT GCGCGGG	317

(2) INFORMATION FOR SEQ ID NO:29:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 182 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:29:

GATCGTGGAG CTGTCGATGA ACAGCGTTGC CGGACGCGCG GCGGCCAGCA CGTCGGTGTA	60
---	----

GCAGCGCCGG ACCACCTCGC CGGTGGGCAG CATGGTGATG ACCACGTCGG CCTCGGCCAC 120
 CGCTTCGGGC GCGCTACGAA ACACCGCGAC ACCGTGCGCG GCGGCGCCGG ACGCCGCCGT 180
 GG 182

(2) INFORMATION FOR SEQ ID NO:30:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 308 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:30:

GATCGCGAAG TTTGGTGAGC AGGTGGTCGA CGCGAAAGTC TGGGCGCCTG CGAAGCGGGT 60
 CGGCGTTCAC GAGGCGAAGA CACGCCTGTC CGAGCTGCTG CGGCTCGTCT ACGGCGGGCA 120
 GAGGTTGAGA TTGCCCCGCCG CGGCGAGCCG GTAGCAAAGC TTGTGCCGCT GCATCCTCAT 180
 GAGACTCGGC GGTTAGGCAT TGACCATGGC GTGTACCGCG TGCCCGACGA TTTGGACGCT 240
 CCGTTGTCAG ACGACGTGCT CGAACGCTTT CACCGGTGAA GCGCTACCTC ATCGACACCC 300
 ACGTTTGG 308

(2) INFORMATION FOR SEQ ID NO:31:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 267 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:31:

CCGACGACGA GCAACTCACG TGGATGATGG TCGGCAGCGG CATTGAGGAC GGAGAGAATC 60
 CGGCCGAAGC TGCCGCGCGG CAAGTGCTCA TAGTGACCGG CCGTAGAGGG CTCCCCCGAT 120
 GGCACCGGAC TATTCTGGTG TGCCGCTGGC CGGTAAGAGC GGGTAAAAGA ATGTGAGGGG 180
 ACACGATGAG CAATCACACC TACCGAGTGA TCGAGATCGT CGGGACCTCG CCCGACGGCG 240
 TCGACGCGGC AATCCAGGGC GGTCTGG 267

(2) INFORMATION FOR SEQ ID NO:32:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 1539 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:32:

CTCGTGCCGA AAGAATGTGA GGGGACACGA TGAGCAATCA CACCTACCGA GTGATCGAGA	60
TCGTCGGGAC CTCGCCCAGAC GCGGTCGACG CGGCAATCCA GGGCGGTCTG GCCCGAGCTG	120
CGCAGACCAT GCGCGCGCTG GACTGGTTTCG AAGTACAGTC AATTCGAGGC CACCTGGTCG	180
ACGGAGCGGT CGCGCACTTC CAGGTGACTA TGAAAGTCGG CTTCGCTGG AGGATTCCTG	240
AACCTTCAAG CGCGGCCGAT AACTGAGGTG CATCATTAAG CGACTTTTCC AGAACATCCT	300
GACGCGCTCG AAACGCGGTT CAGCCGACGG TGGCTCCGCC GAGGCGCTGC CTCCAAAATC	360
CCTGCGACAA TTCGTCGGCG GCGCCTACAA GGAAGTCGGT GCTGAATTCG TCGGGTATCT	420
GGTCGACCTG TGTGGGCTGC AGCCGGACGA AGCGGTGCTC GACGTCGGCT GCGGCTCGGG	480
GCGGATGGCG TTGCCGCTCA CCGGCTATCT GAACAGCGAG GGACGCTACG CCGGCTTCGA	540
TATCTCGCAG AAAGCCATCG CGTGGTGCCA GGAGCACATC ACCTCGGCGC ACCCCAACCT	600
CCAGTTCGAG GTCTCCGACA TCTACAACCTC GCTGTACAAC CCGAAAGGGA AATACCAGTC	660
ACTAGACTTT CGCTTTCCAT ATCCGGATGC GTCGTTTCGAT GTGGTGTTTC TTACCTCGGT	720
GTTACCCAC ATGTTTCCGC CGGACGTGGA GCACTATCTG GACGAGATCT CCCGCTGCT	780
GAAGCCCGGC GGACGATGCC TGTGCACGTA CTTCTTGCTC AATGACGAGT CGTTAGCCCA	840
CATCGCGGAA GGAAAGAGTG CGCACAACTT CCAGCATGAG GGACCGGGTT ATCGGACAAT	900
CCACAAGAAG CGGCCGAAG AAGCAATCGG CTTGCCGGAG ACCTTCGTCA GGGATGTCTA	960
TGGCAAGTTC GGCCTCGCCG TGCACGAACC ATTGCACTAC GGCTCATGGA GTGGCCGGGA	1020
ACCACGCCTA AGCTTCCAGG ACATCGTCAT CGCGACCAA ACCGCGAGCT AGGTCGGCAT	1080
CCGGGAAGCA TCGCGACACC GTGGCGCCGA GCGCCGCTGC CGGCAGGCCG ATTAGGCGGG	1140
CAGATTAGCC CGCCGCGGCT CCCGGCTCCG AGTACGGCGC CCCGAATGGC GTCACCGGCT	1200
GGTAACCACG CTTGCGCGCC TGGGCGGCGG CCTGCCGGAT CAGGTGGTAG ATGCCGACAA	1260

AGCCTGCGTG ATCGGTCATC ACCAACGGTG ACAGCAGCCG GTTGTGCACC AGCGCGAACG	1320
CCACCCCGGT CTCCGGGTCT GTCCAGCCGA TCGAGCCGCC CAAGCCCACA TGACCAAACC	1380
CCGGCATCAC GTTGCCGATC GGCATACCGT GATAGCCAAG ATGAAAATTT AAGGGCACCA	1440
ATAGATTTTCG ATCCGGCAGA ACTTGCCGTC GGTTCGGGGT CAGGCCCGTG ACCAGCTCCC	1500
GCGACAAGAA CCGTATGCCG TCGATCTCGC CTCGTGCCG	1539

(2) INFORMATION FOR SEQ ID NO:33:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 851 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:33:

CTGCAGGGTG GCGTGGATGA GCGTCACCGC GGGGCAGGCC GAGCTGACCG CCGCCCAGGT	60
CCGGGTTGCT GCGGCGGCCT ACGAGACGGC GTATGGGCTG ACGGTGCCCC CGCCGGTGAT	120
CGCCGAGAAC CGTGCTGAAC TGATGATTCT GATAGCGACC AACCTCTTGG GGCAAAACAC	180
CCCGGCGATC GCGGTCAACG AGGCCGAATA CGGCGAGATG TGGGCCCCAAG ACGCCGCCGC	240
GATGTTTGGC TACGCCGCGG CGACGGCGAC GGCGACGGCG ACGTTGCTGC CGTTCGAGGA	300
GGCGCCGGAG ATGACCAGCG CGGGTGGGCT CCTCGAGCAG GCCGCCGCGG TCGAGGAGGC	360
CTCCGACACC GCCGCGGCGA ACCAGTTGAT GAACAATGTG CCCCAGGCGC TGAAACAGTT	420
GGCCCAGCCC ACGCAGGGCA CCACGCCTTC TTCCAAGCTG GGTGGCCTGT GGAAGACGGT	480
CTCGCCGCAT CGGTCGCCGA TCAGCAACAT GGTGTGCGATG GCCAACAACC ACATGTCGAT	540
GACCAACTCG GGTGTGTCGA TGACCAACAC CTTGAGCTCG ATGTTGAAGG GCTTTGCTCC	600
GGCGGCGGCC GCCCAGGCCG TGCAAACCGC GGCGCAAAAC GGGGTCCGGG CGATGAGCTC	660
GCTGGGCAGC TCGCTGGGTT CTTGCGGTCT GGGCGGTGGG GTGGCCGCCA ACTTGGGTCTG	720
GGCGGCCTCG GTACGGTATG GTCACCGGGA TGGCGGAAAA TATGCANAGT CTGGTCGGCG	780
GAACGGTGGT CCGGCGTAAG GTTTACCCCC GTTTTCTGGA TGCGGTGAAC TTCGTCAACG	840
GAAACAGTTA C	851

(2) INFORMATION FOR SEQ ID NO:34:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 254 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:34:

GATCGATCGG GCGGAAATTT GGACCAGATT CGCCTCCGGC GATAACCCAA TCAATCGAAC	60
CTAGATTTAT TCCGTCCAGG GGCCCGAGTA ATGGCTCGCA GGAGAGGAAC CTTACTGCTG	120
CGGGCACCTG TCGTAGGTCC TCGATACGGC GGAAGGCGTC GACATTTTCC ACCGACACCC	180
CCATCCAAAC GTTCGAGGGC CACTCCAGCT TGTGAGCGAG GCGACGCAGT CGCAGGCTGC	240
GCTTGGTCAA GATC	254

(2) INFORMATION FOR SEQ ID NO:35:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 1227 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:35:

GATCCTGACC GAAGCGGCCG CCGCCAAGGC GAAGTCGCTG TTGGACCAGG AGGGACGGGA	60
CGATCTGGCG CTGCGGATCG CGGTTAGCC GGGGGGGTGC GCTGGATTGC GCTATAACCT	120
TTTCTTCGAC GACCGGACGC TGGATGGTGA CCAAACCGCG GAGTTCGGTG GTGTCAGGTT	180
GATCGTGGAC CGGATGAGCG CGCCGTATGT GGAAGGCGCG TCGATCGATT TCGTCGACAC	240
TATTGAGAAG CAAGGTTTAC CATCGACAAT CCCAACGCCA CCGGCTCCTG CGCGTGCGGG	300
GATTCGTTCA ACTGATAAAA CGCTAGTACG ACCCCGCGGT GCGCAACACG TACGAGCACA	360
CCAAGACCTG ACCGCGCTGG AAAAGCAACT GAGCGATGCC TTGCACCTGA CCGCGTGGCG	420
GGCCGCCGGC GGCAGGTGTC ACCTGCATGG TGAACAGCAC CTGGGCCTGA TATTGCGACC	480
AGTACACGAT TTTGTCGATC GAGGTCACCTT CGACCTGGGA GAACTGCTTG CGGAACGCGT	540

CGCTGCTCAG CTTGGCCAAG GCCTGATCGG AGCGCTTGTC GCGCACGCCG TCGTGGATAC 600
CGCACAGCGC ATTGCGAACG ATGGTGTCCA CATCGCGGTT CTCCAGCGCG TTGAGGTATC 660
CCTGAATCGC GGTTTTGGCC GGTCCCTCCG AGAATGTGCC TGCCGTGTTG GCTCCGTTGG 720
TGCGGACCCC GTATATGATC GCCGCCGTCA TAGCCGACAC CAGCGCGAGG GCTACCACAA 780
TGCCGATCAG CAGCCGCTTG TGCCGTCGCT TCGGGTAGGA CACCTGCGGC GGCACGCCGG 840
GATATGCGGC GGGCGGCAGC GCCGCGTCGT CTGCCGGTCC CGGGGCGAAG GCCGGTTCGG 900
CGGCGCCGAG GTCGTGGGGG TAGTCCAGGG CTTGGGGTTC GTGGGATGAG GGCTCGGGGT 960
ACGGCGCCGG TCCGTTGGTG CCGACACCGG GGTTCGGCGA GTGGGGACCG GGCATTGTGG 1020
TTCTCCTAGG GTGGTGGACG GGACCAGCTG CTAGGGCGAC AACCGCCCGT CGCGTCAGCC 1080
GGCAGCATCG GCAATCAGGT GAGCTCCCTA GGCAGGCTAG CGCAACAGCT GCCGTCAGCT 1140
CTCAACGCGA CGGGGCGGGC CGCGGCGCCG ATAATGTTGA AAGACTAGGC AACCTTAGGA 1200
ACGAAGGACG GAGATTTTGT GACGATC 1227

(2) INFORMATION FOR SEQ ID NO:36:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 181 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:36:

GCGGTGTCGG CGGATCCGGC GGGTGGTTGA ACGGCAACGG CGGGGCCGGC GGGGCCGGCG 60
GGACCGGCGC TAACGGTGGT GCCGGCGGCA ACGCCTGGTT GTTCGGGGCC GGCGGGTCCG 120
GCGGNGCCGG CACCAATGGT GGNGTCGGCG GGTCCGGCGG ATTTGTCTAC GGCAACGGCG 180
G 181

(2) INFORMATION FOR SEQ ID NO:37:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 290 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:37:

GCGGTGTCGG CGGATCCGGC GGGTGTTGA ACGGCAACGG CGGTGTCGGC GGCCGGGGCG 60
GCGACGGCGT CTTTGCCGGT GCCGGCGGCC AGGGCGGCCT CGGTGGGCAG GCGGGCAATG 120
GCGGCGGCTC CACCGGCGGC AACGGCGGTC TTGGCGGCGC GGGCGGTGGC GGAGGCAACG 180
CCCCGGACGG CGGCTTCGGT GGCAACGGCG GTAAGGGTGG CCAGGGCGGN ATTGGCGGCG 240
GCACTCAGAG CGCGACCGGC CTCGGNGGTG ACGGCGGTGA CGGCGGTGAC 290

(2) INFORMATION FOR SEQ ID NO:38:

- (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 34 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:38:

GATCCAGTGG CATGGNGGGT GTCAGTGGAA GCAT 34

(2) INFORMATION FOR SEQ ID NO:39:

- (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 155 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:39:

GATCGCTGCT CGTCCCCCCC TTGCCGCCGA CGCCACCGGT CCCACCGTTA CCGAACAAGC 60
TGGCGTGGTC GCCAGACCCC CCGGCACCGC CGACGCCGGA GTCGAACAAT GGCACCGTCG 120
TATCCCCACC ATTGCCGCCG GNCCCACCGG CACCG 155

(2) INFORMATION FOR SEQ ID NO:40:

- (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 53 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:40:

ATGGCGTTCA CGGGGCGCCG GGGACCGGGC AGCCCGGNGG GGCCGGGGGG TGG 53

(2) INFORMATION FOR SEQ ID NO:41:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 132 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:41:

GATCCACCGC GGGTGCAGAC GGTGCCCCGCG GCGCCACCCC GACCAGCGGC GGCAACGGCG 60
GCACCGGCGG CAACGGCGCG AACGCCACCG TCGTCGGNGG GGCCGGCGGG GCCGGCGGCA 120
AGGGCGGCAA CG 132

(2) INFORMATION FOR SEQ ID NO:42:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 132 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:42:

GATCGGCGGC CGGNACGGNC GGGGACGGCG GCAAGGGCGG NAACGGGGGC GCCGNAGCCA 60
CCNGCCAAGA ATCCTCCGNG TCCNCCAATG GCGCGAATGG CGGACAGGGC GGCAACGGCG 120
GCANCGGCGG CA 132

(2) INFORMATION FOR SEQ ID NO:43:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 702 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:43:

```
CGGCACGAGG ATCGGTACCC CGCGGCATCG GCAGCTGCCG ATTCGCCGGG TTTCCCCACC      60
CGAGGAAAGC CGCTACCAGA TGGCGCTGCC GAAGTAGGGC GATCCGTTTCG CGATGCCGGC      120
ATGAACGGGC GGCATCAAAT TAGTGCAGGA ACCTTTCAGT TTAGCGACGA TAATGGCTAT      180
AGCACTAAGG AGGATGATCC GATATGACGC AGTCGCAGAC CGTGACGGTG GATCAGCAAG      240
AGATTTTGAA CAGGGCCAAC GAGGTGGAGG CCCCgatggc GGACCCACCG ACTGATGTCC      300
CCATCACACC GTGCGAACTC ACGGNGGNTA AAAACGCCGC CCAACAGNTG GTNTTGTCCG      360
CCGACAACAT GCGGGAATAC CTGGCGGCCG GTGCCAAAGA GCGGCAGCGT CTGGCGACCT      420
CGCTGCGCAA CGCGGCCAAG GNGTATGGCG AGGTTGATGA GGAGGCTGCG ACCGCGCTGG      480
ACAACGACGG CGAAGGAACT GTGCAGGCAG AATCGGCCGG GGCCGTCGGA GGGGACAGTT      540
CGGCCGAACT AACCgATACG CCGAGGGTGG CCACGGCCGG TGAACCCAAC TTCATGGATC      600
TCAAAGAAGC GGCAAGGAAG CTCGAAACGG GCGACCAAGG CGCATCGCTC GCGCACTGNG      660
GGGATGGGTG GAACACTTNC ACCCTGACGC TGCAAGGCGA CG                          702
```

(2) INFORMATION FOR SEQ ID NO:44:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 298 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:44:

```
GAAGCCGCAG CGCTGTCGGG CGACGTGGCG GTCAAAGCGG CATCGCTCGG TGGCGGTGGA      60
GGCGGCGGGG TGCCGTCGGC GCCGTTGGGA TCCGCGATCG GGGCGCCGA ATCGGTGCGG      120
CCCCTGGCG CTGGTGACAT TGCCGGCTTA GGCCAGGGAA GGGCCGGCGG CGGCGCCGCG      180
```

CTGGGCGGCG GTGGCATGGG AATGCCGATG GGTGCCGCGC ATCAGGGACA AGGGGGCGCC 240
 AAGTCCAAGG GTTCTCAGCA GGAAGACGAG GCGCTCTACA CCGAGGATCC TCGTGCCG 298

(2) INFORMATION FOR SEQ ID NO:45:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 1058 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:45:

CGGCACGAGG ATCGAATCGC GTCGCCGGGA GCACAGCGTC GCACTGCACC AGTGGAGGAG 60
 CCATGACCTA CTCGCCGGGT AACCCCGGAT ACCCGCAAGC GCAGCCCGCA GGCTCCTACG 120
 GAGGCGTCAC ACCCTCGTTC GCCCAGCCG ATGAGGGTGC GAGCAAGCTA CCGATGTACC 180
 TGAACATCGC GGTGGCAGTG CTCGGTCTGG CTGCGTACTT CGCCAGCTTC GGCCCAATGT 240
 TCACCCTCAG TACCGAACTC GGGGGGGGTG ATGGCGCAGT GTCCGGTGAC ACTGGGCTGC 300
 CGGTCGGGGT GGCTCTGCTG GCTGCGCTGC TTGCCGGGGT GGTCTGGTG CCTAAGGCCA 360
 AGAGCCATGT GACGGTAGTT GCGGTGCTCG GGGTACTCGG CGTATTTCTG ATGGTCTCGG 420
 CGACGTTTAA CAAGCCCAGC GCCTATTCGA CCGGTTGGGC ATTGTGGGTT GTGTTGGCTT 480
 TCATCGTGTT CCAGGCGGTT GCGGCAGTCC TGGCGCTCTT GGTGGAGACC GGCCTATCA 540
 CCGCGCCGGC GCCGCGGCCC AAGTTCGACC CGTATGGACA GTACGGGCGG TACGGGCAGT 600
 ACGGGCAGTA CGGGGTGCAG CCGGGTGGGT ACTACGGTCA GCAGGGTGCT CAGCAGGCCG 660
 CGGGACTGCA GTCGCCCGGC CCGCAGCAGT CTCCGCAGCC TCCCGGATAT GGGTCGCACT 720
 ACGGCGGCTA TTCGTCCAGT CCGAGCCAAT CGGGCAGTGG ATACACTGCT CAGCCCCCGG 780
 CCCAGCCGCC GGCAGTCC GGGTCGCAAC AATCGCACCA GGGCCCATCC ACGCCACCTA 840
 CCGGCTTTCC GAGCTTCAGC CCACCACCAC CGGTCAGTGC CGGGACGGGG TCGCAGGCTG 900
 GTTCGGCTCC AGTCAACTAT TCAAACCCCA GCGGGGGCGA GCAGTCGTCG TCCCCGGGG 960
 GGGCGCCGGT CTAACCGGGC GTTCCGCGT CCGGTCGCGC GTGTGCGCGA AGAGTGAACA 1020
 GGGTGTACG AAGCGCGGAC GATCCTCGTG CCGAATTC 1058

(2) INFORMATION FOR SEQ ID NO:46:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 327 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:46:

CGGCACGAGA GACCGATGCC GCTACCCTCG CGCAGGAGGC AGGTAATTTC GAGCGGATCT	60
CCGGCGACCT GAAAACCCAG ATCGACCAGG TGGAGTCGAC GGCAGGTTTC TTGCAGGGCC	120
AGTGGCGCGG CGCGGCGGGG ACGGCCGCCC AGGCCGCGGT GGTGCGCTTC CAAGAAGCAG	180
CCAATAAGCA GAAGCAGGAA CTCGACGAGA TCTCGACGAA TATTCGTCAG GCCGGCGTCC	240
AATACTCGAG GGCCGACGAG GAGCAGCAGC AGGCGCTGTC CTCGCAAATG GGCTTCTGAC	300
CCGCTAATAC GAAAAGAAAC GGAGCAA	327

(2) INFORMATION FOR SEQ ID NO:47:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 170 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:47:

CGGTGCGGAT GATGGCGTTG TCGAACGTGA CCGATTCTGT ACCGCCGTCG TTGAGATCAA	60
CCAACAACGT GTTGGCGTCG GCAAATGTGC CGNACCCGTG GATCTCGGTG ATCTTGTCT	120
TCTTCATCAG GAAGTGCACA CCGGCCACCC TGCCCTCGGN TACCTTTCGG	170

(2) INFORMATION FOR SEQ ID NO:48:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 127 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:48:

GATCCGGCGG CACGGGGGGT GCCGGCGGCA GCACCGCTGG CGCTGGCGGC AACGGCGGGG 60
CCGGGGGTGG CGGCGGAACC GGTGGGTTC TCTTCGGCAA CGGCGGTGCC GGCGGGCACG 120
GGGCCGT 127

(2) INFORMATION FOR SEQ ID NO:49:

- (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 81 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:49:

CGGCGGCAAG GGCGGCACCG CCGGCAACGG GAGCGGCGCG GCCGGCGGCA ACGGCGGCAA 60
CGGCGGCTCC GGCCTCAACG G 81

(2) INFORMATION FOR SEQ ID NO:50:

- (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 149 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:50:

GATCAGGGCT GGCCGGCTCC GGCCAGAAGG GCGGTAACGG AGGAGCTGCC GGATTGTTTG 60
GCAACGGCGG GGCCGGNGGT GCCGGCGCGT CCAACCAAGC CGGTAACGGC GGNGCCGGCG 120
GAAACGGTGG TGCCGGTGGG CTGATCTGG 149

(2) INFORMATION FOR SEQ ID NO:51:

- (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 355 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single

(D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:51:

CGGCACGAGA TCACACCTAC CGAGTGATCG AGATCGTCGG GACCTCGCCC GACGGTGTCTG	60
ACGCGGNAAT CCAGGGCGGT CTGGCCCGAG CTGCGCAGAC CATGCGCGCG CTGGACTGGT	120
TCGAAGTACA GTCAATTCGA GGCCACCTGG TCGACGGAGC GGTCGCGCAC TTCCAGGTGA	180
CTATGAAAGT CGGCTTCCGC CTGGAGGATT CCTGAACCTT CAAGCGCGGC CGATAACTGA	240
GGTGCATCAT TAAGCGACTT TTCCAGAACA TCCTGACGCG CTCGAAACGC GGTTCAGCCG	300
ACGGTGGCTC CGCCGAGGCG CTGCCTCCAA AATCCCTGCG ACAATTCGTC GGCGG	355

(2) INFORMATION FOR SEQ ID NO:52:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 999 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:52:

ATGCATCACC ATCACCATCA CATGCATCAG GTGGACCCCA ACTTGACACG TCGCAAGGGA	60
CGATTGGCGG CACTGGCTAT CGCGGCGATG GCCAGCGCCA GCCTGGTGAC CGTTGCGGTG	120
CCC CGACCG CCAACGCCGA TCCGGAGCCA GCGCCCCCGG TACCCACAAC GGCCGCCTCG	180
CCGCCGTCGA CCGCTGCAGC GCCACCCGCA CCGGCGACAC CTGTTGCCCC CCCACCACCG	240
GCCGCCGCCA ACACGCCGAA TGCCCAGCCG GGCGATCCCA ACGCAGCACC TCCGCCGGCC	300
GACCCGAACG CACCGCCGCC ACCTGTCATT GCCCCAAACG CACCCCAACC TGTCCGGATC	360
GACAACCCGG TTGGAGGATT CAGCTTCGCG CTGCCTGCTG GCTGGGTGGA GTCTGACGCC	420
GCCCATTTCG ACTACGGTTC AGCACTCCTC AGCAAAACCA CCGGGGACCC GCCATTTCCTC	480
GGACAGCCGC CGCCGGTGGC CAATGACACC CGTATCGTGC TCGGCCGGCT AGACCAAAAG	540
CTTTACGCCA GCGCCGAAGC CACCGACTCC AAGGCCGCGG CCCGGTTGGG CTCGGACATG	600
GGTGAGTTCT ATATGCCCTA CCCGGGCACC CGGATCAACC AGGAAACCGT CTCGCTCGAC	660

GCCAACGGGG TGTCTGGAAG CGCGTCGTAT TACGAAGTCA AGTTCAGCGA TCCGAGTAAG 720
 CCGAACGGCC AGATCTGGAC GGGCGTAATC GGCTCGCCCG CGGCGAACGC ACCGGACGCC 780
 GGGCCCCCTC AGCGCTGGTT TGTGGTATGG CTCGGGACCG CCAACAACCC GGTGGACAAG 840
 GGC GCGGCCA AGGCGCTGGC CGAATCGATC CGGCCTTTGG TCGCCCCGCC GCCGGCGCCG 900
 GCACCGGCTC CTGCAGAGCC CGCTCCGGCG CCGGCGCCGG CCGGGGAAGT CGCTCCTACC 960
 CCGACGACAC CGACACCGCA GCGGACCTTA CCGGCCTGA 999

(2) INFORMATION FOR SEQ ID NO:53:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 332 amino acids
- (B) TYPE: amino acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:53:

Met His His His His His His Met His Gln Val Asp Pro Asn Leu Thr
 1 5 10 15
 Arg Arg Lys Gly Arg Leu Ala Ala Leu Ala Ile Ala Ala Met Ala Ser
 20 25 30
 Ala Ser Leu Val Thr Val Ala Val Pro Ala Thr Ala Asn Ala Asp Pro
 35 40 45
 Glu Pro Ala Pro Pro Val Pro Thr Thr Ala Ala Ser Pro Pro Ser Thr
 50 55 60
 Ala Ala Ala Pro Pro Ala Pro Ala Thr Pro Val Ala Pro Pro Pro Pro
 65 70 75 80
 Ala Ala Ala Asn Thr Pro Asn Ala Gln Pro Gly Asp Pro Asn Ala Ala
 85 90 95
 Pro Pro Pro Ala Asp Pro Asn Ala Pro Pro Pro Pro Val Ile Ala Pro
 100 105 110
 Asn Ala Pro Gln Pro Val Arg Ile Asp Asn Pro Val Gly Gly Phe Ser
 115 120 125
 Phe Ala Leu Pro Ala Gly Trp Val Glu Ser Asp Ala Ala His Phe Asp
 130 135 140
 Tyr Gly Ser Ala Leu Leu Ser Lys Thr Thr Gly Asp Pro Pro Phe Pro

[illegible]

(2) INFORMATION FOR SEQ ID NO:54:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 20 amino acids
(B) TYPE: amino acid
(C) STRANDEDNESS:
(D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:54:

```

Asp Pro Val Asp Ala Val Ile Asn Thr Thr Xaa Asn Tyr Gly Gln Val
1          5          10          15
Val Ala Ala Leu
          20

```

WO 98/16645

95

(2) INFORMATION FOR SEQ ID NO:55:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 15 amino acids
 (B) TYPE: amino acid
 (C) STRANDEDNESS:
 (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:55:

Ala Val Glu Ser Gly Met Leu Ala Leu Gly Thr Pro Ala Pro Ser
 1 5 10 15

(2) INFORMATION FOR SEQ ID NO:56:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 19 amino acids
 (B) TYPE: amino acid
 (C) STRANDEDNESS:
 (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:56:

Ala Ala Met Lys Pro Arg Thr Gly Asp Gly Pro Leu Glu Ala Ala Lys
 1 5 10 15
 Glu Gly Arg

(2) INFORMATION FOR SEQ ID NO:57:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 15 amino acids
 (B) TYPE: amino acid
 (C) STRANDEDNESS:
 (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:57:

Tyr Tyr Trp Cys Pro Gly Gln Pro Phe Asp Pro Ala Trp Gly Pro
 1 5 10 15

(2) INFORMATION FOR SEQ ID NO:58:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 14 amino acids
- (B) TYPE: amino acid
- (C) STRANDEDNESS:
- (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:58:

Asp Ile Gly Ser Glu Ser Thr Glu Asp Gln Gln Xaa Ala Val
1 5 10

(2) INFORMATION FOR SEQ ID NO:59:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 13 amino acids
- (B) TYPE: amino acid
- (C) STRANDEDNESS:
- (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:59:

Ala Glu Glu Ser Ile Ser Thr Xaa Glu Xaa Ile Val Pro
1 5 10

(2) INFORMATION FOR SEQ ID NO:60:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 17 amino acids
- (B) TYPE: amino acid
- (C) STRANDEDNESS:
- (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:60:

Asp Pro Glu Pro Ala Pro Pro Val Pro Thr Ala Ala Ala Ala Pro Pro
1 5 10 15

Ala

(2) INFORMATION FOR SEQ ID NO:61:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 15 amino acids
 (B) TYPE: amino acid
 (C) STRANDEDNESS:
 (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:61:

Ala Pro Lys Thr Tyr Xaa Glu Glu Leu Lys Gly Thr Asp Thr Gly
1 5 10 15

(2) INFORMATION FOR SEQ ID NO:62:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 30 amino acids
 (B) TYPE: amino acid
 (C) STRANDEDNESS:
 (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:62:

Asp Pro Ala Ser Ala Pro Asp Val Pro Thr Ala Ala Gln Gln Thr Ser
1 5 10 15

Leu Leu Asn Asn Leu Ala Asp Pro Asp Val Ser Phe Ala Asp
 20 25 30

(2) INFORMATION FOR SEQ ID NO:63:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 24 amino acids
 (B) TYPE: amino acid
 (C) STRANDEDNESS:
 (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:63:

Gly Cys Gly Asp Arg Ser Gly Gly Asn Leu Asp Gln Ile Arg Leu Arg
1 5 10 15

Arg Asp Arg Ser Gly Gly Asn Leu
 20

(2) INFORMATION FOR SEQ ID NO:64:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 187 amino acids
- (B) TYPE: amino acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:64:

```

Thr Gly Ser Leu Asn Gln Thr His Asn Arg Arg Ala Asn Glu Arg Lys
1          5          10          15

Asn Thr Thr Met Lys Met Val Lys Ser Ile Ala Ala Gly Leu Thr Ala
          20          25          30

Ala Ala Ala Ile Gly Ala Ala Ala Ala Gly Val Thr Ser Ile Met Ala
          35          40          45

Gly Gly Pro Val Val Tyr Gln Met Gln Pro Val Val Phe Gly Ala Pro
50          55          60

Leu Pro Leu Asp Pro Ala Ser Ala Pro Asp Val Pro Thr Ala Ala Gln
65          70          75          80

Leu Thr Ser Leu Leu Asn Ser Leu Ala Asp Pro Asn Val Ser Phe Ala
          85          90          95

Asn Lys Gly Ser Leu Val Glu Gly Gly Ile Gly Gly Thr Glu Ala Arg
          100          105          110

Ile Ala Asp His Lys Leu Lys Lys Ala Ala Glu His Gly Asp Leu Pro
          115          120          125

Leu Ser Phe Ser Val Thr Asn Ile Gln Pro Ala Ala Ala Gly Ser Ala
          130          135          140

Thr Ala Asp Val Ser Val Ser Gly Pro Lys Leu Ser Ser Pro Val Thr
145          150          155          160

Gln Asn Val Thr Phe Val Asn Gln Gly Gly Trp Met Leu Ser Arg Ala
          165          170          175

Ser Ala Met Glu Leu Leu Gln Ala Ala Gly Xaa
          180          185

```

(2) INFORMATION FOR SEQ ID NO:65:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 148 amino acids
- (B) TYPE: amino acid

- (C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:65:

```

Asp Glu Val Thr Val Glu Thr Thr Ser Val Phe Arg Ala Asp Phe Leu
1           5           10           15

Ser Glu Leu Asp Ala Pro Ala Gln Ala Gly Thr Glu Ser Ala Val Ser
          20           25           30

Gly Val Glu Gly Leu Pro Pro Gly Ser Ala Leu Leu Val Val Lys Arg
          35           40           45

Gly Pro Asn Ala Gly Ser Arg Phe Leu Leu Asp Gln Ala Ile Thr Ser
          50           55           60

Ala Gly Arg His Pro Asp Ser Asp Ile Phe Leu Asp Asp Val Thr Val
65           70           75           80

Ser Arg Arg His Ala Glu Phe Arg Leu Glu Asn Asn Glu Phe Asn Val
          85           90           95

Val Asp Val Gly Ser Leu Asn Gly Thr Tyr Val Asn Arg Glu Pro Val
          100          105          110

Asp Ser Ala Val Leu Ala Asn Gly Asp Glu Val Gln Ile Gly Lys Leu
          115          120          125

Arg Leu Val Phe Leu Thr Gly Pro Lys Gln Gly Glu Asp Asp Gly Ser
          130          135          140

Thr Gly Gly Pro
145

```

(2) INFORMATION FOR SEQ ID NO:66:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 230 amino acids
 (B) TYPE: amino acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:66:

```

Thr Ser Asn Arg Pro Ala Arg Arg Gly Arg Arg Ala Pro Arg Asp Thr
1           5           10           15

```

100

Gly Pro Asp Arg Ser Ala Ser Leu Ser Leu Val Arg His Arg Arg Gln
 20 25 30
 Gln Arg Asp Ala Leu Cys Leu Ser Ser Thr Gln Ile Ser Arg Gln Ser
 35 40 45
 Asn Leu Pro Pro Ala Ala Gly Gly Ala Ala Asn Tyr Ser Arg Arg Asn
 50 55 60
 Phe Asp Val Arg Ile Lys Ile Phe Met Leu Val Thr Ala Val Val Leu
 65 70 75 80
 Leu Cys Cys Ser Gly Val Ala Thr Ala Ala Pro Lys Thr Tyr Cys Glu
 85 90 95
 Glu Leu Lys Gly Thr Asp Thr Gly Gln Ala Cys Gln Ile Gln Met Ser
 100 105 110
 Asp Pro Ala Tyr Asn Ile Asn Ile Ser Leu Pro Ser Tyr Tyr Pro Asp
 115 120 125
 Gln Lys Ser Leu Glu Asn Tyr Ile Ala Gln Thr Arg Asp Lys Phe Leu
 130 135 140
 Ser Ala Ala Thr Ser Ser Thr Pro Arg Glu Ala Pro Tyr Glu Leu Asn
 145 150 155 160
 Ile Thr Ser Ala Thr Tyr Gln Ser Ala Ile Pro Pro Arg Gly Thr Gln
 165 170 175
 Ala Val Val Leu Xaa Val Tyr His Asn Ala Gly Gly Thr His Pro Thr
 180 185 190
 Thr Thr Tyr Lys Ala Phe Asp Trp Asp Gln Ala Tyr Arg Lys Pro Ile
 195 200 205
 Thr Tyr Asp Thr Leu Trp Gln Ala Asp Thr Asp Pro Leu Pro Val Val
 210 215 220
 Phe Pro Ile Val Ala Arg
 225 230

(2) INFORMATION FOR SEQ ID NO:67:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 132 amino acids
- (B) TYPE: amino acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:67:

Thr Ala Ala Ser Asp Asn Phe Gln Leu Ser Gln Gly Gly Gln Gly Phe
 1 5 10 15
 Ala Ile Pro Ile Gly Gln Ala Met Ala Ile Ala Gly Gln Ile Arg Ser
 20 25 30
 Gly Gly Gly Ser Pro Thr Val His Ile Gly Pro Thr Ala Phe Leu Gly
 35 40 45
 Leu Gly Val Val Asp Asn Asn Gly Asn Gly Ala Arg Val Gln Arg Val
 50 55 60
 Val Gly Ser Ala Pro Ala Ala Ser Leu Gly Ile Ser Thr Gly Asp Val
 65 70 75 80
 Ile Thr Ala Val Asp Gly Ala Pro Ile Asn Ser Ala Thr Ala Met Ala
 85 90 95
 Asp Ala Leu Asn Gly His His Pro Gly Asp Val Ile Ser Val Asn Trp
 100 105 110
 Gln Thr Lys Ser Gly Gly Thr Arg Thr Gly Asn Val Thr Leu Ala Glu
 115 120 125
 Gly Pro Pro Ala
 130

(2) INFORMATION FOR SEQ ID NO:68:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 100 amino acids
- (B) TYPE: amino acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:68:

Val Pro Leu Arg Ser Pro Ser Met Ser Pro Ser Lys Cys Leu Ala Ala
 1 5 10 15
 Ala Gln Arg Asn Pro Val Ile Arg Arg Arg Arg Leu Ser Asn Pro Pro
 20 25 30
 Pro Arg Lys Tyr Arg Ser Met Pro Ser Pro Ala Thr Ala Ser Ala Gly
 35 40 45
 Met Ala Arg Val Arg Arg Arg Ala Ile Trp Arg Gly Pro Ala Thr Xaa
 50 55 60

Ser Ala Gly Met Ala Arg Val Arg Arg Trp Xaa Val Met Pro Xaa Val
 65 70 75 80

Ile Gln Ser Thr Xaa Ile Arg Xaa Xaa Gly Pro Phe Asp Asn Arg Gly
 85 90 95

Ser Glu Arg Lys
 100

(2) INFORMATION FOR SEQ ID NO:69:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 163 amino acids
- (B) TYPE: amino acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:69:

Met Thr Asp Asp Ile Leu Leu Ile Asp Thr Asp Glu Arg Val Arg Thr
 1 5 10 15

Leu Thr Leu Asn Arg Pro Gln Ser Arg Asn Ala Leu Ser Ala Ala Leu
 20 25 30

Arg Asp Arg Phe Phe Ala Xaa Leu Xaa Asp Ala Glu Xaa Asp Asp Asp
 35 40 45

Ile Asp Val Val Ile Leu Thr Gly Ala Asp Pro Val Phe Cys Ala Gly
 50 55 60

Leu Asp Leu Lys Val Ala Gly Arg Ala Asp Arg Ala Ala Gly His Leu
 65 70 75 80

Thr Ala Val Gly Gly His Asp Gln Ala Gly Asp Arg Arg Asp Gln Arg
 85 90 95

Arg Arg Gly His Arg Arg Ala Arg Thr Gly Ala Val Leu Arg His Pro
 100 105 110

Asp Arg Leu Arg Ala Arg Pro Leu Arg Arg His Pro Arg Pro Gly Gly
 115 120 125

Ala Ala Ala His Leu Gly Thr Gln Cys Val Leu Ala Ala Lys Gly Arg
 130 135 140

His Arg Xaa Gly Pro Val Asp Glu Pro Asp Arg Arg Leu Pro Val Arg
 145 150 155 160

Asp Arg Arg

(2) INFORMATION FOR SEQ ID NO:70:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 344 amino acids
- (B) TYPE: amino acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:70:

```

Met Lys Phe Val Asn His Ile Glu Pro Val Ala Pro Arg Arg Ala Gly
 1             5             10             15

Gly Ala Val Ala Glu Val Tyr Ala Glu Ala Arg Arg Glu Phe Gly Arg
      20             25             30

Leu Pro Glu Pro Leu Ala Met Leu Ser Pro Asp Glu Gly Leu Leu Thr
      35             40             45

Ala Gly Trp Ala Thr Leu Arg Glu Thr Leu Leu Val Gly Gln Val Pro
      50             55             60

Arg Gly Arg Lys Glu Ala Val Ala Ala Ala Val Ala Ala Ser Leu Arg
      65             70             75             80

Cys Pro Trp Cys Val Asp Ala His Thr Thr Met Leu Tyr Ala Ala Gly
      85             90             95

Gln Thr Asp Thr Ala Ala Ala Ile Leu Ala Gly Thr Ala Pro Ala Ala
      100             105             110

Gly Asp Pro Asn Ala Pro Tyr Val Ala Trp Ala Ala Gly Thr Gly Thr
      115             120             125

Pro Ala Gly Pro Pro Ala Pro Phe Gly Pro Asp Val Ala Ala Glu Tyr
      130             135             140

Leu Gly Thr Ala Val Gln Phe His Phe Ile Ala Arg Leu Val Leu Val
      145             150             155             160

Leu Leu Asp Glu Thr Phe Leu Pro Gly Gly Pro Arg Ala Gln Gln Leu
      165             170             175

Met Arg Arg Ala Gly Gly Leu Val Phe Ala Arg Lys Val Arg Ala Glu
      180             185             190

His Arg Pro Gly Arg Ser Thr Arg Arg Leu Glu Pro Arg Thr Leu Pro
      195             200             205

Asp Asp Leu Ala Trp Ala Thr Pro Ser Glu Pro Ile Ala Thr Ala Phe

```

104

210	215	220
Ala Ala Leu Ser His His Leu Asp Thr Ala Pro His Leu Pro Pro Pro		
225	230	235 240
Thr Arg Gln Val Val Arg Arg Val Val Gly Ser Trp His Gly Glu Pro		
	245	250 255
Met Pro Met Ser Ser Arg Trp Thr Asn Glu His Thr Ala Glu Leu Pro		
	260	265 270
Ala Asp Leu His Ala Pro Thr Arg Leu Ala Leu Leu Thr Gly Leu Ala		
	275	280 285
Pro His Gln Val Thr Asp Asp Asp Val Ala Ala Ala Arg Ser Leu Leu		
	290	295 300
Asp Thr Asp Ala Ala Leu Val Gly Ala Leu Ala Trp Ala Ala Phe Thr		
305	310	315 320
Ala Ala Arg Arg Ile Gly Thr Trp Ile Gly Ala Ala Ala Glu Gly Gln		
	325	330 335
Val Ser Arg Gln Asn Pro Thr Gly		
	340	

(2) INFORMATION FOR SEQ ID NO:71:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 485 amino acids
- (B) TYPE: amino acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:71:

Asp Asp Pro Asp Met Pro Gly Thr Val Ala Lys Ala Val Ala Asp Ala	
1	5 10 15
Leu Gly Arg Gly Ile Ala Pro Val Glu Asp Ile Gln Asp Cys Val Glu	
	20 25 30
Ala Arg Leu Gly Glu Ala Gly Leu Asp Asp Val Ala Arg Val Tyr Ile	
	35 40 45
Ile Tyr Arg Gln Arg Arg Ala Glu Leu Arg Thr Ala Lys Ala Leu Leu	
	50 55 60
Gly Val Arg Asp Glu Leu Lys Leu Ser Leu Ala Ala Val Thr Val Leu	
65	70 75 80

Arg Glu Arg Tyr Leu Leu His Asp Glu Gln Gly Arg Pro Ala Glu Ser
 85 90 95
 Thr Gly Glu Leu Met Asp Arg Ser Ala Arg Cys Val Ala Ala Glu
 100 105 110
 Asp Gln Tyr Glu Pro Gly Ser Ser Arg Arg Trp Ala Glu Arg Phe Ala
 115 120 125
 Thr Leu Leu Arg Asn Leu Glu Phe Leu Pro Asn Ser Pro Thr Leu Met
 130 135 140
 Asn Ser Gly Thr Asp Leu Gly Leu Leu Ala Gly Cys Phe Val Leu Pro
 145 150 155 160
 Ile Glu Asp Ser Leu Gln Ser Ile Phe Ala Thr Leu Gly Gln Ala Ala
 165 170 175
 Glu Leu Gln Arg Ala Gly Gly Gly Thr Gly Tyr Ala Phe Ser His Leu
 180 185 190
 Arg Pro Ala Gly Asp Arg Val Ala Ser Thr Gly Gly Thr Ala Ser Gly
 195 200 205
 Pro Val Ser Phe Leu Arg Leu Tyr Asp Ser Ala Ala Gly Val Val Ser
 210 215 220
 Met Gly Gly Arg Arg Arg Gly Ala Cys Met Ala Val Leu Asp Val Ser
 225 230 235 240
 His Pro Asp Ile Cys Asp Phe Val Thr Ala Lys Ala Glu Ser Pro Ser
 245 250 255
 Glu Leu Pro His Phe Asn Leu Ser Val Gly Val Thr Asp Ala Phe Leu
 260 265 270
 Arg Ala Val Glu Arg Asn Gly Leu His Arg Leu Val Asn Pro Arg Thr
 275 280 285
 Gly Lys Ile Val Ala Arg Met Pro Ala Ala Glu Leu Phe Asp Ala Ile
 290 295 300
 Cys Lys Ala Ala His Ala Gly Gly Asp Pro Gly Leu Val Phe Leu Asp
 305 310 315 320
 Thr Ile Asn Arg Ala Asn Pro Val Pro Gly Arg Gly Arg Ile Glu Ala
 325 330 335
 Thr Asn Pro Cys Gly Glu Val Pro Leu Leu Pro Tyr Glu Ser Cys Asn
 340 345 350
 Leu Gly Ser Ile Asn Leu Ala Arg Met Leu Ala Asp Gly Arg Val Asp
 355 360 365
 Trp Asp Arg Leu Glu Glu Val Ala Gly Val Ala Val Arg Phe Leu Asp

370		375		380
Asp Val Ile Asp Val Ser Arg Tyr Pro Phe Pro Glu Leu Gly Glu Ala				
385		390		395
Ala Arg Ala Thr Arg Lys Ile Gly Leu Gly Val Met Gly Leu Ala Glu				
	405		410	415
Leu Leu Ala Ala Leu Gly Ile Pro Tyr Asp Ser Glu Glu Ala Val Arg				
	420		425	430
Leu Ala Thr Arg Leu Met Arg Arg Ile Gln Gln Ala Ala His Thr Ala				
	435		440	445
Ser Arg Arg Leu Ala Glu Glu Arg Gly Ala Phe Pro Ala Phe Thr Asp				
	450		455	460
Ser Arg Phe Ala Arg Ser Gly Pro Arg Arg Asn Ala Gln Val Thr Ser				
	465		470	475
Val Ala Pro Thr Gly				
	485			

(2) INFORMATION FOR SEQ ID NO:72:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 267 amino acids
- (B) TYPE: amino acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:72:

Gly Val Ile Val Leu Asp Leu Glu Pro Arg Gly Pro Leu Pro Thr Glu			
1	5	10	15
Ile Tyr Trp Arg Arg Arg Gly Leu Ala Leu Gly Ile Ala Val Val Val			
	20	25	30
Val Gly Ile Ala Val Ala Ile Val Ile Ala Phe Val Asp Ser Ser Ala			
	35	40	45
Gly Ala Lys Pro Val Ser Ala Asp Lys Pro Ala Ser Ala Gln Ser His			
	50	55	60
Pro Gly Ser Pro Ala Pro Gln Ala Pro Gln Pro Ala Gly Gln Thr Glu			
	65	70	75
Gly Asn Ala Ala Ala Ala Pro Pro Gln Gly Gln Asn Pro Glu Thr Pro			
	85	90	95

107

```

Thr Pro Thr Ala Ala Val Gln Pro Pro Pro Val Leu Lys Glu Gly Asp
      100                      105                      110

Asp Cys Pro Asp Ser Thr Leu Ala Val Lys Gly Leu Thr Asn Ala Pro
      115                      120                      125

Gln Tyr Tyr Val Gly Asp Gln Pro Lys Phe Thr Met Val Val Thr Asn
      130                      135                      140

Ile Gly Leu Val Ser Cys Lys Arg Asp Val Gly Ala Ala Val Leu Ala
      145                      150                      155                      160

Ala Tyr Val Tyr Ser Leu Asp Asn Lys Arg Leu Trp Ser Asn Leu Asp
      165                      170                      175

Cys Ala Pro Ser Asn Glu Thr Leu Val Lys Thr Phe Ser Pro Gly Glu
      180                      185                      190

Gln Val Thr Thr Ala Val Thr Trp Thr Gly Met Gly Ser Ala Pro Arg
      195                      200                      205

Cys Pro Leu Pro Arg Pro Ala Ile Gly Pro Gly Thr Tyr Asn Leu Val
      210                      215                      220

Val Gln Leu Gly Asn Leu Arg Ser Leu Pro Val Pro Phe Ile Leu Asn
      225                      230                      235                      240

Gln Pro Pro Pro Pro Pro Gly Pro Val Pro Ala Pro Gly Pro Ala Gln
      245                      250                      255

Ala Pro Pro Pro Glu Ser Pro Ala Gln Gly Gly
      260                      265

```

(2) INFORMATION FOR SEQ ID NO:73:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 97 amino acids
- (B) TYPE: amino acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:73:

```

Leu Ile Ser Thr Gly Lys Ala Ser His Ala Ser Leu Gly Val Gln Val
1           5           10           15

Thr Asn Asp Lys Asp Thr Pro Gly Ala Lys Ile Val Glu Val Val Ala
      20           25           30

Gly Gly Ala Ala Ala Asn Ala Gly Val Pro Lys Gly Val Val Val Thr
      35           40           45

```

Lys Val Asp Asp Arg Pro Ile Asn Ser Ala Asp Ala Leu Val Ala Ala
 50 55 60
 Val Arg Ser Lys Ala Pro Gly Ala Thr Val Ala Leu Thr Phe Gln Asp
 65 70 75 80
 Pro Ser Gly Gly Ser Arg Thr Val Gln Val Thr Leu Gly Lys Ala Glu
 85 90 95
 Gln

(2) INFORMATION FOR SEQ ID NO:74:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 364 amino acids
- (B) TYPE: amino acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:74:

Gly Ala Ala Val Ser Leu Leu Ala Ala Gly Thr Leu Val Leu Thr Ala
 1 5 10 15
 Cys Gly Gly Gly Thr Asn Ser Ser Ser Ser Gly Ala Gly Gly Thr Ser
 20 25 30
 Gly Ser Val His Cys Gly Gly Lys Lys Glu Leu His Ser Ser Gly Ser
 35 40 45
 Thr Ala Gln Glu Asn Ala Met Glu Gln Phe Val Tyr Ala Tyr Val Arg
 50 55 60
 Ser Cys Pro Gly Tyr Thr Leu Asp Tyr Asn Ala Asn Gly Ser Gly Ala
 65 70 75 80
 Gly Val Thr Gln Phe Leu Asn Asn Glu Thr Asp Phe Ala Gly Ser Asp
 85 90 95
 Val Pro Leu Asn Pro Ser Thr Gly Gln Pro Asp Arg Ser Ala Glu Arg
 100 105 110
 Cys Gly Ser Pro Ala Trp Asp Leu Pro Thr Val Phe Gly Pro Ile Ala
 115 120 125
 Ile Thr Tyr Asn Ile Lys Gly Val Ser Thr Leu Asn Leu Asp Gly Pro
 130 135 140
 Thr Thr Ala Lys Ile Phe Asn Gly Thr Ile Thr Val Trp Asn Asp Pro

145		150		155		160
Gln Ile Gln Ala Leu Asn Ser Gly Thr Asp Leu Pro Pro Thr Pro Ile						
		165		170		175
Ser Val Ile Phe Arg Ser Asp Lys Ser Gly Thr Ser Asp Asn Phe Gln						
		180		185		190
Lys Tyr Leu Asp Gly Val Ser Asn Gly Ala Trp Gly Lys Gly Ala Ser						
		195		200		205
Glu Thr Phe Ser Gly Gly Val Gly Val Gly Ala Ser Gly Asn Asn Gly						
		210		215		220
Thr Ser Ala Leu Leu Gln Thr Thr Asp Gly Ser Ile Thr Tyr Asn Glu						
		225		230		235
Trp Ser Phe Ala Val Gly Lys Gln Leu Asn Met Ala Gln Ile Ile Thr						
		245		250		255
Ser Ala Gly Pro Asp Pro Val Ala Ile Thr Thr Glu Ser Val Gly Lys						
		260		265		270
Thr Ile Ala Gly Ala Lys Ile Met Gly Gln Gly Asn Asp Leu Val Leu						
		275		280		285
Asp Thr Ser Ser Phe Tyr Arg Pro Thr Gln Pro Gly Ser Tyr Pro Ile						
		290		295		300
Val Leu Ala Thr Tyr Glu Ile Val Cys Ser Lys Tyr Pro Asp Ala Thr						
		305		310		315
Thr Gly Thr Ala Val Arg Ala Phe Met Gln Ala Ala Ile Gly Pro Gly						
		325		330		335
Gln Glu Gly Leu Asp Gln Tyr Gly Ser Ile Pro Leu Pro Lys Ser Phe						
		340		345		350
Gln Ala Lys Leu Ala Ala Ala Val Asn Ala Ile Ser						
		355		360		

(2) INFORMATION FOR SEQ ID NO:75:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 309 amino acids
- (B) TYPE: amino acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:75:

Gln Ala Ala Ala Gly Arg Ala Val Arg Arg Thr Gly His Ala Glu Asp
 1 5 10 15
 Gln Thr His Gln Asp Arg Leu His His Gly Cys Arg Arg Ala Ala Val
 20 25 30
 Val Val Arg Gln Asp Arg Ala Ser Val Ser Ala Thr Ser Ala Arg Pro
 35 40 45
 Pro Arg Arg His Pro Ala Gln Gly His Arg Arg Arg Val Ala Pro Ser
 50 55 60
 Gly Gly Arg Arg Arg Pro His Pro His His Val Gln Pro Asp Asp Arg
 65 70 75 80
 Arg Asp Arg Pro Ala Leu Leu Asp Arg Thr Gln Pro Ala Glu His Pro
 85 90 95
 Asp Pro His Arg Arg Gly Pro Ala Asp Pro Gly Arg Val Arg Gly Arg
 100 105 110
 Gly Arg Leu Arg Arg Val Asp Asp Gly Arg Leu Gln Pro Asp Arg Asp
 115 120 125
 Ala Asp His Gly Ala Pro Val Arg Gly Arg Gly Pro His Arg Gly Val
 130 135 140
 Gln His Arg Gly Gly Pro Val Phe Val Arg Arg Val Pro Gly Val Arg
 145 150 155 160
 Cys Ala His Arg Arg Gly His Arg Arg Val Ala Ala Pro Gly Gln Gly
 165 170 175
 Asp Val Leu Arg Ala Gly Leu Arg Val Glu Arg Leu Arg Pro Val Ala
 180 185 190
 Ala Val Glu Asn Leu His Arg Gly Ser Gln Arg Ala Asp Gly Arg Val
 195 200 205
 Phe Arg Pro Ile Arg Arg Gly Ala Arg Leu Pro Ala Arg Arg Ser Arg
 210 215 220
 Ala Gly Pro Gln Gly Arg Leu His Leu Asp Gly Ala Gly Pro Ser Pro
 225 230 235 240
 Leu Pro Ala Arg Ala Gly Gln Gln Gln Pro Ser Ser Ala Gly Gly Arg
 245 250 255
 Arg Ala Gly Gly Ala Glu Arg Ala Asp Pro Gly Gln Arg Gly Arg His
 260 265 270
 His Gln Gly Gly His Asp Pro Gly Arg Gln Gly Ala Gln Arg Gly Thr
 275 280 285
 Ala Gly Val Ala His Ala Ala Ala Gly Pro Arg Arg Ala Ala Val Arg

290

295

300

Asn Arg Pro Arg Arg
305

(2) INFORMATION FOR SEQ ID NO:76:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 580 amino acids
- (B) TYPE: amino acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:76:

Ser	Ala	Val	Trp	Cys	Leu	Asn	Gly	Phe	Thr	Gly	Arg	His	Arg	His	Gly	1	5	10	15
Arg	Cys	Arg	Val	Arg	Ala	Ser	Gly	Trp	Arg	Ser	Ser	Asn	Arg	Trp	Cys	20	25	30	
Ser	Thr	Thr	Ala	Asp	Cys	Cys	Ala	Ser	Lys	Thr	Pro	Thr	Gln	Ala	Ala	35	40	45	
Ser	Pro	Leu	Glu	Arg	Arg	Phe	Thr	Cys	Cys	Ser	Pro	Ala	Val	Gly	Cys	50	55	60	
Arg	Phe	Arg	Ser	Phe	Pro	Val	Arg	Arg	Leu	Ala	Leu	Gly	Ala	Arg	Thr	65	70	75	80
Ser	Arg	Thr	Leu	Gly	Val	Arg	Arg	Thr	Leu	Ser	Gln	Trp	Asn	Leu	Ser	85	90	95	
Pro	Arg	Ala	Gln	Pro	Ser	Cys	Ala	Val	Thr	Val	Glu	Ser	His	Thr	His	100	105	110	
Ala	Ser	Pro	Arg	Met	Ala	Lys	Leu	Ala	Arg	Val	Val	Gly	Leu	Val	Gln	115	120	125	
Glu	Glu	Gln	Pro	Ser	Asp	Met	Thr	Asn	His	Pro	Arg	Tyr	Ser	Pro	Pro	130	135	140	
Pro	Gln	Gln	Pro	Gly	Thr	Pro	Gly	Tyr	Ala	Gln	Gly	Gln	Gln	Gln	Thr	145	150	155	160
Tyr	Ser	Gln	Gln	Phe	Asp	Trp	Arg	Tyr	Pro	Pro	Ser	Pro	Pro	Pro	Gln	165	170	175	
Pro	Thr	Gln	Tyr	Arg	Gln	Pro	Tyr	Glu	Ala	Leu	Gly	Gly	Thr	Arg	Pro	180	185	190	

Gly Leu Ile Pro Gly Val Ile Pro Thr Met Thr Pro Pro Pro Gly Met
 195 200 205
 Val Arg Gln Arg Pro Arg Ala Gly Met Leu Ala Ile Gly Ala Val Thr
 210 215 220
 Ile Ala Val Val Ser Ala Gly Ile Gly Gly Ala Ala Ala Ser Leu Val
 225 230 235 240
 Gly Phe Asn Arg Ala Pro Ala Gly Pro Ser Gly Gly Pro Val Ala Ala
 245 250 255
 Ser Ala Ala Pro Ser Ile Pro Ala Ala Asn Met Pro Pro Gly Ser Val
 260 265 270
 Glu Gln Val Ala Ala Lys Val Val Pro Ser Val Val Met Leu Glu Thr
 275 280 285
 Asp Leu Gly Arg Gln Ser Glu Glu Gly Ser Gly Ile Ile Leu Ser Ala
 290 295 300
 Glu Gly Leu Ile Leu Thr Asn Asn His Val Ile Ala Ala Ala Ala Lys
 305 310 315 320
 Pro Pro Leu Gly Ser Pro Pro Pro Lys Thr Thr Val Thr Phe Ser Asp
 325 330 335
 Gly Arg Thr Ala Pro Phe Thr Val Val Gly Ala Asp Pro Thr Ser Asp
 340 345 350
 Ile Ala Val Val Arg Val Gln Gly Val Ser Gly Leu Thr Pro Ile Ser
 355 360 365
 Leu Gly Ser Ser Ser Asp Leu Arg Val Gly Gln Pro Val Leu Ala Ile
 370 375 380
 Gly Ser Pro Leu Gly Leu Glu Gly Thr Val Thr Thr Gly Ile Val Ser
 385 390 395 400
 Ala Leu Asn Arg Pro Val Ser Thr Thr Gly Glu Ala Gly Asn Gln Asn
 405 410 415
 Thr Val Leu Asp Ala Ile Gln Thr Asp Ala Ala Ile Asn Pro Gly Asn
 420 425 430
 Ser Gly Gly Ala Leu Val Asn Met Asn Ala Gln Leu Val Gly Val Asn
 435 440 445
 Ser Ala Ile Ala Thr Leu Gly Ala Asp Ser Ala Asp Ala Gln Ser Gly
 450 455 460
 Ser Ile Gly Leu Gly Phe Ala Ile Pro Val Asp Gln Ala Lys Arg Ile
 465 470 475 480
 Ala Asp Glu Leu Ile Ser Thr Gly Lys Ala Ser His Ala Ser Leu Gly

	485		490		495										
Val	Gln	Val	Thr	Asn	Asp	Lys	Asp	Thr	Pro	Gly	Ala	Lys	Ile	Val	Glu
			500					505					510		
Val	Val	Ala	Gly	Gly	Ala	Ala	Ala	Asn	Ala	Gly	Val	Pro	Lys	Gly	Val
		515					520					525			
Val	Val	Thr	Lys	Val	Asp	Asp	Arg	Pro	Ile	Asn	Ser	Ala	Asp	Ala	Leu
		530				535					540				
Val	Ala	Ala	Val	Arg	Ser	Lys	Ala	Pro	Gly	Ala	Thr	Val	Ala	Leu	Thr
545					550					555					560
Phe	Gln	Asp	Pro	Ser	Gly	Gly	Ser	Arg	Thr	Val	Gln	Val	Thr	Leu	Gly
				565					570					575	
Lys	Ala	Glu	Gln												
			580												

(2) INFORMATION FOR SEQ ID NO:77:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 233 amino acids
- (B) TYPE: amino acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:77:

Met	Asn	Asp	Gly	Lys	Arg	Ala	Val	Thr	Ser	Ala	Val	Leu	Val	Val	Leu
1				5					10					15	
Gly	Ala	Cys	Leu	Ala	Leu	Trp	Leu	Ser	Gly	Cys	Ser	Ser	Pro	Lys	Pro
			20					25					30		
Asp	Ala	Glu	Glu	Gln	Gly	Val	Pro	Val	Ser	Pro	Thr	Ala	Ser	Asp	Pro
		35					40					45			
Ala	Leu	Leu	Ala	Glu	Ile	Arg	Gln	Ser	Leu	Asp	Ala	Thr	Lys	Gly	Leu
	50					55					60				
Thr	Ser	Val	His	Val	Ala	Val	Arg	Thr	Thr	Gly	Lys	Val	Asp	Ser	Leu
65					70					75					80
Leu	Gly	Ile	Thr	Ser	Ala	Asp	Val	Asp	Val	Arg	Ala	Asn	Pro	Leu	Ala
				85					90					95	
Ala	Lys	Gly	Val	Cys	Thr	Tyr	Asn	Asp	Glu	Gln	Gly	Val	Pro	Phe	Arg
			100					105					110		

114

Val Gln Gly Asp Asn Ile Ser Val Lys Leu Phe Asp Asp Trp Ser Asn
 115 120 125

Leu Gly Ser Ile Ser Glu Leu Ser Thr Ser Arg Val Leu Asp Pro Ala
 130 135 140

Ala Gly Val Thr Gln Leu Leu Ser Gly Val Thr Asn Leu Gln Ala Gln
 145 150 155 160

Gly Thr Glu Val Ile Asp Gly Ile Ser Thr Thr Lys Ile Thr Gly Thr
 165 170 175

Ile Pro Ala Ser Ser Val Lys Met Leu Asp Pro Gly Ala Lys Ser Ala
 180 185 190

Arg Pro Ala Thr Val Trp Ile Ala Gln Asp Gly Ser His His Leu Val
 195 200 205

Arg Ala Ser Ile Asp Leu Gly Ser Gly Ser Ile Gln Leu Thr Gln Ser
 210 215 220

Lys Trp Asn Glu Pro Val Asn Val Asp
 225 230

(2) INFORMATION FOR SEQ ID NO:78:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 66 amino acids
- (B) TYPE: amino acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:78:

Val Ile Asp Ile Ile Gly Thr Ser Pro Thr Ser Trp Glu Gln Ala Ala
 1 5 10 15

Ala Glu Ala Val Gln Arg Ala Arg Asp Ser Val Asp Asp Ile Arg Val
 20 25 30

Ala Arg Val Ile Glu Gln Asp Met Ala Val Asp Ser Ala Gly Lys Ile
 35 40 45

Thr Tyr Arg Ile Lys Leu Glu Val Ser Phe Lys Met Arg Pro Ala Gln
 50 55 60

Pro Arg
 65

(2) INFORMATION FOR SEQ ID NO:79:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 69 amino acids
- (B) TYPE: amino acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:79:

```

Val Pro Pro Ala Pro Pro Leu Pro Pro Leu Pro Pro Ser Pro Ile Ser
1           5           10           15

Cys Ala Ser Pro Pro Ser Pro Pro Leu Pro Pro Ala Pro Pro Val Ala
          20           25           30

Pro Gly Pro Pro Met Pro Pro Leu Asp Pro Trp Pro Pro Ala Pro Pro
          35           40           45

Leu Pro Tyr Ser Thr Pro Pro Gly Ala Pro Leu Pro Pro Ser Pro Pro
          50           55           60

Ser Pro Pro Leu Pro
          65

```

(2) INFORMATION FOR SEQ ID NO:80:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 355 amino acids
- (B) TYPE: amino acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:80:

```

Met Ser Asn Ser Arg Arg Arg Ser Leu Arg Trp Ser Trp Leu Leu Ser
1           5           10           15

Val Leu Ala Ala Val Gly Leu Gly Leu Ala Thr Ala Pro Ala Gln Ala
          20           25           30

Ala Pro Pro Ala Leu Ser Gln Asp Arg Phe Ala Asp Phe Pro Ala Leu
          35           40           45

Pro Leu Asp Pro Ser Ala Met Val Ala Gln Val Ala Pro Gln Val Val
          50           55           60

Asn Ile Asn Thr Lys Leu Gly Tyr Asn Asn Ala Val Gly Ala Gly Thr
          65           70           75           80

```

116

Gly Ile Val Ile Asp Pro Asn Gly Val Val Leu Thr Asn Asn His Val
 85 90 95
 Ile Ala Gly Ala Thr Asp Ile Asn Ala Phe Ser Val Gly Ser Gly Gln
 100 105 110
 Thr Tyr Gly Val Asp Val Val Gly Tyr Asp Arg Thr Gln Asp Val Ala
 115 120 125
 Val Leu Gln Leu Arg Gly Ala Gly Gly Leu Pro Ser Ala Ala Ile Gly
 130 135 140
 Gly Gly Val Ala Val Gly Glu Pro Val Val Ala Met Gly Asn Ser Gly
 145 150 155 160
 Gly Gln Gly Gly Thr Pro Arg Ala Val Pro Gly Arg Val Val Ala Leu
 165 170 175
 Gly Gln Thr Val Gln Ala Ser Asp Ser Leu Thr Gly Ala Glu Glu Thr
 180 185 190
 Leu Asn Gly Leu Ile Gln Phe Asp Ala Ala Ile Gln Pro Gly Asp Ser
 195 200 205
 Gly Gly Pro Val Val Asn Gly Leu Gly Gln Val Val Gly Met Asn Thr
 210 215 220
 Ala Ala Ser Asp Asn Phe Gln Leu Ser Gln Gly Gly Gln Gly Phe Ala
 225 230 235 240
 Ile Pro Ile Gly Gln Ala Met Ala Ile Ala Gly Gln Ile Arg Ser Gly
 245 250 255
 Gly Gly Ser Pro Thr Val His Ile Gly Pro Thr Ala Phe Leu Gly Leu
 260 265 270
 Gly Val Val Asp Asn Asn Gly Asn Gly Ala Arg Val Gln Arg Val Val
 275 280 285
 Gly Ser Ala Pro Ala Ala Ser Leu Gly Ile Ser Thr Gly Asp Val Ile
 290 295 300
 Thr Ala Val Asp Gly Ala Pro Ile Asn Ser Ala Thr Ala Met Ala Asp
 305 310 315 320
 Ala Leu Asn Gly His His Pro Gly Asp Val Ile Ser Val Asn Trp Gln
 325 330 335
 Thr Lys Ser Gly Gly Thr Arg Thr Gly Asn Val Thr Leu Ala Glu Gly
 340 345 350
 Pro Pro Ala
 355

(2) INFORMATION FOR SEQ ID NO:81:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 205 amino acids
- (B) TYPE: amino acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:81:

```

Ser Pro Lys Pro Asp Ala Glu Glu Gln Gly Val Pro Val Ser Pro Thr
1           5           10           15
Ala Ser Asp Pro Ala Leu Leu Ala Glu Ile Arg Gln Ser Leu Asp Ala
20           25           30
Thr Lys Gly Leu Thr Ser Val His Val Ala Val Arg Thr Thr Gly Lys
35           40           45
Val Asp Ser Leu Leu Gly Ile Thr Ser Ala Asp Val Asp Val Arg Ala
50           55           60
Asn Pro Leu Ala Ala Lys Gly Val Cys Thr Tyr Asn Asp Glu Gln Gly
65           70           75           80
Val Pro Phe Arg Val Gln Gly Asp Asn Ile Ser Val Lys Leu Phe Asp
85           90           95
Asp Trp Ser Asn Leu Gly Ser Ile Ser Glu Leu Ser Thr Ser Arg Val
100          105          110
Leu Asp Pro Ala Ala Gly Val Thr Gln Leu Leu Ser Gly Val Thr Asn
115          120          125
Leu Gln Ala Gln Gly Thr Glu Val Ile Asp Gly Ile Ser Thr Thr Lys
130          135          140
Ile Thr Gly Thr Ile Pro Ala Ser Ser Val Lys Met Leu Asp Pro Gly
145          150          155          160
Ala Lys Ser Ala Arg Pro Ala Thr Val Trp Ile Ala Gln Asp Gly Ser
165          170          175
His His Leu Val Arg Ala Ser Ile Asp Leu Gly Ser Gly Ser Ile Gln
180          185          190
Leu Thr Gln Ser Lys Trp Asn Glu Pro Val Asn Val Asp
195          200          205

```

(2) INFORMATION FOR SEQ ID NO:82:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 286 amino acids
- (B) TYPE: amino acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:82:

Gly	Asp	Ser	Phe	Trp	Ala	Ala	Ala	Asp	Gln	Met	Ala	Arg	Gly	Phe	Val	1	5	10	15
Leu	Gly	Ala	Thr	Ala	Gly	Arg	Thr	Thr	Leu	Thr	Gly	Glu	Gly	Leu	Gln	20	25	30	
His	Ala	Asp	Gly	His	Ser	Leu	Leu	Leu	Asp	Ala	Thr	Asn	Pro	Ala	Val	35	40	45	
Val	Ala	Tyr	Asp	Pro	Ala	Phe	Ala	Tyr	Glu	Ile	Gly	Tyr	Ile	Xaa	Glu	50	55	60	
Ser	Gly	Leu	Ala	Arg	Met	Cys	Gly	Glu	Asn	Pro	Glu	Asn	Ile	Phe	Phe	65	70	75	80
Tyr	Ile	Thr	Val	Tyr	Asn	Glu	Pro	Tyr	Val	Gln	Pro	Pro	Glu	Pro	Glu	85	90	95	
Asn	Phe	Asp	Pro	Glu	Gly	Val	Leu	Gly	Gly	Ile	Tyr	Arg	Tyr	His	Ala	100	105	110	
Ala	Thr	Glu	Gln	Arg	Thr	Asn	Lys	Xaa	Gln	Ile	Leu	Ala	Ser	Gly	Val	115	120	125	
Ala	Met	Pro	Ala	Ala	Leu	Arg	Ala	Ala	Gln	Met	Leu	Ala	Ala	Glu	Trp	130	135	140	
Asp	Val	Ala	Ala	Asp	Val	Trp	Ser	Val	Thr	Ser	Trp	Gly	Glu	Leu	Asn	145	150	155	160
Arg	Asp	Gly	Val	Val	Ile	Glu	Thr	Glu	Lys	Leu	Arg	His	Pro	Asp	Arg	165	170	175	
Pro	Ala	Gly	Val	Pro	Tyr	Val	Thr	Arg	Ala	Leu	Glu	Asn	Ala	Arg	Gly	180	185	190	
Pro	Val	Ile	Ala	Val	Ser	Asp	Trp	Met	Arg	Ala	Val	Pro	Glu	Gln	Ile	195	200	205	
Arg	Pro	Trp	Val	Pro	Gly	Thr	Tyr	Leu	Thr	Leu	Gly	Thr	Asp	Gly	Phe	210	215	220	
Gly	Phe	Ser	Asp	Thr	Arg	Pro	Ala	Gly	Arg	Arg	Tyr	Phe	Asn	Thr	Asp				

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 173 amino acids
(B) TYPE: amino acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

Thr 1	Lys	Phe	His 5	Ala	Leu	Met	Gln	Glu 10	Ile	His	Asn	Glu 15	Phe	Thr	
Ala	Ala	Gln 20	Gln	Tyr	Val	Ala	Ile 25	Ala	Val	Tyr	Phe	Asp 30	Ser	Glu	Asp
Leu	Pro 35	Gln	Leu	Ala	Lys	His	Phe 40	Tyr	Ser	Gln	Ala	Val 45	Glu	Glu	Arg
Asn 50	His	Ala	Met	Met	Leu 55	Val	Gln	His	Leu	Leu 60	Asp	Arg	Asp	Leu	Arg
Val 65	Glu	Ile	Pro	Gly 70	Val	Asp	Thr	Val	Arg 75	Asn	Gln	Phe	Asp	Arg	Pro 80
Arg	Glu	Ala	Leu 85	Ala	Leu	Ala	Leu	Asp 90	Gln	Glu	Arg	Thr	Val	Thr 95	Asp
Gln	Val	Gly 100	Arg	Leu	Thr	Ala	Val	Ala 105	Arg	Asp	Glu	Gly	Asp 110	Phe	Leu
Gly	Glu 115	Gln	Phe	Met	Gln	Trp	Phe 120	Leu	Gln	Glu	Gln	Ile 125	Glu	Glu	Val
Ala 130	Leu	Met	Ala	Thr	Leu 135	Val	Arg	Val	Ala	Asp 140	Arg	Ala	Gly	Ala	Asn
Leu 145	Phe	Glu	Leu	Glu 150	Asn	Phe	Val	Ala	Arg 155	Glu	Val	Asp	Val	Ala	Pro 160

Ala Ala Ser Gly Ala Pro His Ala Ala Gly Gly Arg Leu
 165 170

(2) INFORMATION FOR SEQ ID NO:84:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 107 amino acids
- (B) TYPE: amino acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:84:

Arg Ala Asp Glu Arg Lys Asn Thr Thr Met Lys Met Val Lys Ser Ile
 1 5 10 15
 Ala Ala Gly Leu Thr Ala Ala Ala Ala Ile Gly Ala Ala Ala Ala Gly
 20 25 30
 Val Thr Ser Ile Met Ala Gly Gly Pro Val Val Tyr Gln Met Gln Pro
 35 40 45
 Val Val Phe Gly Ala Pro Leu Pro Leu Asp Pro Xaa Ser Ala Pro Xaa
 50 55 60
 Val Pro Thr Ala Ala Gln Trp Thr Xaa Leu Leu Asn Xaa Leu Xaa Asp
 65 70 75 80
 Pro Asn Val Ser Phe Xaa Asn Lys Gly Ser Leu Val Glu Gly Gly Ile
 85 90 95
 Gly Gly Xaa Glu Gly Xaa Xaa Arg Arg Xaa Gln
 100 105

(2) INFORMATION FOR SEQ ID NO:85:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 125 amino acids
- (B) TYPE: amino acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:85:

Val Leu Ser Val Pro Val Gly Asp Gly Phe Trp Xaa Arg Val Val Asn
 1 5 10 15
 Pro Leu Gly Gln Pro Ile Asp Gly Arg Gly Asp Val Asp Ser Asp Thr

121

	20		25		30	
Arg	Arg	Ala	Leu	Glu	Leu	Gln
	35					40
Pro	Ser	Val	Val	Xaa	Arg	Gln
				45		
Val	Lys	Glu	Pro	Leu	Xaa	Thr
	50					55
Gly	Ile	Lys	Ala	Ile	Asp	Ala
				60		
Pro	Ile	Gly	Arg	Gly	Gln	Arg
	65				70	
Gln	Leu	Ile	Ile	Gly	Asp	Arg
				75		80
Gly	Lys	Asn	Arg	Arg	Leu	Cys
				85		
Thr	Pro	Ser	Ser	Asn	Gln	Arg
				90		95
Glu	Leu	Gly	Val	Arg	Trp	Ile
				100		
Pro	Arg	Ser	Arg	Cys	Ala	Cys
				105		110
Val	Gly	His	Arg	Ala	Arg	Arg
	115					120
						125

(2) INFORMATION FOR SEQ ID NO:86:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 117 amino acids
- (B) TYPE: amino acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:86:

Cys	Asp	Ala	Val	Met	Gly	Phe	Leu	Gly	Gly	Ala	Gly	Pro	Leu	Ala	Val
1			5					10					15		
Val	Asp	Gln	Gln	Leu	Val	Thr	Arg	Val	Pro	Gln	Gly	Trp	Ser	Phe	Ala
		20						25					30		
Gln	Ala	Ala	Ala	Val	Pro	Val	Val	Phe	Leu	Thr	Ala	Trp	Tyr	Gly	Leu
		35					40					45			
Ala	Asp	Leu	Ala	Glu	Ile	Lys	Ala	Gly	Glu	Ser	Val	Leu	Ile	His	Ala
	50					55					60				
Gly	Thr	Gly	Gly	Val	Gly	Met	Ala	Ala	Val	Gln	Leu	Ala	Arg	Gln	Trp
	65				70					75				80	
Gly	Val	Glu	Val	Phe	Val	Thr	Ala	Ser	Arg	Gly	Lys	Trp	Asp	Thr	Leu
				85					90					95	
Arg	Ala	Xaa	Xaa	Phe	Asp	Asp	Xaa	Pro	Tyr	Arg	Xaa	Phe	Pro	His	Xaa
		100						105					110		

Arg Ser Ser Xaa Gly
115

(2) INFORMATION FOR SEQ ID NO:87:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 103 amino acids
- (B) TYPE: amino acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:87:

Met	Tyr	Arg	Phe	Ala	Cys	Arg	Thr	Leu	Met	Leu	Ala	Ala	Cys	Ile	Leu	1	5	10	15
Ala	Thr	Gly	Val	Ala	Gly	Leu	Gly	Val	Gly	Ala	Gln	Ser	Ala	Ala	Gln	20	25	30	
Thr	Ala	Pro	Val	Pro	Asp	Tyr	Tyr	Trp	Cys	Pro	Gly	Gln	Pro	Phe	Asp	35	40	45	
Pro	Ala	Trp	Gly	Pro	Asn	Trp	Asp	Pro	Tyr	Thr	Cys	His	Asp	Asp	Phe	50	55	60	
His	Arg	Asp	Ser	Asp	Gly	Pro	Asp	His	Ser	Arg	Asp	Tyr	Pro	Gly	Pro	65	70	75	80
Ile	Leu	Glu	Gly	Pro	Val	Leu	Asp	Asp	Pro	Gly	Ala	Ala	Pro	Pro	Pro	85	90	95	
Pro	Ala	Ala	Gly	Gly	Gly	Ala										100			

(2) INFORMATION FOR SEQ ID NO:88:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 88 amino acids
- (B) TYPE: amino acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:88:

Val	Gln	Cys	Arg	Val	Trp	Leu	Glu	Ile	Gln	Trp	Arg	Gly	Met	Leu	Gly	1	5	10	15
-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	---	---	----	----

123

Ala Asp Gln Ala Arg Ala Gly Gly Pro Ala Arg Ile Trp Arg Glu His
 20 25 30

Ser Met Ala Ala Met Lys Pro Arg Thr Gly Asp Gly Pro Leu Glu Ala
 35 40 45

Thr Lys Glu Gly Arg Gly Ile Val Met Arg Val Pro Leu Glu Gly Gly
 50 55 60

Gly Arg Leu Val Val Glu Leu Thr Pro Asp Glu Ala Ala Ala Leu Gly
 65 70 75 80

Asp Glu Leu Lys Gly Val Thr Ser
 85

(2) INFORMATION FOR SEQ ID NO:89:

- (i) SEQUENCE CHARACTERISTICS:
- (A) LENGTH: 95 amino acids
 - (B) TYPE: amino acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:89:

Thr Asp Ala Ala Thr Leu Ala Gln Glu Ala Gly Asn Phe Glu Arg Ile
 1 5 10 15

Ser Gly Asp Leu Lys Thr Gln Ile Asp Gln Val Glu Ser Thr Ala Gly
 20 25 30

Ser Leu Gln Gly Gln Trp Arg Gly Ala Ala Gly Thr Ala Ala Gln Ala
 35 40 45

Ala Val Val Arg Phe Gln Glu Ala Ala Asn Lys Gln Lys Gln Glu Leu
 50 55 60

Asp Glu Ile Ser Thr Asn Ile Arg Gln Ala Gly Val Gln Tyr Ser Arg
 65 70 75 80

Ala Asp Glu Glu Gln Gln Gln Ala Leu Ser Ser Gln Met Gly Phe
 85 90 95

(2) INFORMATION FOR SEQ ID NO:90:

- (i) SEQUENCE CHARACTERISTICS:
- (A) LENGTH: 166 amino acids
 - (B) TYPE: amino acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:90:

```

Met Thr Gln Ser Gln Thr Val Thr Val Asp Gln Gln Glu Ile Leu Asn
1           5           10           15

Arg Ala Asn Glu Val Glu Ala Pro Met Ala Asp Pro Pro Thr Asp Val
          20           25           30

Pro Ile Thr Pro Cys Glu Leu Thr Xaa Xaa Lys Asn Ala Ala Gln Gln
          35           40           45

Xaa Val Leu Ser Ala Asp Asn Met Arg Glu Tyr Leu Ala Ala Gly Ala
          50           55           60

Lys Glu Arg Gln Arg Leu Ala Thr Ser Leu Arg Asn Ala Ala Lys Xaa
65           70           75           80

Tyr Gly Glu Val Asp Glu Glu Ala Ala Thr Ala Leu Asp Asn Asp Gly
          85           90           95

Glu Gly Thr Val Gln Ala Glu Ser Ala Gly Ala Val Gly Gly Asp Ser
          100          105          110

Ser Ala Glu Leu Thr Asp Thr Pro Arg Val Ala Thr Ala Gly Glu Pro
          115          120          125

Asn Phe Met Asp Leu Lys Glu Ala Ala Arg Lys Leu Glu Thr Gly Asp
          130          135          140

Gln Gly Ala Ser Leu Ala His Xaa Gly Asp Gly Trp Asn Thr Xaa Thr
145          150          155          160

Leu Thr Leu Gln Gly Asp
          165

```

(2) INFORMATION FOR SEQ ID NO:91:

- (i) SEQUENCE CHARACTERISTICS:
- (A) LENGTH: 5 amino acids
 - (B) TYPE: amino acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:91:

```

Arg Ala Glu Arg Met
1           5

```

(2) INFORMATION FOR SEQ ID NO:92:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 263 amino acids
- (B) TYPE: amino acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:92:

```

Val Ala Trp Met Ser Val Thr Ala Gly Gln Ala Glu Leu Thr Ala Ala
1           5           10           15/
Gln Val Arg Val Ala Ala Ala Ala Tyr Glu Thr Ala Tyr Gly Leu Thr
          20           25           30
Val Pro Pro Pro Val Ile Ala Glu Asn Arg Ala Glu Leu Met Ile Leu
          35           40           45
Ile Ala Thr Asn Leu Leu Gly Gln Asn Thr Pro Ala Ile Ala Val Asn
          50           55           60
Glu Ala Glu Tyr Gly Glu Met Trp Ala Gln Asp Ala Ala Ala Met Phe
          65           70           75           80
Gly Tyr Ala Ala Ala Thr Ala Thr Ala Thr Ala Thr Leu Leu Pro Phe
          85           90           95
Glu Glu Ala Pro Glu Met Thr Ser Ala Gly Gly Leu Leu Glu Gln Ala
          100          105          110
Ala Ala Val Glu Glu Ala Ser Asp Thr Ala Ala Ala Asn Gln Leu Met
          115          120          125
Asn Asn Val Pro Gln Ala Leu Lys Gln Leu Ala Gln Pro Thr Gln Gly
          130          135          140
Thr Thr Pro Ser Ser Lys Leu Gly Gly Leu Trp Lys Thr Val Ser Pro
          145          150          155          160
His Arg Ser Pro Ile Ser Asn Met Val Ser Met Ala Asn Asn His Met
          165          170          175
Ser Met Thr Asn Ser Gly Val Ser Met Thr Asn Thr Leu Ser Ser Met
          180          185          190
Leu Lys Gly Phe Ala Pro Ala Ala Ala Ala Gln Ala Val Gln Thr Ala
          195          200          205
Ala Gln Asn Gly Val Arg Ala Met Ser Ser Leu Gly Ser Ser Leu Gly

```

210	215	220
Ser Ser Gly Leu Gly Gly Gly Val Ala Ala Asn Leu Gly Arg Ala Ala		
225	230	235 240
Ser Val Arg Tyr Gly His Arg Asp Gly Gly Lys Tyr Ala Xaa Ser Gly		
	245	250 255
Arg Arg Asn Gly Gly Pro Ala		
	260	

(2) INFORMATION FOR SEQ ID NO:93:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 303 amino acids
- (B) TYPE: amino acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:93:

Met Thr Tyr Ser Pro Gly Asn Pro Gly Tyr Pro Gln Ala Gln Pro Ala	
1	5 10 15
Gly Ser Tyr Gly Gly Val Thr Pro Ser Phe Ala His Ala Asp Glu Gly	
	20 25 30
Ala Ser Lys Leu Pro Met Tyr Leu Asn Ile Ala Val Ala Val Leu Gly	
	35 40 45
Leu Ala Ala Tyr Phe Ala Ser Phe Gly Pro Met Phe Thr Leu Ser Thr	
	50 55 60
Glu Leu Gly Gly Gly Asp Gly Ala Val Ser Gly Asp Thr Gly Leu Pro	
65	70 75 80
Val Gly Val Ala Leu Leu Ala Ala Leu Leu Ala Gly Val Val Leu Val	
	85 90 95
Pro Lys Ala Lys Ser His Val Thr Val Val Ala Val Leu Gly Val Leu	
	100 105 110
Gly Val Phe Leu Met Val Ser Ala Thr Phe Asn Lys Pro Ser Ala Tyr	
	115 120 125
Ser Thr Gly Trp Ala Leu Trp Val Val Leu Ala Phe Ile Val Phe Gln	
	130 135 140
Ala Val Ala Ala Val Leu Ala Leu Leu Val Glu Thr Gly Ala Ile Thr	
145	150 155 160

127

Ala Pro Ala Pro Arg Pro Lys Phe Asp Pro Tyr Gly Gln Tyr Gly Arg
165 170 175

Tyr Gly Gln Tyr Gly Gln Tyr Gly Val Gln Pro Gly Gly Tyr Tyr Gly
180 185 190

Gln Gln Gly Ala Gln Gln Ala Ala Gly Leu Gln Ser Pro Gly Pro Gln
195 200 205

Gln Ser Pro Gln Pro Pro Gly Tyr Gly Ser Gln Tyr Gly Gly Tyr Ser
210 215 220

Ser Ser Pro Ser Gln Ser Gly Ser Gly Tyr Thr Ala Gln Pro Pro Ala
225 230 235 240

Gln Pro Pro Ala Gln Ser Gly Ser Gln Gln Ser His Gln Gly Pro Ser
245 250 255

Thr Pro Pro Thr Gly Phe Pro Ser Phe Ser Pro Pro Pro Pro Val Ser
260 265 270

Ala Gly Thr Gly Ser Gln Ala Gly Ser Ala Pro Val Asn Tyr Ser Asn
275 280 285

Pro Ser Gly Gly Glu Gln Ser Ser Ser Pro Gly Gly Ala Pro Val
290 295 300

(2) INFORMATION FOR SEQ ID NO:94:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 507 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:94:

ATGAAGATGG TGAAATCGAT CGCCGCAGGT CTGACCGCCG CGGCTGCAAT CGGCGCCGCT	60
GCGGCCGGTG TGACTTCGAT CATGGCTGGC GGCCCGGTCG TATACCAGAT GCAGCCGGTC	120
GTCTTCGGCG CGCCACTGCC GTTGGACCCG GCATCCGCCC CTGACGTCCC GACCGCCGCC	180
CAGTTGACCA GCCTGCTCAA CAGCCTCGCC GATCCCAACG TGTCGTTTGC GAACAAGGGC	240
AGTCTGGTCG AGGGCGGCAT CGGGGGCACC GAGGCGCGCA TCGCCGACCA CAAGCTGAAG	300
AAGGCCGCCG AGCACGGGGA TCTGCCGCTG TCGTTCAGCG TGACGAACAT CCAGCCGGCG	360
GCCGCCGGTT CGGCCACCGC CGACGTTTCC GTCTCGGGTC CGAAGCTCTC GTCGCCGGTC	420

ACGCAGAACG TCACGTTTCGT GAATCAAGGC GGCTGGATGC TGTCACGCGC ATCGGCGATG 480
 GAGTTGCTGC AGGCCGCGAGG GAACTGA 507

(2) INFORMATION FOR SEQ ID NO:95:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 168 amino acids
- (B) TYPE: amino acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:95:

Met Lys Met Val Lys Ser Ile Ala Ala Gly Leu Thr Ala Ala Ala Ala
 1 5 10 15
 Ile Gly Ala Ala Ala Ala Gly Val Thr Ser Ile Met Ala Gly Gly Pro
 20 25 30
 Val Val Tyr Gln Met Gln Pro Val Val Phe Gly Ala Pro Leu Pro Leu
 35 40 45
 Asp Pro Ala Ser Ala Pro Asp Val Pro Thr Ala Ala Gln Leu Thr Ser
 50 55 60
 Leu Leu Asn Ser Leu Ala Asp Pro Asn Val Ser Phe Ala Asn Lys Gly
 65 70 75 80
 Ser Leu Val Glu Gly Gly Ile Gly Gly Thr Glu Ala Arg Ile Ala Asp
 85 90 95
 His Lys Leu Lys Lys Ala Ala Glu His Gly Asp Leu Pro Leu Ser Phe
 100 105 110
 Ser Val Thr Asn Ile Gln Pro Ala Ala Ala Gly Ser Ala Thr Ala Asp
 115 120 125
 Val Ser Val Ser Gly Pro Lys Leu Ser Ser Pro Val Thr Gln Asn Val
 130 135 140
 Thr Phe Val Asn Gln Gly Gly Trp Met Leu Ser Arg Ala Ser Ala Met
 145 150 155 160
 Glu Leu Leu Gln Ala Ala Gly Asn
 165

(2) INFORMATION FOR SEQ ID NO:96:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 500 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:96:

```

CGTGGCAATG TCGTTGACCG TCGGGGCCGG GGTGCGCTCC GCAGATCCCG TGGACGCGGT      60
CATTAAACACC ACCTGCAATT ACGGGCAGGT AGTAGCTGCG CTCAACGCGA CGGATCCGGG      120
GGCTGCCGCA CAGTTCAACG CCTCACCGGT GGCGCAGTCC TATTTGCGCA ATTTCTCTCGC      180
CGCACCGCCA CCTCAGCGCG CTGCCATGGC CGCGCAATTG CAAGCTGTGC CGGGGGCGGC      240
ACAGTACATC GGCCTTGTCG AGTCGGTTGC CGGCTCCTGC AACAACTATT AAGCCCATGC      300
GGGCCCCATC CCGCGACCCG GCATCGTCGC CGGGGCTAGG CCAGATTGCC CCGCTCCTCA      360
ACGGGGCCGCA TCCGCGGACC CGGCATCGTC GCCGGGGCTA GGCCAGATTG CCCCCTCCT      420
CAACGGGGCCG CATCTCGTGC CGAATTCCTG CAGCCCGGGG GATCCACTAG TTCTAGAGCG      480
GCCGCCACCG CGGTGGAGCT                                         500

```

(2) INFORMATION FOR SEQ ID NO:97:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 96 amino acids
 - (B) TYPE: amino acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:97:

```

Val Ala Met Ser Leu Thr Val Gly Ala Gly Val Ala Ser Ala Asp Pro
1           5           10           15
Val Asp Ala Val Ile Asn Thr Thr Cys Asn Tyr Gly Gln Val Val Ala
20           25           30
Ala Leu Asn Ala Thr Asp Pro Gly Ala Ala Ala Gln Phe Asn Ala Ser
35           40           45
Pro Val Ala Gln Ser Tyr Leu Arg Asn Phe Leu Ala Ala Pro Pro Pro
50           55           60

```

Gln Arg Ala Ala Met Ala Ala Gln Leu Gln Ala Val Pro Gly Ala Ala
 65 70 75 80
 Gln Tyr Ile Gly Leu Val Glu Ser Val Ala Gly Ser Cys Asn Asn Tyr
 85 90 95

(2) INFORMATION FOR SEQ ID NO:98:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 154 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:98:

ATGACAGAGC AGCAGTGGAA TTTCGCGGGT ATCGAGGCCG CGGCAAGCGC AATCCAGGGA 60
 AATGTCACGT CCATTCATTC CCTCCTTGAC GAGGGGAAGC AGTCCCTGAC CAAGCTCGCA 120
 GCGGCCTGGG GCGGTAGCGG TTCGGAAGCG TACC 154

(2) INFORMATION FOR SEQ ID NO:99:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 51 amino acids
 (B) TYPE: amino acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:99:

Met Thr Glu Gln Gln Trp Asn Phe Ala Gly Ile Glu Ala Ala Ala Ser
 1 5 10 15
 Ala Ile Gln Gly Asn Val Thr Ser Ile His Ser Leu Leu Asp Glu Gly
 20 25 30
 Lys Gln Ser Leu Thr Lys Leu Ala Ala Ala Trp Gly Gly Ser Gly Ser
 35 40 45
 Glu Ala Tyr
 50

(2) INFORMATION FOR SEQ ID NO:100:

- (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 282 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:100:

```

CGGTCGCGCA CTTCCAGGTG ACTATGAAAG TCGGCTTCCG NCTGGAGGAT TCCTGAACCT      60
TCAAGCGCGG CCGATAACTG AGGTGCATCA TTAAGCGACT TTTCCAGAAC ATCCTGACGC      120
GCTCGAAACG CGGCACAGCC GACGGTGGCT CCGNCGAGGC GCTGNCTCCA AAATCCCTGA      180
GACAATTTCGN CGGGGGCGCC TACAAGGAAG TCGGTGCTGA ATTCGNCNG TATCTGGTTCG      240
ACCTGTGTGG TCTGNAGCCG GACGAAGCGG TGCTCGACGT CG                          282

```

(2) INFORMATION FOR SEQ ID NO:101:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 3058 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:101:

```

GATCGTACCC GTGCGAGTGC TCGGGCCGTT TGAGGATGGA GTGCACGTGT CTTTCGTGAT      60
GGCATACCCA GAGATGTTGG CGGCGGCGGC TGACACCCTG CAGAGCATCG GTGCTACCAC      120
TGTGGCTAGC AATGCCGCTG CGGCGGCCCC GACGACTGGG GTGGTGCCCC CCGCTGCCGA      180
TGAGGTGTCG GCGCTGACTG CGGCGCACTT CGCCGCACAT GCGGCGATGT ATCAGTCCGT      240
GAGCGCTCGG GCTGCTGCGA TTCATGACCA GTTCGTGGCC ACCCTTGCCA GCAGCGCCAG      300
CTCGTATGCG GCCACTGAAG TCGCCAATGC GGCGGCGGCC AGCTAAGCCA GGAACAGTCG      360
GCACGAGAAA CCACGAGAAA TAGGGACACG TAATGGTGGA TTTCGGGGCG TTACCACCGG      420
AGATCAACTC CGCGAGGATG TACGCCGGCC CGGGTTCGGC CTCGCTGGTG GCCGCGGCTC      480
AGATGTGGGA CAGCGTGGCG AGTGACCTGT TTTCGGCCGC GTCGGCGTTT CAGTCGGTGG      540
TCTGGGGTCT GACGGTGGGG TCGTGGATAG GTTCGTGCGC GGGTCTGATG GTGGCGGCGG      600

```

CCTCGCCGTA	TGTGGCGTGG	ATGAGCGTCA	CCGCGGGGCA	GGCCGAGCTG	ACCGCCGCCC	660
AGGTCCGGGT	TGCTGCGGCG	GCCTACGAGA	CGGCGTATGG	GCTGACGGTG	CCCCCGCCGG	720
TGATCGCCGA	GAACCGTGCT	GAAGTATGA	TTCTGATAGC	GACCAACCTC	TTGGGGCAAA	780
ACACCCCGGC	GATCGCGGTC	AACGAGGCCG	AATACGGCGA	GATGTGGGCC	CAAGACGCCG	840
CCGCGATGTT	TGGCTACGCC	GCGGCGACGG	CGACGGCGAC	GGCGACGTTG	CTGCCGTTTCG	900
AGGAGGCGCC	GGAGATGACC	AGCGCGGGTG	GGCTCCTCGA	GCAGGCCGCC	GCGGTTCGAGG	960
AGGCCTCCGA	CACCGCCGCG	GCGAACCAGT	TGATGAACAA	TGTGCCCCAG	GCGCTGCAAC	1020
AGCTGGCCCA	GCCCACGCAG	GGCACCACGC	CTTCTTCCAA	GCTGGGTGGC	CTGTGGAAGA	1080
CGGTCTCGCC	GCATCGGTCTG	CCGATCAGCA	ACATGGTGTC	GATGGCCAAC	AACCACATGT	1140
CGATGACCAA	CTCGGGTGTG	TCGATGACCA	ACACCTTGAG	CTCGATGTTG	AAGGGCTTTG	1200
CTCCGGCGGC	GGCCGCCCAG	GCCGTGCAAA	CCGCGGCGCA	AAACGGGGTC	CGGGCGATGA	1260
GCTCGCTGGG	CAGCTCGCTG	GGTTCTTCGG	GTCTGGGCGG	TGGGGTGGCC	GCCAACTTGG	1320
GTCGGGCGGC	CTCGGTCTGGT	TCGTTGTCTG	TGCCGCAGGC	CTGGGCCGCG	GCCAACCAGG	1380
CAGTCACCCC	GGCGGCGCGG	GCGCTGCCGC	TGACCAGCCT	GACCAGCGCC	GCGGAAAGAG	1440
GGCCCGGGCA	GATGCTGGGC	GGGCTGCCGG	TGGGGCAGAT	GGGCGCCAGG	GCCGGTGGTG	1500
GGCTCAGTGG	TGTGCTGCGT	GTTCCGCCGC	GACCCATATG	GATGCCGCAT	TCTCCGGCGG	1560
CCGGCTAGGA	GAGGGGGCGC	AGACTGTCGT	TATTTGACCA	GTGATCGGCG	GTCTCGGTGT	1620
TTCCGCGGCC	GGCTATGACA	ACAGTCAATG	TGCATGACAA	GTTACAGGTA	TTAGGTCCAG	1680
GTTCAACAAG	GAGACAGGCA	ACATGGCCTC	ACGTTTTATG	ACGGATCCGC	ACGCGATGCG	1740
GGACATGGCG	GGCCGTTTTG	AGGTGCACGC	CCAGACGGTG	GAGGACGAGG	CTCGCCGGAT	1800
GTGGGCGTCC	GCGCAAAACA	TTTCCGGTGC	GGGCTGGAGT	GGCATGGCCG	AGGCGACCTC	1860
GCTAGACACC	ATGGCCCAGA	TGAATCAGGC	GTTTCGCAAC	ATCGTGAACA	TGCTGCACGG	1920
GGTGCGTGAC	GGGCTGGTTC	GCGACGCCAA	CAACTACGAG	CAGCAAGAGC	AGGCCTCCCA	1980
GCAGATCCTC	AGCAGCTAAC	GTCAGCCGCT	GCAGCACAAT	ACTTTTACAA	GCGAAGGAGA	2040
ACAGGTTCTGA	TGACCATCAA	CTATCAATTC	GGGGATGTCTG	ACGCTCACGG	CGCCATGATC	2100
CGCGCTCAGG	CCGGGTTGCT	GGAGGCCGAG	CATCAGGCCA	TCATTCGTGA	TGTGTTGACC	2160
GCGAGTGAAT	TTTGGGGCGG	CGCCGGTTCG	GCGGCCTGCC	AGGGGTTCAT	TACCCAGTTG	2220
GGCCGTAACT	TCCAGGTGAT	CTACGAGCAG	GCCAACGCCC	ACGGGCAGAA	GGTGCAGGCT	2280

```

GCCGGCAACA ACATGGCGCA AACCGACAGC GCCGTCGGCT CCAGCTGGGC CTGACACCAG      2340
GCCAAGGCCA GGGACGTGGT GTACGAGTGA AGTTCCTCGC GTGATCCTTC GGGTGGCAGT      2400
CTAAGTGGTC AGTGCTGGGG TGTGGTGGT TTGCTGCTTG GCGGGTTCTT CGGTGCTGGT      2460
CAGTGCTGCT CGGGCTCGGG TGAGGACCTC GAGGCCCAGG TAGCGCCGTC CTCGATCCA      2520
TTCGTCGTGT TGTTCCGGCA GGACGGCTCC GACGAGGCGG ATGATCGAGG CGCGGTCGGG      2580
GAAGATGCCC ACGACGTCGG TTCGGCGTCG TACCTCTCGG TTGAGGCGTT CCTGGGGGTT      2640
GTTGGACCAG ATTTGGCGCC AGATCTGCTT GGGGAAGGCG GTGAACGCCA GCAGGTCGGT      2700
GCGGGCGGTG TCGAGGTGCT CGGCCACCGC GGGGAGTTTG TCGGTCAGAG CGTCGAGTAC      2760
CCGATCATAT TGGGCAACAA CTGATTCGGC GTCGGGCTGG TCGTAGATGG AGTGCAGCAG      2820
GGTGCGCACC CACGGCCAGG AGGGCTTCGG GGTGGCTGCC ATCAGATTGG CTGCGTAGTG      2880
GGTTCTGCAG CGCTGCCAGG CCGCTGCGGG CAGGGTGGCG CCGATCGCGG CCACCAGGCC      2940
GGCGTGGGCG TCGCTGGTGA CCAGCGCGAC CCCGGACAGG CCGCGGGCGA CCAGGTCGCG      3000
GAAGAACGCC AGCCAGCCGG CCCCGTCCTC GGCGGAGGTG ACCTGGATGC CCAGGATC      3058

```

(2) INFORMATION FOR SEQ ID NO:102:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 391 amino acids
- (B) TYPE: amino acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:102:

```

Met Val Asp Phe Gly Ala Leu Pro Pro Glu Ile Asn Ser Ala Arg Met
1           5           10           15
Tyr Ala Gly Pro Gly Ser Ala Ser Leu Val Ala Ala Ala Gln Met Trp
20          25          30
Asp Ser Val Ala Ser Asp Leu Phe Ser Ala Ala Ser Ala Phe Gln Ser
35          40          45
Val Val Trp Gly Leu Thr Val Gly Ser Trp Ile Gly Ser Ser Ala Gly
50          55          60
Leu Met Val Ala Ala Ala Ser Pro Tyr Val Ala Trp Met Ser Val Thr
65          70          75          80

```

Ala Gly Gln Ala Glu Leu Thr Ala Ala Gln Val Arg Val Ala Ala Ala
 85 90 95
 Ala Tyr Glu Thr Ala Tyr Gly Leu Thr Val Pro Pro Pro Val Ile Ala
 100 105 110
 Glu Asn Arg Ala Glu Leu Met Ile Leu Ile Ala Thr Asn Leu Leu Gly
 115 120 125
 Gln Asn Thr Pro Ala Ile Ala Val Asn Glu Ala Glu Tyr Gly Glu Met
 130 135 140
 Trp Ala Gln Asp Ala Ala Ala Met Phe Gly Tyr Ala Ala Ala Thr Ala
 145 150 155 160
 Thr Ala Thr Ala Thr Leu Leu Pro Phe Glu Glu Ala Pro Glu Met Thr
 165 170 175
 Ser Ala Gly Gly Leu Leu Glu Gln Ala Ala Ala Val Glu Glu Ala Ser
 180 185 190
 Asp Thr Ala Ala Ala Asn Gln Leu Met Asn Asn Val Pro Gln Ala Leu
 195 200 205
 Gln Gln Leu Ala Gln Pro Thr Gln Gly Thr Thr Pro Ser Ser Lys Leu
 210 215 220
 Gly Gly Leu Trp Lys Thr Val Ser Pro His Arg Ser Pro Ile Ser Asn
 225 230 235 240
 Met Val Ser Met Ala Asn Asn His Met Ser Met Thr Asn Ser Gly Val
 245 250 255
 Ser Met Thr Asn Thr Leu Ser Ser Met Leu Lys Gly Phe Ala Pro Ala
 260 265 270
 Ala Ala Ala Gln Ala Val Gln Thr Ala Ala Gln Asn Gly Val Arg Ala
 275 280 285
 Met Ser Ser Leu Gly Ser Ser Leu Gly Ser Ser Gly Leu Gly Gly Gly
 290 295 300
 Val Ala Ala Asn Leu Gly Arg Ala Ala Ser Val Gly Ser Leu Ser Val
 305 310 315 320
 Pro Gln Ala Trp Ala Ala Ala Asn Gln Ala Val Thr Pro Ala Ala Arg
 325 330 335
 Ala Leu Pro Leu Thr Ser Leu Thr Ser Ala Ala Glu Arg Gly Pro Gly
 340 345 350
 Gln Met Leu Gly Gly Leu Pro Val Gly Gln Met Gly Ala Arg Ala Gly
 355 360 365

Gly Gly Leu Ser Gly Val Leu Arg Val Pro Pro Arg Pro Tyr Val Met
 370 375 380

Pro His Ser Pro Ala Ala Gly
 385 390

(2) INFORMATION FOR SEQ ID NO:103:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 1725 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:103:

GACGTCAGCA CCCGCCGTGC AGGGCTGGAG CGTGGTCGGT TTTGATCTGC GGTCAAGGTG	60
ACGTCCCTCG GCGTGTGCGC GCGTGGATG CAGACTCGAT GCCGCTCTTT AGTGCAACTA	120
ATTTCTGTGA AGTGCCTGCG AGGTATAGGA CTTACAGATT GGTTAATGTA GCGTTCACCC	180
CGTGTTGGGG TCGATTGGC CGGACCAGTC GTCACCAACG CTTGGCGTGC GCGCCAGGCG	240
GGCGATCAGA TCGCTTGACT ACCAATCAAT CTTGAGCTCC CGGGCCGATG CTCGGGCTAA	300
ATGAGGAGGA GCACGCGTGT CTTTCACTGC GCAACCGGAG ATGTTGGCGG CCGCGGCTGG	360
CGAACTTCGT TCCCTGGGGG CAACGCTGAA GGCTAGCAAT GCCGCCGCAG CCGTGCCGAC	420
GA CTGGGGTG GTGCCCCCGG CTGCCGACGA GGTGTGCTG CTGCTTGCCA CACAATTCCG	480
TACGCATGCG GCGACGTATC AGACGGCCAG CGCCAAGGCC GCGGTGATCC ATGAGCAGTT	540
TGTGACCACG CTGGCCACCA GCGCTAGTTC ATATGCGGAC ACCGAGGCCG CCAACGCTGT	600
GGTCACCGGC TAGCTGACCT GACGGTATTC GAGCGGAAGG ATTATCGAAG TGGTGGATTT	660
CGGGGCGTTA CCACCGGAGA TCAACTCCGC GAGGATGTAC GCCGGCCCGG GTTCGGCCTC	720
GCTGGTGGCC GCCGCGAAGA TGTGGGACAG CGTGGCGAGT GACCTGTTTT CGGCCGCGTC	780
GGCGTTTCAG TCGGTGGTCT GGGGTCTGAC GGTGGGGTCG TGGATAGGTT CGTCGGCGGG	840
TCTGATGGCG GCGGCGGCCT CGCCGTATGT GGCGTGGATG AGCGTCACCG CGGGGCAGGC	900
CCAGCTGACC GCCGCCCAGG TCCGGGTTCG TGCGGCGGCC TACGAGACAG CGTATAGGCT	960
GACGGTGCCC CCGCCGGTGA TCGCCGAGAA CCGTACCGAA CTGATGACGC TGACCGCGAC	1020
CAACCTCTTG GGGCAAAACA CGCCGGCGAT CGAGGCCAAT CAGGCCGCAT ACAGCCAGAT	1080

GTGGGGCCAA GACGCGGAGG CGATGTATGG CTACGCCGCC ACGGCGGCGA CGGCGACCGA 1140
 GGC GTTGCTG CCGTTCGAGG ACGCCCCACT GATCACCAAC CCCGGCGGGC TCCTTGAGCA 1200
 GGCCGTCGCG GTCGAGGAGG CCATCGACAC CGCCGCGGCG AACCAGTTGA TGAACAATGT 1260
 GCCCCAAGCG CTGCAACAGC TGGCCCAGCC AGCGCAGGGC GTCGTACCTT CTTCCAAGCT 1320
 GGGTGGGCTG TGGACGGCGG TCTCGCCGCA TCTGTGCGCCG CTCAGCAACG TCAGTTCGAT 1380
 AGCCAACAAC CACATGTCTG TGATGGGCAC GGGTGTGTCTG ATGACCAACA CCTTGCACTC 1440
 GATGTTGAAG GGCTTAGCTC CGGCGGCGGC TCAGGCCGTG GAAACCGCGG CGGAAAACGG 1500
 GGTCTGGGCG ATGAGCTCGC TGGGCAGCCA GCTGGGTTCG TCGCTGGGTT CTTCGGGTCT 1560
 GGGCGCTGGG GTGGCCGCCA ACTTGGGTCTG GGCGGCCTCG GTCGGTTCGT TGTCGGGTGCC 1620
 GCCAGCATGG GCCGCGGCCA ACCAGGCGGT CACCCCGGCG GCGCGGGCGC TGCCGCTGAC 1680
 CAGCCTGACC AGCGCCGCC AAACCGCCCC CGGACACATG CTGGG 1725

(2) INFORMATION FOR SEQ ID NO:104:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 359 amino acids
- (B) TYPE: amino acid
- (C) STRANDEDNESS:
- (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:104:

Val Val Asp Phe Gly Ala Leu Pro Pro Glu Ile Asn Ser Ala Arg Met
 1 5 10 15
 Tyr Ala Gly Pro Gly Ser Ala Ser Leu Val Ala Ala Ala Lys Met Trp
 20 25 30
 Asp Ser Val Ala Ser Asp Leu Phe Ser Ala Ala Ser Ala Phe Gln Ser
 35 40 45
 Val Val Trp Gly Leu Thr Val Gly Ser Trp Ile Gly Ser Ser Ala Gly
 50 55 60
 Leu Met Ala Ala Ala Ala Ser Pro Tyr Val Ala Trp Met Ser Val Thr
 65 70 75 80
 Ala Gly Gln Ala Gln Leu Thr Ala Ala Gln Val Arg Val Ala Ala Ala
 85 90 95

137

Ala Tyr Glu Thr Ala Tyr Arg Leu Thr Val Pro Pro Pro Val Ile Ala
100 105 110

Glu Asn Arg Thr Glu Leu Met Thr Leu Thr Ala Thr Asn Leu Leu Gly
115 120 125

Gln Asn Thr Pro Ala Ile Glu Ala Asn Gln Ala Ala Tyr Ser Gln Met
130 135 140

Trp Gly Gln Asp Ala Glu Ala Met Tyr Gly Tyr Ala Ala Thr Ala Ala
145 150 155 160

Thr Ala Thr Glu Ala Leu Leu Pro Phe Glu Asp Ala Pro Leu Ile Thr
165 170 175

Asn Pro Gly Gly Leu Leu Glu Gln Ala Val Ala Val Glu Glu Ala Ile
180 185 190

Asp Thr Ala Ala Ala Asn Gln Leu Met Asn Asn Val Pro Gln Ala Leu
195 200 205

Gln Gln Leu Ala Gln Pro Ala Gln Gly Val Val Pro Ser Ser Lys Leu
210 215 220

Gly Gly Leu Trp Thr Ala Val Ser Pro His Leu Ser Pro Leu Ser Asn
225 230 235 240

Val Ser Ser Ile Ala Asn Asn His Met Ser Met Met Gly Thr Gly Val
245 250 255

Ser Met Thr Asn Thr Leu His Ser Met Leu Lys Gly Leu Ala Pro Ala
260 265 270

Ala Ala Gln Ala Val Glu Thr Ala Ala Glu Asn Gly Val Trp Ala Met
275 280 285

Ser Ser Leu Gly Ser Gln Leu Gly Ser Ser Leu Gly Ser Ser Gly Leu
290 295 300

Gly Ala Gly Val Ala Ala Asn Leu Gly Arg Ala Ala Ser Val Gly Ser
305 310 315 320

Leu Ser Val Pro Pro Ala Trp Ala Ala Ala Asn Gln Ala Val Thr Pro
325 330 335

Ala Ala Arg Ala Leu Pro Leu Thr Ser Leu Thr Ser Ala Ala Gln Thr
340 345 350

Ala Pro Gly His Met Leu Gly
355

(2) INFORMATION FOR SEQ ID NO:105:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 3027 base pairs

(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:105:

AGTTCAGTCG AGAATGATAC TGACGGGCTG TATCCACGAT GGCTGAGACA ACCGAACCAC	60
CGTCGGACGC GGGGACATCG CAAGCCGACG CGATGGCGTT GGCCGCCGAA GCCGAAGCCG	120
CCGAAGCCGA AGCGCTGGCC GCCGCGGCGC GGGCCCCGTGC CCGTGCCGCC CGGTTGAAGC	180
GTGAGGCGCT GGCGATGGCC CCAGCCGAGG ACGAGAACGT CCCCAGGAT ATGCAGACTG	240
GGAAGACGCC GAAGACTATG ACGACTATGA CGACTATGAG GCCGCAGACC AGGAGGCCGC	300
ACGGTCGGCA TCCTGGCGAC GGCGGTTGCG GGTGCGGTTA CCAAGACTGT CCACGATTGC	360
CATGGCGGCC GCAGTCGTCA TCATCTGCGG CTTACCGGG CTCAGCGGAT ACATTGTGTG	420
GCAACACCAT GAGGCCACCG AACGCCAGCA GCGCGCCGCG GCGTTCGCCG CCGGAGCCAA	480
GCAAGGTGTC ATCAACATGA CCTCGCTGGA CTTCAACAAG GCCAAAGAAG ACGTCGCGCG	540
TGTGATCGAC AGCTCCACCG GCGAATTCAG GGATGACTTC CAGCAGCGGG CAGCCGATTT	600
CACCAAGGTT GTCGAACAGT CCAAAGTGGT CACCGAAGGC ACGGTGAACG CGACAGCCGT	660
CGAATCCATG AACGAGCATT CCGCCGTGGT GCTCGTCGCG GCGACTTCAC GGGTCACCAA	720
TTCCGCTGGG GCGAAAGACG AACCACGTGC GTGGCGGCTC AAAGTGACCG TGACCGAAGA	780
GGGGGGACAG TACAAGATGT CGAAAGTTGA GTTCGTACCG TGACCGATGA CGTACGCGAC	840
GTCAACACCG AAACCACTGA CGCCACCGAA GTCGCTGAGA TCGACTCAGC CGCAGGCGAA	900
GCCGGTGATT CGGCGACCGA GGCATTTGAC ACCGACTCTG CAACGGAATC TACCGCGCAG	960
AAGGGTCAGC GGCACCGTGA CCTGTGGCGA ATGCAGGTTA CCTTGAAACC CGTTCCGGTG	1020
ATTCTCATCC TGCTCATGTT GATCTCTGGG GGCGCGACGG GATGGCTATA CCTTGAGCAA	1080
TACGACCCGA TCAGCAGACG GACTCCGGCG CCGCCCGTGC TGCCGTCGCC GCGGCGTCTG	1140
ACGGGACAAT CGCGCTGTTG TGTATTACCG CGACACGTCG ACCAAGACTT CGCTACCGCC	1200
AGGTGCGACC TCGCCGGCGA TTTCTGTCC TATACGACCA GTTCACGCAG CAGATCGTGG	1260
CTCCGGCGGC CAAACAGAAG TCACTGAAAA CCACCGCCAA GGTGGTGCGC GCGGCCGTGT	1320
CGGAGCTACA TCCGATTCTG GCCGTCGTTT TGGTTTTTGT CGACCAGAGC ACTACCACTA	1380

AGGACAGCCC	CAATCCGTCG	ATGGCGGCCA	GCAGCGTGAT	GGTGACCCTA	GCCAAGGTCG	1440
ACGGCAATTG	GCTGATCACC	AAGTTCACCC	CGGTTTAGGT	TGCCGTAGGC	GGTCGCCAAG	1500
TCTGACGGGG	GCGCGGGTGG	CTGCTCGTGC	GAGATAACCGG	CCGTTCTCCG	GACAATCACG	1560
GCCCGACCTC	AAACAGATCT	CGGCCGCTGT	CTAATCGGCC	GGGTTATTTA	AGATTAGTTG	1620
CCACTGTATT	TACCTGATGT	TCAGATTGTT	CAGCTGGATT	TAGCTTCGCG	GCAGGGCGGC	1680
TGGTGCACTT	TGCATCTGGG	GTTGTGACTA	CTTGAGAGAA	TTTGACCTGT	TGCCGACGTT	1740
GTTTGCTGTC	CATCATTGGT	GCTAGTTATG	GCCGAGCGGA	AGGATTATCG	AAGTGGTGGA	1800
CTTCGGGGCG	TTACCACCGG	AGATCAACTC	CGCGAGGATG	TACGCCGGCC	CGGGTTCGGC	1860
CTCGCTGGTG	GCCGCCGCGA	AGATGTGGGA	CAGCGTGGCG	AGTGACCTGT	TTTCGGCCGC	1920
GTCGGCGTTT	CAGTCGGTGG	TCTGGGGTCT	GACGACGGGA	TCGTGGATAG	GTTCTGTCGGC	1980
GGGTCTGATG	GTGGCGGCGG	CCTCGCCGTA	TGTGGCGTGG	ATGAGCGTCA	CCGCGGGGCA	2040
GGCCGAGCTG	ACCGCCGCCC	AGGTCCGGGT	TGCTGCGGCG	GCCTACGAGA	CGGCGTATGG	2100
GCTGACGGTG	CCCCGCGCGG	TGATCGCCGA	GAACCGTGCT	GAACCTGATGA	TTCTGATAGC	2160
GACCAACCTC	TTGGGGCAAA	ACACCCCGGC	GATCGCGGTC	AACGAGGCCG	AATACGGGGA	2220
GATGTGGGCC	CAAGACGCCG	CCGCGATGTT	TGGCTACGCC	GCCACGGCGG	CGACGGCGAC	2280
CGAGGCGTTG	CTGCCGTTTCG	AGGACGCCCC	ACTGATCACC	AACCCGCGCG	GGCTCCTTGA	2340
GCAGGCCGTC	GCGGTCGAGG	AGGCCATCGA	CACCGCCGCG	GCGAACCAGT	TGATGAACAA	2400
TGTGCCCCAA	GCGCTGCAAC	AACTGGCCCA	GCCCACGAAA	AGCATCTGGC	CGTTCGACCA	2460
ACTGAGTGAA	CTCTGGAAAG	CCATCTCGCC	GCATCTGTCG	CCGCTCAGCA	ACATCGTGTC	2520
GATGCTCAAC	AACCACGTGT	CGATGACCAA	CTCGGGTGTG	TCGATGGCCA	GCACCTTGCA	2580
CTCAATGTTG	AAGGGCTTTG	CTCCGGCGGC	GGCTCAGGCC	GTGGAAACCG	CGGCGCAAAA	2640
CGGGGTCCAG	GCGATGAGCT	CGCTGGGCAG	CCAGCTGGGT	TCGTCGCTGG	GTTCTTCGGG	2700
TCTGGGCGCT	GGGGTGGCCG	CCAACCTGGG	TCGGGCGGCC	TCGGTCGGTT	CGTTGTCCGT	2760
GCCGCAGGCC	TGGGCCGCGG	CCAACCAGGC	GGTCACCCCG	GCGGCGCGGG	CGCTGCCGCT	2820
GACCAGCCTG	ACCAGCGCCG	CCCAAACCGC	CCCCGGACAC	ATGCTGGGCG	GGCTACCGCT	2880
GGGGCAACTG	ACCAATAGCG	GCGGCGGGTT	CGGCGGGGTT	AGCAATGCGT	TGCGGATGCC	2940
GCCGCGGGCG	TACGTAATGC	CCCGTGTGCC	CGCCGCCGGG	TAACGCCGAT	CCGCACGCAA	3000

TGCGGGCCCT CTATGCGGGC AGCGATC

3027

(2) INFORMATION FOR SEQ ID NO:106:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 396 amino acids
- (B) TYPE: amino acid
- (C) STRANDEDNESS:
- (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:106:

```

Val Val Asp Phe Gly Ala Leu Pro Pro Glu Ile Asn Ser Ala Arg Met
1           5           10           15
Tyr Ala Gly Pro Gly Ser Ala Ser Leu Val Ala Ala Ala Lys Met Trp
          20           25           30
Asp Ser Val Ala Ser Asp Leu Phe Ser Ala Ala Ser Ala Phe Gln Ser
          35           40           45
Val Val Trp Gly Leu Thr Thr Gly Ser Trp Ile Gly Ser Ser Ala Gly
          50           55           60
Leu Met Val Ala Ala Ala Ser Pro Tyr Val Ala Trp Met Ser Val Thr
          65           70           75           80
Ala Gly Gln Ala Glu Leu Thr Ala Ala Gln Val Arg Val Ala Ala Ala
          85           90           95
Ala Tyr Glu Thr Ala Tyr Gly Leu Thr Val Pro Pro Pro Val Ile Ala
          100          105          110
Glu Asn Arg Ala Glu Leu Met Ile Leu Ile Ala Thr Asn Leu Leu Gly
          115          120          125
Gln Asn Thr Pro Ala Ile Ala Val Asn Glu Ala Glu Tyr Gly Glu Met
          130          135          140
Trp Ala Gln Asp Ala Ala Ala Met Phe Gly Tyr Ala Ala Thr Ala Ala
          145          150          155          160
Thr Ala Thr Glu Ala Leu Leu Pro Phe Glu Asp Ala Pro Leu Ile Thr
          165          170          175
Asn Pro Gly Gly Leu Leu Glu Gln Ala Val Ala Val Glu Glu Ala Ile
          180          185          190
Asp Thr Ala Ala Ala Asn Gln Leu Met Asn Asn Val Pro Gln Ala Leu
          195          200          205

```

141

Gln Gln Leu Ala Gln Pro Thr Lys Ser Ile Trp Pro Phe Asp Gln Leu
 210 215 220
 Ser Glu Leu Trp Lys Ala Ile Ser Pro His Leu Ser Pro Leu Ser Asn
 225 230 235 240
 Ile Val Ser Met Leu Asn Asn His Val Ser Met Thr Asn Ser Gly Val
 245 250 255
 Ser Met Ala Ser Thr Leu His Ser Met Leu Lys Gly Phe Ala Pro Ala
 260 265 270
 Ala Ala Gln Ala Val Glu Thr Ala Ala Gln Asn Gly Val Gln Ala Met
 275 280 285
 Ser Ser Leu Gly Ser Gln Leu Gly Ser Ser Leu Gly Ser Ser Gly Leu
 290 295 300
 Gly Ala Gly Val Ala Ala Asn Leu Gly Arg Ala Ala Ser Val Gly Ser
 305 310 315 320
 Leu Ser Val Pro Gln Ala Trp Ala Ala Ala Asn Gln Ala Val Thr Pro
 325 330 335
 Ala Ala Arg Ala Leu Pro Leu Thr Ser Leu Thr Ser Ala Ala Gln Thr
 340 345 350
 Ala Pro Gly His Met Leu Gly Gly Leu Pro Leu Gly Gln Leu Thr Asn
 355 360 365
 Ser Gly Gly Gly Phe Gly Gly Val Ser Asn Ala Leu Arg Met Pro Pro
 370 375 380
 Arg Ala Tyr Val Met Pro Arg Val Pro Ala Ala Gly
 385 390 395

(2) INFORMATION FOR SEQ ID NO:107:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 1616 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:107:

CATCGGAGGG AGTGATCACC ATGCTGTGGC ACGCAATGCC ACCGGAGTAA ATACCGCACG	60
GCTGATGGCC GGCGCGGGTC CGGCTCCAAT GCTTGCGGCG GCCGCGGGAT GGCAGACGCT	120
TTCGGCGGCT CTGGACGCTC AGGCCGTCGA GTTGACCGCG CGCCTGAACT CTCTGGGAGA	180

AGCCTGGACT GGAGGTGGCA GCGACAAGGC GCTTGCGGCT GCAACGCCGA TGGTGGTCTG	240
GCTACAAACC GCGTCAACAC AGGCCAAGAC CCGTGCGATG CAGGCGACGG CGCAAGCCGC	300
GGCATAACACC CAGGCCATGG CCACGACGCC GTCGCTGCCG GAGATCGCCG CCAACCACAT	360
CACCCAGGCC GTCCTTACGG CCACCAACTT CTTCGGTATC AACACGATCC CGATCGCGTT	420
GACCGAGATG GATTATTTCA TCCGTATGTG GAACCAGGCA GCCCTGGCAA TGGAGGTCTA	480
CCAGGCCGAG ACCGCGGTTA ACACGCTTTT CGAGAAGCTC GAGCCGATGG CGTCGATCCT	540
TGATCCCGGC GCGAGCCAGA GCACGACGAA CCCGATCTTC GGAATGCCCT CCCCTGGCAG	600
CTCAACACCG GTTGCCAGT TGCCGCCGGC GGCTACCCAG ACCCTCGGCC AACTGGGTGA	660
GATGAGCGGC CCGATGCAGC AGCTGACCCA GCCGCTGCAG CAGGTGACGT CGTTGTTTACG	720
CCAGGTGGGC GGCACCGGCG GCGGCAACCC AGCCGACGAG GAAGCCGCGC AGATGGGCCT	780
GCTCGGCACC AGTCCGCTGT CGAACCATCC GCTGGCTGGT GGATCAGGCC CCAGCGCGGG	840
CGCGGGCCTG CTGCGCGCGG AGTCGCTACC TGGCGCAGGT GGGTCGTTGA CCCGCACGCC	900
GCTGATGTCT CAGCTGATCG AAAAGCCGGT TGCCCCCTCG GTGATGCCGG CGGCTGCTGC	960
CGGATCGTCG GCGACGGGTG GCGCCGCTCC GGTGGGTGCG GGAGCGATGG GCCAGGGTGC	1020
GCAATCCGGC GGCTCCACCA GGCCGGGTCT GGTGCGGCCG GCACCGCTCG CGCAGGAGCG	1080
TGAAGAAGAC GACGAGGACG ACTGGGACGA AGAGGACGAC TGGTGAGCTC CCGTAATGAC	1140
AACAGACTTC CCGGCCACCC GGGCCGGAAG ACTTGCCAAC ATTTTGGCGA GGAAGGTAAA	1200
GAGAGAAAGT AGTCCAGCAT GGCAGAGATG AAGACCGATG CCGTACCCT CGCGCAGGAG	1260
GCAGGTAATT TCGAGCGGAT CTCCGGCGAC CTGAAAACCC AGATCGACCA GGTGGAGTCG	1320
ACGGCAGGTT CGTTGCAGGG CCAGTGGCGC GGCGCGGCGG GGACGGCCGC CCAGGCCGCG	1380
GTGGTGCGCT TCCAAGAAGC AGCCAATAAG CAGAAGCAGG AACTCGACGA GATCTCGACG	1440
AATATTCGTC AGGCCGGCGT CCAATACTCG AGGGCCGACG AGGAGCAGCA GCAGGCGCTG	1500
TCCTCGCAA TGGGCTTCTG ACCCGCTAAT ACGAAAAGAA ACGGAGCAAA AACATGACAG	1560
AGCAGCAGTG GAATTTGCGG GGTATCGAGG CCGCGGCAAG CGCAATCCAG GGAAAT	1616

(2) INFORMATION FOR SEQ ID NO:108:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 432 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: single

CTAGTGGATG	GGACCATGGC	CATTTTCTGC	AGTCTCACTG	CCTTCTGTGT	TGACATTTTG	60
GCACGCCGGC	GGAAACGAAG	CACTGGGGTC	GAAGAACGGC	TGCGCTGCCA	TATCGTCCGG	120
AGCTTCCATA	CCTTCGTGCG	GCCGGAAGAG	CTTGTCGTAG	TCGGCCGCCA	TGACAACCTC	180
TCAGAGTGCG	CTCAAACGTA	TAAACACGAG	AAAGGGCGAG	ACCGACGGAA	GGTCGAACTC	240
GCCCGATCCC	GTGTTTCGCT	ATTCTACGCG	AACTCGGCGT	TGCCCTATGC	GAACATCCCA	300
GTGACGTTGC	CTTCGGTCGA	AGCCATTGCC	TGACCGGCTT	CGCTGATCGT	CCGCGCCAGG	360
TTCTGCAGCG	CGTTGTTTCT	CTCGGTAGCC	GTGGCGTCCC	ATTTTTGCTG	GACACCTTGG	420
TACGCCTCCG	AA					432

(A) LENGTH: 368 amino acids
(B) TYPE: amino acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

Met 1	Leu	Trp	His 5	Ala	Met	Pro	Pro	Glu	Xaa 10	Asn	Thr	Ala	Arg	Leu 15	Met
Ala	Gly	Ala	Gly 20	Pro	Ala	Pro	Met	Leu 25	Ala	Ala	Ala	Ala	Gly 30	Trp	Gln
Thr	Leu	Ser 35	Ala	Ala	Leu	Asp	Ala 40	Gln	Ala	Val	Glu	Leu 45	Thr	Ala	Arg
Leu	Asn 50	Ser	Leu	Gly	Glu	Ala 55	Trp	Thr	Gly	Gly	Gly 60	Ser	Asp	Lys	Ala
Leu 65	Ala	Ala	Ala	Thr	Pro 70	Met	Val	Val	Trp	Leu 75	Gln	Thr	Ala	Ser	Thr 80
Gln	Ala	Lys	Thr 85	Arg	Ala	Met	Gln	Ala 90	Thr	Ala	Gln	Ala	Ala	Ala 95	Tyr

144

Thr Gln Ala Met Ala Thr Thr Pro Ser Leu Pro Glu Ile Ala Ala Asn
 100 105 110
 His Ile Thr Gln Ala Val Leu Thr Ala Thr Asn Phe Phe Gly Ile Asn
 115 120 125
 Thr Ile Pro Ile Ala Leu Thr Glu Met Asp Tyr Phe Ile Arg Met Trp
 130 135 140
 Asn Gln Ala Ala Leu Ala Met Glu Val Tyr Gln Ala Glu Thr Ala Val
 145 150 155 160
 Asn Thr Leu Phe Glu Lys Leu Glu Pro Met Ala Ser Ile Leu Asp Pro
 165 170 175
 Gly Ala Ser Gln Ser Thr Thr Asn Pro Ile Phe Gly Met Pro Ser Pro
 180 185 190
 Gly Ser Ser Thr Pro Val Gly Gln Leu Pro Pro Ala Ala Thr Gln Thr
 195 200 205
 Leu Gly Gln Leu Gly Glu Met Ser Gly Pro Met Gln Gln Leu Thr Gln
 210 215 220
 Pro Leu Gln Gln Val Thr Ser Leu Phe Ser Gln Val Gly Gly Thr Gly
 225 230 235 240
 Gly Gly Asn Pro Ala Asp Glu Glu Ala Ala Gln Met Gly Leu Leu Gly
 245 250 255
 Thr Ser Pro Leu Ser Asn His Pro Leu Ala Gly Gly Ser Gly Pro Ser
 260 265 270
 Ala Gly Ala Gly Leu Leu Arg Ala Glu Ser Leu Pro Gly Ala Gly Gly
 275 280 285
 Ser Leu Thr Arg Thr Pro Leu Met Ser Gln Leu Ile Glu Lys Pro Val
 290 295 300
 Ala Pro Ser Val Met Pro Ala Ala Ala Ala Gly Ser Ser Ala Thr Gly
 305 310 315 320
 Gly Ala Ala Pro Val Gly Ala Gly Ala Met Gly Gln Gly Ala Gln Ser
 325 330 335
 Gly Gly Ser Thr Arg Pro Gly Leu Val Ala Pro Ala Pro Leu Ala Gln
 340 345 350
 Glu Arg Glu Glu Asp Asp Glu Asp Asp Trp Asp Glu Glu Asp Asp Trp
 355 360 365

(2) INFORMATION FOR SEQ ID NO:110:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 100 amino acids
- (B) TYPE: amino acid
- (C) STRANDEDNESS:
- (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:110:

```

Met Ala Glu Met Lys Thr Asp Ala Ala Thr Leu Ala Gln Glu Ala Gly
1           5           10           15

Asn Phe Glu Arg Ile Ser Gly Asp Leu Lys Thr Gln Ile Asp Gln Val
          20           25           30

Glu Ser Thr Ala Gly Ser Leu Gln Gly Gln Trp Arg Gly Ala Ala Gly
          35           40           45

Thr Ala Ala Gln Ala Ala Val Val Arg Phe Gln Glu Ala Ala Asn Lys
          50           55           60

Gln Lys Gln Glu Leu Asp Glu Ile Ser Thr Asn Ile Arg Gln Ala Gly
65           70           75           80

Val Gln Tyr Ser Arg Ala Asp Glu Glu Gln Gln Gln Ala Leu Ser Ser
          85           90           95

Gln Met Gly Phe
          100

```

(2) INFORMATION FOR SEQ ID NO:111:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 396 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:111:

```

GATCTCCGGC GACCTGAAAA CCCAGATCGA CCAGGTGGAG TCGACGGCAG GTTCGTTGCA      60
GGGCCAGTGG CGCGGCGCGG CGGGGACGGC CGCCCAGGCC GCGGTGGTGC GCTTCCAAGA      120
AGCAGCCAAT AAGCAGAAGC AGGAACTCGA CGAGATCTCG ACGAATATTC GTCAGGCCGG      180
CGTCCAATAC TCGAGGGCCG ACGAGGAGCA GCAGCAGGCG CTGTCCTCGC AAATGGGCTT      240

```

CTGACCCGCT AATACGAAAA GAAACGGAGC AAAACATGA CAGAGCAGCA GTGGAATTTC 300
 GCGGGTATCG AGGCCGCGGC AAGCGCAATC CAGGGAAATG TCACGTCCAT TCATTCCCTC 360
 CTTGACGAGG GGAAGCAGTC CCTGACCAAG CTCGCA 396

(2) INFORMATION FOR SEQ ID NO:112:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 80 amino acids
 (B) TYPE: amino acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:112:

Ile Ser Gly Asp Leu Lys Thr Gln Ile Asp Gln Val Glu Ser Thr Ala
 1 5 10 15
 Gly Ser Leu Gln Gly Gln Trp Arg Gly Ala Ala Gly Thr Ala Ala Gln
 20 25 30
 Ala Ala Val Val Arg Phe Gln Glu Ala Ala Asn Lys Gln Lys Gln Glu
 35 40 45
 Leu Asp Glu Ile Ser Thr Asn Ile Arg Gln Ala Gly Val Gln Tyr Ser
 50 55 60
 Arg Ala Asp Glu Glu Gln Gln Gln Ala Leu Ser Ser Gln Met Gly Phe
 65 70 75 80

(2) INFORMATION FOR SEQ ID NO:113:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 387 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:113:

GTGGATCCCG ATCCCGTGTT TCGCTATTCT ACGCGAACTC GCGGTTGCCC TATGCGAACA 60
 TCCCAGTGAC GTTGCCCTTCG GTCGAAGCCA TTGCCTGACC GGCTTCGCTG ATCGTCCGCG 120
 CCAGGTTCTG CAGCGCGTTG TTCAGCTCGG TAGCCGTGGC GTCCCATTTT TGCTGGACAC 180

CCTGGTACGC CTCCGAACCG CTACCGCCCC AGGCCGCTGC GAGCTTGGTC AGGGACTGCT 240
 TCCCCTCGTC AAGGAGGGAA TGAATGGACG TGACATTTCC CTGGATTGCG CTTGCCGCGG 300
 CCTCGATACC CGCGAAATTC CACTGCTGCT CTGTCATGTT TTTGCTCCGT TTCTTTTCGT 360
 ATTAGCGGGT CAGAAGCCCA TTTGCGA 387

(2) INFORMATION FOR SEQ ID NO:114:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 272 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:114:

CGGCACGAGG ATCTCGGTTG GCCCAACGGC GCTGGCGAGG GCTCCGTTCC GGGGGCGAGC 60
 TGC GCGCCGG ATGCTTCCTC TGCCCGCAGC CGCGCCTGGA TGGATGGACC AGTTGCTACC 120
 TTCCCACGT TTCGTTGGT GTCTGTGCGA TAGCGGTGAC CCCGGCGCGC ACGTCGGGAG 180
 TGTGGGGGG CAGGCCGGT CGGTGGTTTCG GCCGGGGACG CAGACGGTCT GGACGGAACG 240
 GCGGGGGGTT CGCCGATTGG CATCTTTGCC CA 272

(2) INFORMATION FOR SEQ ID NO:115:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 20 amino acids
 (B) TYPE: amino acid
 (C) STRANDEDNESS:
 (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:115:

Asp Pro Val Asp Ala Val Ile Asn Thr Thr Cys Asn Tyr Gly Gln Val
 1 5 10 15
 Val Ala Ala Leu
 20

(2) INFORMATION FOR SEQ ID NO:116:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 15 amino acids
- (B) TYPE: amino acid
- (C) STRANDEDNESS:
- (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:116:

Ala Val Glu Ser Gly Met Leu Ala Leu Gly Thr Pro Ala Pro Ser
1 5 10 15

(2) INFORMATION FOR SEQ ID NO:117:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 19 amino acids
- (B) TYPE: amino acid
- (C) STRANDEDNESS:
- (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:117:

Ala Ala Met Lys Pro Arg Thr Gly Asp Gly Pro Leu Glu Ala Ala Lys
1 5 10 15

Glu Gly Arg

(2) INFORMATION FOR SEQ ID NO:118:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 15 amino acids
- (B) TYPE: amino acid
- (C) STRANDEDNESS:
- (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:118:

Tyr Tyr Trp Cys Pro Gly Gln Pro Phe Asp Pro Ala Trp Gly Pro
1 5 10 15

(2) INFORMATION FOR SEQ ID NO:119:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 14 amino acids
(B) TYPE: amino acid
(C) STRANDEDNESS:
(D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:119:

Asp Ile Gly Ser Glu Ser Thr Glu Asp Gln Gln Xaa Ala Val
1 5 10

(2) INFORMATION FOR SEQ ID NO:120:

- (i) SEQUENCE CHARACTERISTICS:
- (A) LENGTH: 13 amino acids
 - (B) TYPE: amino acid
 - (C) STRANDEDNESS:
 - (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:120:

Ala Glu Glu Ser Ile Ser Thr Xaa Glu Xaa Ile Val Pro
1 5 10

(2) INFORMATION FOR SEQ ID NO:121:

- (i) SEQUENCE CHARACTERISTICS:
- (A) LENGTH: 17 amino acids
 - (B) TYPE: amino acid
 - (C) STRANDEDNESS:
 - (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:121:

Asp Pro Glu Pro Ala Pro Pro Val Pro Thr Thr Ala Ala Ser Pro Pro
1 5 10 15

Ser

(2) INFORMATION FOR SEQ ID NO:122:

- (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 15 amino acids

- (B) TYPE: amino acid
- (C) STRANDEDNESS:
- (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:122:

Ala Pro Lys Thr Tyr Xaa Glu Glu Leu Lys Gly Thr Asp Thr Gly
1 5 10 15

(2) INFORMATION FOR SEQ ID NO:123:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 30 amino acids
 - (B) TYPE: amino acid
 - (C) STRANDEDNESS:
 - (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:123:

Asp Pro Ala Ser Ala Pro Asp Val Pro Thr Ala Ala Gln Leu Thr Ser
1 5 10 15

Leu Leu Asn Ser Leu Ala Asp Pro Asn Val Ser Phe Ala Asn
 20 25 30

(2) INFORMATION FOR SEQ ID NO:124:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 22 amino acids
 - (B) TYPE: amino acid
 - (C) STRANDEDNESS:
 - (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:124:

Asp Pro Pro Asp Pro His Gln Xaa Asp Met Thr Lys Gly Tyr Tyr Pro
1 5 10 15

Gly Gly Arg Arg Xaa Phe
 20

(2) INFORMATION FOR SEQ ID NO:125:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 7 amino acids
 (B) TYPE: amino acid
 (C) STRANDEDNESS:
 (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:125:

Asp Pro Gly Tyr Thr Pro Gly
1 5

(2) INFORMATION FOR SEQ ID NO:126:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 10 amino acids
 (B) TYPE: amino acid
 (C) STRANDEDNESS:
 (D) TOPOLOGY: linear

(ix) FEATURE:

(D) OTHER INFORMATION: /note= "The Second Residue Can Be Either a Pro or Thr"

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:126:

Xaa Xaa Gly Phe Thr Gly Pro Gln Tyr
1 5 10

(2) INFORMATION FOR SEQ ID NO:127:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 9 amino acids
 (B) TYPE: amino acid
 (C) STRANDEDNESS:
 (D) TOPOLOGY: linear

(ix) FEATURE:

(D) OTHER INFORMATION: /note= "The Third Residue Can Be Either a Gln or Leu"

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:127:

Xaa Pro Xaa Val Thr Ala Tyr Ala Gly
1 5

(2) INFORMATION FOR SEQ ID NO:128:

- (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 9 amino acids
(B) TYPE: amino acid
(C) STRANDEDNESS:
(D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:128:

Xaa Xaa Xaa Glu Lys Pro Phe Leu Arg
1 5

(2) INFORMATION FOR SEQ ID NO:129:

- (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 15 amino acids
(B) TYPE: amino acid
(C) STRANDEDNESS:
(D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:129:

Xaa Asp Ser Glu Lys Ser Ala Thr Ile Lys Val Thr Asp Ala Ser
1 5 10 15

(2) INFORMATION FOR SEQ ID NO:130:

- (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 15 amino acids
(B) TYPE: amino acid
(C) STRANDEDNESS:
(D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:130:

Ala Gly Asp Thr Xaa Ile Tyr Ile Val Gly Asn Leu Thr Ala Asp
1 5 10 15

(2) INFORMATION FOR SEQ ID NO:131:

- (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 15 amino acids

- (B) TYPE: amino acid
- (C) STRANDEDNESS:
- (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:131:

Ala	Pro	Glu	Ser	Gly	Ala	Gly	Leu	Gly	Gly	Thr	Val	Gln	Ala	Gly
1				5				10						15

(2) INFORMATION FOR SEQ ID NO:132:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 21 amino acids
 - (B) TYPE: amino acid
 - (C) STRANDEDNESS:
 - (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:132:

Xaa	Tyr	Ile	Ala	Tyr	Xaa	Thr	Thr	Ala	Gly	Ile	Val	Pro	Gly	Lys	Ile
1				5					10					15	

Asn	Val	His	Leu	Val
				20

(2) INFORMATION FOR SEQ ID NO:133:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 882 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:133:

GCAACGCTGT	CGTGGCCTTT	GCGGTGATCG	GTTTCGCCTC	GCTGGCGGTG	GCGGTGGCGG	60
TCACCATCCG	ACCGACCGCG	GCCTCAAAAC	CGGTAGAGGG	ACACCAAAAC	GCCCAGCCAG	120
GGAAGTTCAT	GCCGTTGTTG	CCGACGCAAC	AGCAGGCGCC	GGTCCCGCCG	CCTCCGCCCG	180
ATGATCCCAC	CGCTGGATTC	CAGGGCGGCA	CCATTCCGGC	TGTACAGAAC	GTGGTGCCGC	240

GGCCGGGTAC CTCACCCGGG GTGGGTGGGA CGCCGGCTTC GCCTGCGCCG GAAGCGCCGG	300
CCGTGCCCCG TGTGTGCCT GCCCCGGTGC CAATCCCGGT CCCGATCATC ATTCCCCCGT	360
TCCCGGGTTG GCAGCCTGGA ATGCCGACCA TCCCCACCGC ACCGCCGACG ACGCCGGTGA	420
CCACGTCGGC GACGACGCCG CCGACCACGC CGCCGACCAC GCCGGTGACC ACGCCGCCAA	480
CGACGCCGCC GACCACGCCG GTGACCACGC CGCCAACGAC GCCGCCGACC ACGCCGGTGA	540
CCACGCCACC AACGACCGTC GCCCCGACGA CCGTCGCCCC GACGACGGTC GCTCCGACCA	600
CCGTGCCCCC GACCACGGTC GCTCCAGCCA CCGCCACGCC GACGACCGTC GCTCCGCAGC	660
CGACGCAGCA GCCCAGCAA CAACCAACCC AACAGATGCC AACCCAGCAG CAGACCGTGG	720
CCCCGCAGAC GGTGGCGCCG GCTCCGCAGC CGCCGTCCGG TGGCCGCAAC GGCAGCGGCG	780
GGGGCGACTT ATTGCGCGG TTCTGATCAC GGTGCGGGCT TCACTACGGT CGGAGGACAT	840
GGCCGGTGAT GCGGTGACGG TGGTGCTGCC CTGTCTCAAC GA	882

(2) INFORMATION FOR SEQ ID NO:134:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 815 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:134:

CCATCAACCA ACCGCTCGCG CCGCCCGCGC CGCCGGATCC GCCGTCGCCG CCACGCCCGC	60
CGGTGCCTCC GGTGCCCCCG TTGCCGCCGT CGCCGCCGTC GCCGCCGACC GGCTGGGTGC	120
CTAGGGCGCT GTTACCGCCC TGTTGGCGG GGACGCCGCC GGCACCACCG GTACCGCCGA	180
TGGCGCCGTT GCCGCCGGCG GCACCGTTGC CACCGTTGCC ACCGTTGCCA CCGTTGCCGA	240
CCAGCCACCC GCCGCGACCA CCGGCACCGC CGGCGCCGCC CGCACCGCCG GCGTGCCCGT	300
TCGTGCCCCG ACCGCCGGCA CCGCCGTTGC CGCCGTCACC GCCGACGGAA CTACCGGCGG	360
ACGCGGCCTG CCCGCCGGCG CCGCCCGCAC CGCCATTGGC ACCGCCGTCA CCGCCGGCTG	420
GGAGTGCCGC GATTAGGGCA CTGACCGGCG CAACCAGCGC AAGTACTCTC GGTCACCGAG	480
CACTTCCAGA CGACACCACA GCACGGGGTT GTCGGCGGAC TGGGTGAAAT GGCAGCCGAT	540

AGCGGCTAGC TGTCGGCTGC GGTCAACCTC GATCATGATG TCGAGGTGAC CGTGACCGCG 600
CCCCCGAAG GAGGCGCTGA ACTCGGCGTT GAGCCGATCG GCGATCGGTT GGGGCAGTGC 660
CCAGGCCAAT ACGGGGATAC CGGGTGTCTA AGCCGCCGCG AGCGCAGCTT CGGTTGCGCG 720
ACNGTGGTCG GGGTGGCCTG TTACGCCGTT GTCNTCGAAC ACGAGTAGCA GGTCTGCTCC 780
GGCGAGGGCA TCCACCACGC GTTGCCTCAG CTCGT 815

(2) INFORMATION FOR SEQ ID NO:135:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 1152 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:135:

ACCAGCCGCC GGCTGAGGTC TCAGATCAGA GAGTCTCCGG ACTCACCGGG GCGGTTTCAGC 60
CTTCTCCAG AACAACTGCT GAAGATCCTC GCCCGCGAAA CAGGCGCTGA TTTGACGCTC 120
TATGACCGGT TGAACGACGA GATCATCCGG CAGATTGATA TGGCACCCTG GGGCTAACAG 180
GTGCGCAAGA TGGTGCAGCT GTATGTCTCG GACTCCGTGT CGCGGATCAG CTTTGCCGAC 240
GGCCGGGTGA TCGTGTGGAG CGAGGAGCTC GGCGAGAGCC AGTATCCGAT CGAGACGCTG 300
GACGGCATCA CGCTGTTTGG GCGGCCGACG ATGACAACGC CCTTCATCGT TGAGATGCTC 360
AAGCGTGAGC GCGACATCCA GCTCTTCACG ACCGACGGCC ACTACCAGGG CCGGATCTCA 420
ACACCCGACG TGTCATACGC GCCGCGGCTC CGTCAGCAAG TTCACCGCAC CGACGATCCT 480
GCGTTCTGCC TGTCGTTAAG CAAGCGGATC GTGTCGAGGA AGATCCTGAA TCAGCAGGCC 540
TTGATTTCGGG CACACACGTC GGGGCAAGAC GTTGCTGAGA GCATCCGCAC GATGAAGCAC 600
TCGCTGGCCT GGGTCGATCG ATCGGGCTCC CTGGCGGAGT TGAACGGGTT CGAGGGAAAT 660
GCCGCAAAGG CATACTTCAC CGCGCTGGGG CATCTCGTCC CGCAGGAGTT CGCATTCCAG 720
GGCCGCTCGA CTCGGCCGCC GTTGACGCC TTCAACTCGA TGGTCAGCCT CGGCTATTCG 780
CTGCTGTACA AGAACATCAT AGGGGCGATC GAGCGTCACA GCCTGAACGC GTATATCGGT 840
TTCCTACACC AGGATTCACG AGGGCACGCA ACGTCTCGTG CCGAATTCGG CACGAGCTCC 900

GCTGAAACCG CTGGCCGGCT GCTCAGTGCC CGTACGTAAT CCGCTGCGCC CAGGCCGGCC 960
 CGCCGGCCGA ATACCAGCAG ATCGGACAGC GAATTGCCGC CCAGCCGGTT GGAGCCGTGC 1020
 ATACCGCCGG CACACTCACC GGCAGCGAAC AGGCCTGGCA CCGTGGCGGC GCCGGTGTCC 1080
 GCGTCTACTT CGACACCGCC CATCACGTAG TGACACGTCG GCCCGACTTC CATTGCCTGC 1140
 GTTCGGCACG AG 1152

(2) INFORMATION FOR SEQ ID NO:136:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 655 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:136:

CTCGTGCCGA TTCGGCAGGG TGTAATTGCC GGTGGTGTAN GCCGCATGAG TGCCGACGAC 60
 CAGCAATGCG GCAACAGCAC GGATCCCGGT CAACGACGCC ACCCGGTCCA CGTGGGCGAT 120
 CCGCTCGAGT CCGCCCTGGG CGGCTCTTTC CTTGGGCAGG GTCATCCGAC GTGTTTCCGC 180
 CGTGGTTTGC CGCCATTATG CCGGCGCGCC GCGTCGGGCG GCCGGTATGG CCGAANGTCG 240
 ATCAGCACAC CCGAGATACG GGTCTGTGCA AGCTTTTTGA GCGTCGCGCG GGGCAGCTTC 300
 GCCGGCAATT CTACTAGCGA GAAGTCTGGC CCGATACGGA TCTGACCGAA GTCGCTGCGG 360
 TGCAGCCCAC CCTCATTGGC GATGGCGCCG ACGATGGCGC CTGGACCGAT CTTGTGCCGC 420
 TTGCCGACGG CGACGCGGTA GGTGGTCAAG TCCGGTCTAC GCTTGGGCCT TTGCGGACGG 480
 TCCCGACGCT GGTCGCGGTT GCGCCGCGAA AGCGGCGGGT CGGGTGCCAT CAGGAATGCC 540
 TCACCGCCGC GGCAGTGCAC GGCCAGTGCC GCGGCGATGT CAGCCATCGG GACATCATGC 600
 TCGCGTTCAT ACTCCTCGAC CAGTCGGCGG AACAGCTCGA TTCCCGGACC GCCCA 655

(2) INFORMATION FOR SEQ ID NO:137:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 267 amino acids
- (B) TYPE: amino acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:137:

Asn	Ala	Val	Val	Ala	Phe	Ala	Val	Ile	Gly	Phe	Ala	Ser	Leu	Ala	Val	1	5	10	15
Ala	Val	Ala	Val	Thr	Ile	Arg	Pro	Thr	Ala	Ala	Ser	Lys	Pro	Val	Glu	20	25	30	
Gly	His	Gln	Asn	Ala	Gln	Pro	Gly	Lys	Phe	Met	Pro	Leu	Leu	Pro	Thr	35	40	45	
Gln	Gln	Gln	Ala	Pro	Val	Pro	Pro	Pro	Pro	Pro	Asp	Asp	Pro	Thr	Ala	50	55	60	
Gly	Phe	Gln	Gly	Gly	Thr	Ile	Pro	Ala	Val	Gln	Asn	Val	Val	Pro	Arg	65	70	75	80
Pro	Gly	Thr	Ser	Pro	Gly	Val	Gly	Gly	Thr	Pro	Ala	Ser	Pro	Ala	Pro	85	90	95	
Glu	Ala	Pro	Ala	Val	Pro	Gly	Val	Val	Pro	Ala	Pro	Val	Pro	Ile	Pro	100	105	110	
Val	Pro	Ile	Ile	Ile	Pro	Pro	Phe	Pro	Gly	Trp	Gln	Pro	Gly	Met	Pro	115	120	125	
Thr	Ile	Pro	Thr	Ala	Pro	Pro	Thr	Thr	Pro	Val	Thr	Thr	Ser	Ala	Thr	130	135	140	
Thr	Pro	Pro	Thr	Thr	Pro	Pro	Thr	Thr	Pro	Val	Thr	Thr	Pro	Pro	Thr	145	150	155	160
Thr	Pro	Pro	Thr	Thr	Pro	Val	Thr	Thr	Pro	Pro	Thr	Thr	Pro	Pro	Thr	165	170	175	
Thr	Pro	Val	Thr	Thr	Pro	Pro	Thr	Thr	Val	Ala	Pro	Thr	Thr	Val	Ala	180	185	190	
Pro	Thr	Thr	Val	Ala	Pro	Thr	Thr	Val	Ala	Pro	Thr	Thr	Val	Ala	Pro	195	200	205	
Ala	Thr	Ala	Thr	Pro	Thr	Thr	Val	Ala	Pro	Gln	Pro	Thr	Gln	Gln	Pro	210	215	220	
Thr	Gln	Gln	Pro	Thr	Gln	Gln	Met	Pro	Thr	Gln	Gln	Gln	Thr	Val	Ala	225	230	235	240
Pro	Gln	Thr	Val	Ala	Pro	Ala	Pro	Gln	Pro	Pro	Ser	Gly	Gly	Arg	Asn	245	250	255	

Gly Ser Gly Gly Gly Asp Leu Phe Gly Gly Phe
 260 265

(2) INFORMATION FOR SEQ ID NO:138:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 174 amino acids
- (B) TYPE: amino acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:138:

Ile Asn Gln Pro Leu Ala Pro Pro Ala Pro Pro Asp Pro Pro Ser Pro
 1 5 10 15
 Pro Arg Pro Pro Val Pro Pro Val Pro Pro Leu Pro Pro Ser Pro Pro
 20 25 30
 Ser Pro Pro Thr Gly Trp Val Pro Arg Ala Leu Leu Pro Pro Trp Leu
 35 40 45
 Ala Gly Thr Pro Pro Ala Pro Pro Val Pro Pro Met Ala Pro Leu Pro
 50 55 60
 Pro Ala Ala Pro Leu Pro Pro Leu Pro Pro Leu Pro Pro Leu Pro Thr
 65 70 75 80
 Ser His Pro Pro Arg Pro Pro Ala Pro Pro Ala Pro Pro Ala Pro Pro
 85 90 95
 Ala Cys Pro Phe Val Pro Val Pro Pro Ala Pro Pro Leu Pro Pro Ser
 100 105 110
 Pro Pro Thr Glu Leu Pro Ala Asp Ala Ala Cys Pro Pro Ala Pro Pro
 115 120 125
 Ala Pro Pro Leu Ala Pro Pro Ser Pro Pro Ala Gly Ser Ala Ala Ile
 130 135 140
 Arg Ala Leu Thr Gly Ala Thr Ser Ala Ser Thr Leu Gly His Arg Ala
 145 150 155 160
 Leu Pro Asp Asp Thr Thr Ala Arg Gly Cys Arg Arg Thr Gly
 165 170

(2) INFORMATION FOR SEQ ID NO:139:

(i) SEQUENCE CHARACTERISTICS:

159

- (A) LENGTH: 35 amino acids
- (B) TYPE: amino acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:139:

```

Gln Pro  Pro Ala Glu Val Ser Asp Gln Arg Val Ser Gly Leu Thr Gly
1          5          10          15
Ala Val  Gln Pro Ser Pro Arg Thr Thr Ala Glu Asp Pro Arg Pro Arg
          20          25          30
Asn Arg  Arg
          35

```

(2) INFORMATION FOR SEQ ID NO:140:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 104 amino acids
- (B) TYPE: amino acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:140:

```

Arg Ala Asp Ser Ala Gly Cys Thr Cys Arg Trp Cys Xaa Pro His Glu
1          5          10          15
Cys Arg Arg Pro Ala Met Arg Gln Gln His Gly Ser Arg Ser Thr Thr
          20          25          30
Pro Pro Gly Pro Arg Gly Arg Ser Ala Arg Val Arg Pro Gly Arg Leu
          35          40          45
Phe Pro Trp Ala Gly Ser Ser Asp Val Phe Pro Pro Trp Phe Ala Ala
          50          55          60
Ile Met Pro Ala Arg Arg Val Gly Arg Pro Val Trp Pro Xaa Val Asp
          65          70          75          80
Gln His Thr Arg Asp Thr Gly Leu Cys Lys Leu Phe Glu Arg Arg Ala
          85          90          95
Gly Gln Leu Arg Arg Gln Phe Tyr

```

100

(2) INFORMATION FOR SEQ ID NO:141:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 53 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: other nucleic acid

- (A) DESCRIPTION: /desc = "PCR primer"

(vi) ORIGINAL SOURCE:

- (A) ORGANISM: Mycobacterium tuberculosis

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:141:

GGATCCATAT GGGCCATCAT CATCATCATC ACGTGATCGA CATCATCGGG ACC

53

(2) INFORMATION FOR SEQ ID NO:142:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 42 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: other nucleic acid

- (A) DESCRIPTION: /desc = "PCR Primer"

(vi) ORIGINAL SOURCE:

- (A) ORGANISM: Mycobacterium tuberculosis

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:142:

CCTGAATTCA GGCCTCGGTT GCGCCGGCCT CATCTTGAAC GA

42

(2) INFORMATION FOR SEQ ID NO:143:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 31 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: other nucleic acid

- (A) DESCRIPTION: /desc = "PCR Primer"

(vi) ORIGINAL SOURCE:

- (A) ORGANISM: Mycobacterium tuberculosis

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:143:

GGATCCTGCA GGCTCGAAAC CACCGAGCGG T

31

(2) INFORMATION FOR SEQ ID NO:144:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 31 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: other nucleic acid

(A) DESCRIPTION: /desc = "PCR primer"

(vi) ORIGINAL SOURCE:

(A) ORGANISM: Mycobacterium tuberculosis

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:144:

CTCTGAATTC AGCGCTGGAA ATCGTCGCGA T

31

(2) INFORMATION FOR SEQ ID NO:145:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 33 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: other nucleic acid

(A) DESCRIPTION: /desc = "PCR primer"

(vi) ORIGINAL SOURCE:

(A) ORGANISM: Mycobacterium tuberculosis

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:145:

GGATCCAGCG CTGAGATGAA GACCGATGCC GCT

33

(2) INFORMATION FOR SEQ ID NO:146:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 33 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: other nucleic acid

(A) DESCRIPTION: /desc = "PCR primer"

(vi) ORIGINAL SOURCE:

(A) ORGANISM: Mycobacterium tuberculosis

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:146:

GAGAGAATTC TCAGAAGCCC ATTTGCGAGG ACA

33

(2) INFORMATION FOR SEQ ID NO:147:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 1993 base pairs

(B) TYPE: nucleic acid

(C) STRANDEDNESS: single

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

(vi) ORIGINAL SOURCE:

(A) ORGANISM: Mycobacterium tuberculosis

(ix) FEATURE:

(A) NAME/KEY: CDS

(B) LOCATION: 152..1273

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:147:

TGTTCTTCGA CGGCAGGCTG GTGGAGGAAG GGCCCACCGA ACAGCTGTTC TCCTCGCCGA	60
AGCATGCGGA AACCGCCCGA TACGTCGCCG GACTGTCGGG GGACGTCAAG GACGCCAAGC	120
GCGGAAATTG AAGAGCACAG AAAGGTATGG C GTG AAA ATT CGT TTG CAT ACG	172
Val Lys Ile Arg Leu His Thr	
1 5	
CTG TTG GCC GTG TTG ACC GCT GCG CCG CTG CTG CTA GCA GCG GCG GGC	220
Leu Leu Ala Val Leu Thr Ala Ala Pro Leu Leu Leu Ala Ala Ala Gly	
10 15 20	
TGT GGC TCG AAA CCA CCG AGC GGT TCG CCT GAA ACG GGC GCC GGC GCC	268
Cys Gly Ser Lys Pro Pro Ser Gly Ser Pro Glu Thr Gly Ala Gly Ala	
25 30 35	
GGT ACT GTC GCG ACT ACC CCC GCG TCG TCG CCG GTG ACG TTG GCG GAG	316
Gly Thr Val Ala Thr Thr Pro Ala Ser Ser Pro Val Thr Leu Ala Glu	
40 45 50 55	
ACC GGT AGC ACG CTG CTC TAC CCG CTG TTC AAC CTG TGG GGT CCG GCC	364
Thr Gly Ser Thr Leu Leu Tyr Pro Leu Phe Asn Leu Trp Gly Pro Ala	
60 65 70	

TTT CAC GAG AGG TAT CCG AAC GTC ACG ATC ACC GCT CAG GGC ACC GGT	412
Phe His Glu Arg Tyr Pro Asn Val Thr Ile Thr Ala Gln Gly Thr Gly	
75 80 85	
TCT GGT GCC GGG ATC GCG CAG GCC GCC GCC GGG ACG GTC AAC ATT GGG	460
Ser Gly Ala Gly Ile Ala Gln Ala Ala Ala Gly Thr Val Asn Ile Gly	
90 95 100	
GCC TCC GAC GCC TAT CTG TCG GAA GGT GAT ATG GCC GCG CAC AAG GGG	508
Ala Ser Asp Ala Tyr Leu Ser Glu Gly Asp Met Ala Ala His Lys Gly	
105 110 115	
CTG ATG AAC ATC GCG CTA GCC ATC TCC GCT CAG CAG GTC AAC TAC AAC	556
Leu Met Asn Ile Ala Leu Ala Ile Ser Ala Gln Gln Val Asn Tyr Asn	
120 125 130 135	
CTG CCC GGA GTG AGC GAG CAC CTC AAG CTG AAC GGA AAA GTC CTG GCG	604
Leu Pro Gly Val Ser Glu His Leu Lys Leu Asn Gly Lys Val Leu Ala	
140 145 150	
GCC ATG TAC CAG GGC ACC ATC AAA ACC TGG GAC GAC CCG CAG ATC GCT	652
Ala Met Tyr Gln Gly Thr Ile Lys Thr Trp Asp Asp Pro Gln Ile Ala	
155 160 165	
GCG CTC AAC CCC GGC GTG AAC CTG CCC GGC ACC GCG GTA GTT CCG CTG	700
Ala Leu Asn Pro Gly Val Asn Leu Pro Gly Thr Ala Val Val Pro Leu	
170 175 180	
CAC CGC TCC GAC GGG TCC GGT GAC ACC TTC TTG TTC ACC CAG TAC CTG	748
His Arg Ser Asp Gly Ser Gly Asp Thr Phe Leu Phe Thr Gln Tyr Leu	
185 190 195	
TCC AAG CAA GAT CCC GAG GGC TGG GGC AAG TCG CCC GGC TTC GGC ACC	796
Ser Lys Gln Asp Pro Glu Gly Trp Gly Lys Ser Pro Gly Phe Gly Thr	
200 205 210 215	
ACC GTC GAC TTC CCG GCG GTG CCG GGT GCG CTG GGT GAG AAC GGC AAC	844
Thr Val Asp Phe Pro Ala Val Pro Gly Ala Leu Gly Glu Asn Gly Asn	
220 225 230	
GGC GGC ATG GTG ACC GGT TGC GCC GAG ACA CCG GGC TGC GTG GCC TAT	892
Gly Gly Met Val Thr Gly Cys Ala Glu Thr Pro Gly Cys Val Ala Tyr	
235 240 245	
ATC GGC ATC AGC TTC CTC GAC CAG GCC AGT CAA CGG GGA CTC GGC GAG	940
Ile Gly Ile Ser Phe Leu Asp Gln Ala Ser Gln Arg Gly Leu Gly Glu	
250 255 260	
GCC CAA CTA GGC AAT AGC TCT GGC AAT TTC TTG TTG CCC GAC GCG CAA	988
Ala Gln Leu Gly Asn Ser Ser Gly Asn Phe Leu Pro Asp Ala Gln	
265 270 275	
AGC ATT CAG GCC GCG GCG GCT GGC TTC GCA TCG AAA ACC CCG GCG AAC	1036
Ser Ile Gln Ala Ala Ala Ala Gly Phe Ala Ser Lys Thr Pro Ala Asn	

280	285	290	295	
CAG GCG ATT TCG ATG ATC GAC GGG CCC GCC CCG GAC GGC TAC CCG ATC				1084
Gln Ala Ile Ser Met Ile Asp Gly Pro Ala Pro Asp Gly Tyr Pro Ile				
300	305	310		
ATC AAC TAC GAG TAC GCC ATC GTC AAC AAC CGG CAA AAG GAC GCC GCC				1132
Ile Asn Tyr Glu Tyr Ala Ile Val Asn Asn Arg Gln Lys Asp Ala Ala				
315	320	325		
ACC GCG CAG ACC TTG CAG GCA TTT CTG CAC TGG GCG ATC ACC GAC GGC				1180
Thr Ala Gln Thr Leu Gln Ala Phe Leu His Trp Ala Ile Thr Asp Gly				
330	335	340		
AAC AAG GCC TCG TTC CTC GAC CAG GTT CAT TTC CAG CCG CTG CCG CCC				1228
Asn Lys Ala Ser Phe Leu Asp Gln Val His Phe Gln Pro Leu Pro Pro				
345	350	355		
GCG GTG GTG AAG TTG TCT GAC GCG TTG ATC GCG ACG ATT TCC AGC				1273
Ala Val Val Lys Leu Ser Asp Ala Leu Ile Ala Thr Ile Ser Ser				
360	365	370		
TAGCCTCGTT GACCACCACG CGACAGCAAC CTCCGTCGGG CCATCGGGCT GCTTTGCGGA				
1333				
GCATGCTGGC CCGTGCCGGT GAAGTCGGCC GCGCTGGCCC GGCCATCCGG TGGTTGGGTG				
1393				
GGATAGGTGC GGTGATCCCG CTGCTTGCGC TGGTCTTGGT GCTGGTGGTG CTGGTCATCG				
1453				
AGGCGATGGG TGCGATCAGG CTCAACGGGT TGCATTTCTT CACCGCCACC GAATGGAATC				
1513				
CAGGCAACAC CTACGGCGAA ACCGTTGTCA CCGACGCGTC GCCCATCCGG TCGGCGCCTA				
1573				
CTACGGGGCG TTGCCGCTGA TCGTCGGGAC GCTGGCGACC TCGGCAATCG CCCTGATCAT				
1633				
CGCGGTGCCG GTCTCTGTAG GAGCGGCGCT GGTGATCGTG GAACGGCTGC CGAAACGGTT				
1693				
GGCCGAGGCT GTGGGAATAG TCCTGGAATT GCTCGCCGGA ATCCCCAGCG TGGTCGTCGG				
1753				
TTTGTGGGGG GCAATGACGT TCGGGCCGTT CATCGCTCAT CACATCGCTC CGGTGATCGC				
1813				
TCACAACGCT CCCGATGTGC CGGTGCTGAA CTA CTGCTGCGC GGCGACCCGG GCAACGGGGA				
1873				
GGGCATGTTG GTGTCCGGTC TGGTGTGGC GGTGATGGTC GTTCCCATTA TCGCCACCAC				
1933				
CACTCATGAC CTGTTCCGGC AGGTGCCGGT GTTGCCCCGG GAGGGCGCGA TCGGGAATTC				
1993				

(2) INFORMATION FOR SEQ ID NO:148:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 374 amino acids
- (B) TYPE: amino acid
- (D) TOPOLOGY: linear

165

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:148:

```

Val Lys Ile Arg Leu His Thr Leu Leu Ala Val Leu Thr Ala Ala Pro
 1           5           10           15
Leu Leu Leu Ala Ala Ala Gly Cys Gly Ser Lys Pro Pro Ser Gly Ser
          20           25           30
Pro Glu Thr Gly Ala Gly Ala Gly Thr Val Ala Thr Thr Pro Ala Ser
          35           40           45
Ser Pro Val Thr Leu Ala Glu Thr Gly Ser Thr Leu Leu Tyr Pro Leu
          50           55           60
Phe Asn Leu Trp Gly Pro Ala Phe His Glu Arg Tyr Pro Asn Val Thr
 65           70           75           80
Ile Thr Ala Gln Gly Thr Gly Ser Gly Ala Gly Ile Ala Gln Ala Ala
          85           90           95
Ala Gly Thr Val Asn Ile Gly Ala Ser Asp Ala Tyr Leu Ser Glu Gly
          100          105          110
Asp Met Ala Ala His Lys Gly Leu Met Asn Ile Ala Leu Ala Ile Ser
          115          120          125
Ala Gln Gln Val Asn Tyr Asn Leu Pro Gly Val Ser Glu His Leu Lys
          130          135          140
Leu Asn Gly Lys Val Leu Ala Ala Met Tyr Gln Gly Thr Ile Lys Thr
          145          150          155          160
Trp Asp Asp Pro Gln Ile Ala Ala Leu Asn Pro Gly Val Asn Leu Pro
          165          170          175
Gly Thr Ala Val Val Pro Leu His Arg Ser Asp Gly Ser Gly Asp Thr
          180          185          190
Phe Leu Phe Thr Gln Tyr Leu Ser Lys Gln Asp Pro Glu Gly Trp Gly
          195          200          205
Lys Ser Pro Gly Phe Gly Thr Thr Val Asp Phe Pro Ala Val Pro Gly
          210          215          220
Ala Leu Gly Glu Asn Gly Asn Gly Gly Met Val Thr Gly Cys Ala Glu
          225          230          235          240
Thr Pro Gly Cys Val Ala Tyr Ile Gly Ile Ser Phe Leu Asp Gln Ala
          245          250          255
Ser Gln Arg Gly Leu Gly Glu Ala Gln Leu Gly Asn Ser Ser Gly Asn
          260          265          270

```

166

Phe Leu Leu Pro Asp Ala Gln Ser Ile Gln Ala Ala Ala Gly Phe
 275 280 285
 Ala Ser Lys Thr Pro Ala Asn Gln Ala Ile Ser Met Ile Asp Gly Pro
 290 295 300
 Ala Pro Asp Gly Tyr Pro Ile Ile Asn Tyr Glu Tyr Ala Ile Val Asn
 305 310 315 320
 Asn Arg Gln Lys Asp Ala Ala Thr Ala Gln Thr Leu Gln Ala Phe Leu
 325 330 335
 His Trp Ala Ile Thr Asp Gly Asn Lys Ala Ser Phe Leu Asp Gln Val
 340 345 350
 His Phe Gln Pro Leu Pro Pro Ala Val Val Lys Leu Ser Asp Ala Leu
 355 360 365
 Ile Ala Thr Ile Ser Ser
 370

(2) INFORMATION FOR SEQ ID NO:149:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 1993 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:149:

TGTTCTTCGA CGGCAGGCTG GTGGAGGAAG GGCCACCGA ACAGCTGTTC TCCTCGCCGA	60
AGCATGCGGA AACCGCCCGA TACGTCGCCG GACTGTCGGG GGACGTCAAG GACGCCAAGC	120
GCGGAAATTG AAGAGCACAG AAAGGTATGG CGTGAAAATT CGTTTGCATA CGCTGTTGGC	180
CGTGTTGACC GCTGCGCCGC TGCTGCTAGC AGCGGCGGGC TGTGGCTCGA AACCACCGAG	240
CGGTTCGCCT GAAACGGGCG CCGGCGCCGG TACTGTCGCG ACTACCCCGG CGTCGTCGCC	300
GGTGACGTTG GCGGAGACCG GTAGCACGCT GCTCTACCCG CTGTTCAACC TGTGGGGTCC	360
GGCCTTTCAC GAGAGGTATC CGAACGTCAC GATCACCGCT CAGGGCACCG GTTCTGGTGC	420
CGGGATCGCG CAGGCCGCCG CCGGGACGGT CAACATTGGG GCCTCCGACG CCTATCTGTC	480
GGAAGGTGAT ATGGCCGCGC ACAAGGGGCT GATGAACATC GCGCTAGCCA TCTCCGCTCA	540
GCAGGTCAAC TACAACCTGC CCGGAGTGAG CGAGCACCTC AAGCTGAACG GAAAAGTCCT	600

GGCGGCCATG TACCAGGGCA CCATCAAAAC CTGGGACGAC CCGCAGATCG CTGCGCTCAA	660
CCCCGGCGTG AACCTGCCCC GCACCGCGGT AGTTCCGCTG CACCGCTCCG ACGGGTCCGG	720
TGACACCTTC TTGTTACCC AGTACCTGTC CAAGCAAGAT CCCGAGGGCT GGGGCAAGTC	780
GCCCCGCTTC GGCACCACCG TCGACTTCCC GGCGGTGCCG GGTGCGCTGG GTGAGAACGG	840
CAACGGCGGC ATGGTGACCG GTTGCGCCGA GACACGGGC TCGGTGGCCT ATATCGGCAT	900
CAGCTTCCTC GACCAGGCCA GTCAACGGGG ACTCGGCGAG GCCCAACTAG GCAATAGCTC	960
TGGCAATTTC TTGTTGCCCG ACGCGCAAAG CATTGAGGCC GCGGCGGCTG GCTTCGCATC	1020
GAAAACCCCG GCGAACCAGG CGATTTTCGAT GATCGACGGG CCCGCCCCGG ACGGCTACCC	1080
GATCATCAAC TACGAGTACG CCATCGTCAA CAACCGGCAA AAGGACGCCG CCACCGCGCA	1140
GACCTTGCA GCAATTTCTGC ACTGGGCGAT CACCGACGGC AACAAGGCCT CGTTCCTCGA	1200
CCAGGTTTCAT TTCCAGCCGC TGCCGCCCGC GGTGGTGAAG TTGTCTGACG CGTTGATCGC	1260
GACGATTTCC AGCTAGCCTC GTTGACCACC ACGCGACAGC AACCTCCGTC GGGCCATCGG	1320
GCTGCTTTGC GGAGCATGCT GGCCCGTGCC GGTGAAGTCG GCCGCGCTGG CCCGGCCATC	1380
CGGTGGTTGG GTGGGATAGG TGCGGTGATC CCGCTGCTTG CGCTGGTCTT GGTGCTGGTG	1440
GTGCTGGTCA TCGAGGCGAT GGTGCGATC AGGCTCAACG GGTGTCATTT CTTACCGCC	1500
ACCGAATGGA ATCCAGGCAA CACCTACGGC GAAACCGTTG TCACCGACGC GTCGCCCATC	1560
CGGTGCGCGC CTACTACGGG GCGTTGCCGC TGATCGTCGG GACGCTGGCG ACCTCGGCAA	1620
TCGCCCTGAT CATCGCGGTG CCGGTCTCTG TAGGAGCGGC GCTGGTGATC GTGGAACGGC	1680
TGCCGAAACG GTTGGCCGAG GCTGTGGGAA TAGTCCTGGA ATTGCTCGCC GGAATCCCCA	1740
GCGTGGTCGT CGGTTTGTGG GGGGCAATGA CGTTCGGGCC GTTCATCGCT CATCACATCG	1800
CTCCGGTGAT CGCTCACAAC GCTCCCGATG TGCCGGTGCT GAACTACTTG CGCGGCGACC	1860
CGGGCAACGG GGAGGGCATG TTGGTGTCCG GTCTGGTGTT GGCGGTGATG GTCGTTCCCA	1920
TTATCGCCAC CACCACTCAT GACCTGTTCC GGCAGGTGCC GGTGTTGCCC CGGGAGGGCG	1980
CGATCGGGAA TTC	1993

(2) INFORMATION FOR SEQ ID NO:150:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 374 amino acids
- (B) TYPE: amino acid
- (C) STRANDEDNESS:
- (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:150:

Met	Lys	Ile	Arg	Leu	His	Thr	Leu	Leu	Ala	Val	Leu	Thr	Ala	Ala	Pro	1	5	10	15
Leu	Leu	Leu	Ala	Ala	Ala	Gly	Cys	Gly	Ser	Lys	Pro	Pro	Ser	Gly	Ser	20	25	30	
Pro	Glu	Thr	Gly	Ala	Gly	Ala	Gly	Thr	Val	Ala	Thr	Thr	Pro	Ala	Ser	35	40	45	
Ser	Pro	Val	Thr	Leu	Ala	Glu	Thr	Gly	Ser	Thr	Leu	Leu	Tyr	Pro	Leu	50	55	60	
Phe	Asn	Leu	Trp	Gly	Pro	Ala	Phe	His	Glu	Arg	Tyr	Pro	Asn	Val	Thr	65	70	75	80
Ile	Thr	Ala	Gln	Gly	Thr	Gly	Ser	Gly	Ala	Gly	Ile	Ala	Gln	Ala	Ala	85	90	95	
Ala	Gly	Thr	Val	Asn	Ile	Gly	Ala	Ser	Asp	Ala	Tyr	Leu	Ser	Glu	Gly	100	105	110	
Asp	Met	Ala	Ala	His	Lys	Gly	Leu	Met	Asn	Ile	Ala	Leu	Ala	Ile	Ser	115	120	125	
Ala	Gln	Gln	Val	Asn	Tyr	Asn	Leu	Pro	Gly	Val	Ser	Glu	His	Leu	Lys	130	135	140	
Leu	Asn	Gly	Lys	Val	Leu	Ala	Ala	Met	Tyr	Gln	Gly	Thr	Ile	Lys	Thr	145	150	155	160
Trp	Asp	Asp	Pro	Gln	Ile	Ala	Ala	Leu	Asn	Pro	Gly	Val	Asn	Leu	Pro	165	170	175	
Gly	Thr	Ala	Val	Val	Pro	Leu	His	Arg	Ser	Asp	Gly	Ser	Gly	Asp	Thr	180	185	190	
Phe	Leu	Phe	Thr	Gln	Tyr	Leu	Ser	Lys	Gln	Asp	Pro	Glu	Gly	Trp	Gly	195	200	205	
Lys	Ser	Pro	Gly	Phe	Gly	Thr	Thr	Val	Asp	Phe	Pro	Ala	Val	Pro	Gly	210	215	220	
Ala	Leu	Gly	Glu	Asn	Gly	Asn	Gly	Gly	Met	Val	Thr	Gly	Cys	Ala	Glu	225	230	235	240
Thr	Pro	Gly	Cys	Val	Ala	Tyr	Ile	Gly	Ile	Ser	Phe	Leu	Asp	Gln	Ala	245	250	255	

169

Ser Gln Arg Gly Leu Gly Glu Ala Gln Leu Gly Asn Ser Ser Gly Asn
 260 265 270
 Phe Leu Leu Pro Asp Ala Gln Ser Ile Gln Ala Ala Ala Gly Phe
 275 280 285
 Ala Ser Lys Thr Pro Ala Asn Gln Ala Ile Ser Met Ile Asp Gly Pro
 290 295 300
 Ala Pro Asp Gly Tyr Pro Ile Ile Asn Tyr Glu Tyr Ala Ile Val Asn
 305 310 315 320
 Asn Arg Gln Lys Asp Ala Ala Thr Ala Gln Thr Leu Gln Ala Phe Leu
 325 330 335
 His Trp Ala Ile Thr Asp Gly Asn Lys Ala Ser Phe Leu Asp Gln Val
 340 345 350
 His Phe Gln Pro Leu Pro Pro Ala Val Val Lys Leu Ser Asp Ala Leu
 355 360 365
 Ile Ala Thr Ile Ser Ser
 370

(2) INFORMATION FOR SEQ ID NO:151:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 1777 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:151:

GGTCTTGACC ACCACCTGGG TGTCGAAGTC GGTGCCCCGA TTGAAGTCCA GGTACTCGTG	60
GGTGGGGCGG GCGAAACAAT AGCGACAAGC ATGCGAGCAG CCGCGGTAGC CGTTGACGGT	120
GTAGCGAAAC GGCAACGCGG CCGCGTTGGG CACCTTGTTT AGCGCTGATT TGCACAACAC	180
CTCGTGGAAG GTGATGCCGT CGAATTGTGG CGCGCGAACG CTGCGGACCA GGCCGATCCG	240
CTGCAACCCG GCAGCGCCCG TCGTCAACGG GCATCCCGTT CACCGCGACG GCTTGCCGGG	300
CCCAACGCAT ACCATTATTC GAACAACCGT TCTATACTTT GTCAACGCTG GCCGCTACCG	360
AGCGCCGCAC AGGATGTGAT ATGCCATCTC TGCCCGCACA GACAGGAGCC AGGCCTTATG	420
ACAGCATTCG GCGTCGAGCC CTACGGGCAG CCGAAGTACC TAGAAATCGC CGGGAAGCGC	480
ATGGCGTATA TCGACGAAGG CAAGGGTGAC GCCATCGTCT TTCAGCACGG CAACCCACG	540

TCGTCTTACT TGTGGCGCAA CATCATGCCG CACTTGGAAG GGCTGGGCCG GCTGGTGGCC	600
TGCGATCTGA TCGGGATGGG CGCGTCGGAC AAGCTCAGCC CATCGGGACC CGACCGCTAT	660
AGCTATGGCG AGCAACGAGA CTTTTTGTTC GCGCTCTGGG ATGCGCTCGA CCTCGGCGAC	720
CACGTGGTAC TGGTGCTGCA CGACTGGGGC TCGGCGCTCG GCTTCGACTG GGCTAACCAG	780
CATCGCGACC GAGTGCAGGG GATCGCGTTC ATGGAAGCGA TCGTCACCCC GATGACGTGG	840
GCGGACTGGC CGCCGGCCGT GCGGGGTGTG TTCCAGGGTT TCCGATCGCC TCAAGGCGAG	900
CCAATGGCGT TGGAGCACAA CATCTTTGTC GAACGGGTGC TGCCCGGGGC GATCCTGCGA	960
CAGCTCAGCG ACGAGGAAAT GAACCACTAT CGGCGGCCAT TCGTGAACGG CGGCGAGGAC	1020
CGTCGCCCCA CGTTGTCGTG GCCACGAAAC CTTCCAATCG ACGGTGAGCC CGCCGAGGTC	1080
GTCGCGTTGG TCAACGAGTA CCGGAGCTGG CTCGAGGAAA CCGACATGCC GAAACTGTTC	1140
ATCAACGCCG AGCCCGGCGC GATCATCACC GGCCGCATCC GTGACTATGT CAGGAGCTGG	1200
CCCAACCAGA CCGAAATCAC AGTGCCCGGC GTGCATTTTCG TTCAGGAGGA CAGCGATGGC	1260
GTCGTATCGT GGGCGGGCGC TCGGCAGCAT CGGCGACCTG GGAGCGCTCT CATTTACGA	1320
GACCAAGAAT GTGATTTCCG GCGAAGGCGG CGCCCTGCTT GTCAACTCAT AAGACTTCCT	1380
GCTCCGGGCA GAGATTCTCA GGGAAAAGG CACCAATCGC AGCCGCTTCC TTCGCAACGA	1440
GGTCGACAAA TATACGTGGC AGGACAAAGG TCTTCTATT TGCCAGCGA ATTAGTCGCT	1500
GCCTTTCTAT GGGCTCAGTT CGAGGAAGCC GAGCGGATCA CGCGTATCCG ATTGGACCTA	1560
TGGAACCGGT ATCATGAAAG CTTCGAATCA TTGGAACAGC GGGGGCTCCT GCGCCGTCCG	1620
ATCATCCAC AGGGCTGCTC TCACAACGCC CACATGTACT ACGTGTTACT AGCGCCCAGC	1680
GCCGATCGGG AGGAGGTGCT GCGCGTCTG ACGAGCGAAG GTATAGGCGC GGTCTTTCAT	1740
TACGTGCCGC TTCACGATTC GCCGGCCGGG CGTCGCT	1777

(2) INFORMATION FOR SEQ ID NO:152:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 324 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:152:

GAGATTGAAT CGTACCGGTC TCCTTAGCGG CTCCGTCCCG TGAATGCCCA TATCACGCAC	60
GGCCATGTTC TGGCTGTCGA CCTTCGCCCC ATGCCCGGAC GTTGGTAAAC CCAGGGTTTG	120
ATCAGTAATT CCGGGGACG GTTGCGGGAA GGCGCCAGG ATGTGCGTGA GCCGCGCGC	180
CGCCGTCGCC CAGGCGACCG CTGGATGCTC AGCCCCGGTG CGGCGACGTA GCCAGCGTTT	240
GGCGCGTGTC GTCCACAGTG GTACTCCGGT GACGACGCGG CGCGGTGCCT GGGTGAAGAC	300
CGTGACCGAC GCCGCCGATT CAGA	324

(2) INFORMATION FOR SEQ ID NO:153:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 1338 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:153:

GCGGTACCGC CGCGTTGCGC TGGCACGGGA CCTGTACGAC CTGAACCACT TCGCCTCGCG	60
AACGATTGAC GAACCGCTCG TGCGGCGGCT GTGGGTGCTC AAGGTGTGGG GTGATGTCGT	120
CGATGACCGG CGCGGCACCC GGCCACTACG CGTCGAAGAC GTCCTCGCCG CCCGCAGCGA	180
GCACGACTTC CAGCCCGACT CGATCGGCGT GCTGACCCGT CCTGTGCTA TGGCTGCCTG	240
GGAAGCTCGC GTTCGGAAGC GATTTGCGTT CCTCACTGAC CTCGACGCCG ACGAGCAGCG	300
GTGGGCCGCC TGCACGAAC GGCACCGCCG CGAAGTGGAG AACGCGCTGG CGGTGCTGCG	360
GTCCTGATCA ACCTGCCGGC GATCGTGCCG TTCCGCTGGC ACGGTTGCGG CTGGACGCGG	420
CTGAATCGAC TAGATGAGAG CAGTTGGGCA CGAATCCGGC TGTGGTGGTG AGCAAGACAC	480
GAGTACTGTC ATCACTATTG GATGCACTGG ATGACCGGCC TGATTCAGCA GGACCAATGG	540
AACTGCCCCG GGCAAAACGT CTCGGAGATG ATCGGCGTCC CCTCGGAACC CTGCGGTGCT	600
GGCGTCATTC GGACATCGGT CCGGCTCGCG GGATCGTGGT GACGCCAGCG CTGAAGGAGT	660
GGAGCGCGGC GGTGCACGCG CTGCTGGACG GCCGGCAGAC GGTGCTGCTG CGTAAGGGCG	720
GGATCGGCGA GAAGCGCTTC GAGGTGGCGG CCCACGAGTT CTTGTTGTTC CCGACGGTCG	780
CGCACAGCCA CGCCGAGCGG GTTCGCCCCG AGCACCGCGA CCTGCTGGGC CCGGCGGCCG	840

CCGACAGCAC CGACGAGTGT GTGCTACTGC GGGCCGCAGC GAAAGTTGTT GCCGCACTGC	900
CGGTAAACCG GCCAGAGGGT CTGGACGCCA TCGAGGATCT GCACATCTGG ACCGCCGAGT	960
CGGTGCGCGC CGACCGGCTC GACTTTCGGC CCAAGCACAA ACTGGCCGTC TTGGTGGTCT	1020
CGGCGATCCC GCTGGCCGAG CCGGTCCGGC TGGCGCGTAG GCCCAGGTAC GGCGGTTGCA	1080
CCAGCTGGGT GCAGCTGCCG GTGACGCCGA CGTTGGCGGC GCCGGTGAC GACGAGGCCG	1140
CGCTGGCCGA GGTGCGCCGC CGGGTCCGCG AGGCCGTGGG TTGACTGGGC GGCATCGCTT	1200
GGGTCTGAGC TGTACGCCCA GTCGGCGCTG CGAGTGATCT GCTGTCGGTT CGGTCCCTGC	1260
TGGCGTCAAT TGACGGCGCG GGCAACAGCA GCATTGGCGG CGCCATCCTC CGCGCGGCCG	1320
GCGCCACCG CTACAACC	1338

(2) INFORMATION FOR SEQ ID NO:154:

- (i) SEQUENCE CHARACTERISTICS:
- (A) LENGTH: 321 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:154:

CCGGCGGCAC CGGCGGCACC GCGGTACCG GCGGCAACGG CGCTGACGCC GCTGCTGTGG	60
TGGGCTTCGG CGCGAACGGC GACCCTGGCT TCGTGGCGG CAAAGGCGGT AACGGCGGAA	120
TAGGTGGGGC CGCGGTGACA GCGGGGTCG CCGGCGACGG CGGCACCGGC GGCAAAGGTG	180
GCACCGGCGG TGCCGGCGGC GCCGGCAACG ACGCCGGCAG CACCGGCAAT CCCGGCGGTA	240
AGGGCGGCGA CGGCGGGATC GCGGTGCCG GCGGGGCCG CGGCGCGGCC GGCACCGGCA	300
ACGGCGGCCA TGCCGGCAAC C	321

(2) INFORMATION FOR SEQ ID NO:155:

- (i) SEQUENCE CHARACTERISTICS:
- (A) LENGTH: 492 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:155:

```

GAAGACCCGG CCCC GCCATA TCGATCGGCT CGCCGACTAC TTTCGCCGAA CGTGCACGCG      60
GCGGCGTCGG GCTGATCATC ACCGGTGGCT ACGCGCCCAA CCGCACCGGA TGGCTGCTGC      120
CGTTCGCCTC CGAACTCGTC ACTTCGGCGC AAGCCCACG GCACCGCCGA ATCACCAGGG      180
CGGTCCACGA TTCGGGTGCA AAGATCCTGC TGCAAATCCT GCACGCCGGA CGCTACGCCT      240
ACCACCCACT TCGGGTCAGC GCCTCGCCGA TCAAGGCGCC GATCACCCCG TTTCGTCCGC      300
GAGCACTATC GGCTCGCGGG GTCGAAGCGA CCATCGCGGA TTTCGCCCGC TGCGCGCAGT      360
TGGCCCGCGA TGCCGGCTAC GACGGCGTCG AAATCATGGG CAGCGAAGGG TATCTGCTCA      420
ATCAGTTCCT GGCGCCGCGC ACCAACAAGC GCACCGACTC GTGGGGCGGC ACACCGGCCA      480
ACCGTCGCCG GT                                         492

```

(2) INFORMATION FOR SEQ ID NO:156:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 536 amino acids
- (B) TYPE: amino acid
- (C) STRANDEDNESS:
- (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:156:

```

Phe Ala Gln His Leu Val Glu Gly Asp Ala Val Glu Leu Trp Arg Ala
1           5           10           15
Asn Ala Ala Asp Gln Ala Asp Pro Leu Gln Pro Gly Ser Ala Arg Arg
20          25          30
Gln Arg Ala Ser Arg Ser Pro Arg Arg Leu Ala Gly Pro Asn Ala Tyr
35          40          45
His Tyr Ser Asn Asn Arg Ser Ile Leu Cys Gln Arg Trp Pro Leu Pro
50          55          60
Ser Ala Ala Gln Asp Val Ile Cys His Leu Cys Pro His Arg Gln Glu
65          70          75          80
Pro Gly Leu Met Thr Ala Phe Gly Val Glu Pro Tyr Gly Gln Pro Lys
85          90          95
Tyr Leu Glu Ile Ala Gly Lys Arg Met Ala Tyr Ile Asp Glu Gly Lys

```


175

```

Ile Arg Leu Pro Ala Pro Gly Arg Asp Ser Gln Gly Lys Gly His Gln
      405                      410                      415

Ser Gln Pro Leu Pro Ser Gln Arg Gly Arg Gln Ile Tyr Val Ala Gly
      420                      425                      430

Gln Arg Ser Ser Tyr Leu Pro Ser Glu Leu Val Ala Ala Phe Leu Trp
      435                      440                      445

Ala Gln Phe Glu Glu Ala Glu Arg Ile Thr Arg Ile Arg Leu Asp Leu
      450                      455                      460

Trp Asn Arg Tyr His Glu Ser Phe Glu Ser Leu Glu Gln Arg Gly Leu
      465                      470                      475                      480

Leu Arg Arg Pro Ile Ile Pro Gln Gly Cys Ser His Asn Ala His Met
      485                      490                      495

Tyr Tyr Val Leu Leu Ala Pro Ser Ala Asp Arg Glu Glu Val Leu Ala
      500                      505                      510

Arg Leu Thr Ser Glu Gly Ile Gly Ala Val Phe His Tyr Val Pro Leu
      515                      520                      525

His Asp Ser Pro Ala Gly Arg Arg
      530                      535

```

(2) INFORMATION FOR SEQ ID NO:157:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 284 amino acids
- (B) TYPE: amino acid
- (C) STRANDEDNESS:
- (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:157:

```

Asn Glu Ser Ala Pro Arg Ser Pro Met Leu Pro Ser Ala Arg Pro Arg
1              5              10              15

Tyr Asp Ala Ile Ala Val Leu Leu Asn Glu Met His Ala Gly His Cys
      20              25              30

Asp Phe Gly Leu Val Gly Pro Ala Pro Asp Ile Val Thr Asp Ala Ala
      35              40              45

Gly Asp Asp Arg Ala Gly Leu Gly Val Asp Glu Gln Phe Arg His Val
      50              55              60

Gly Phe Leu Glu Pro Ala Pro Val Leu Val Asp Gln Arg Asp Asp Leu

```

176

65		70		75		80
Gly Gly Leu Thr Val Asp Trp Lys Val Ser Trp Pro Arg Gln Arg Gly						
	85			90		95
Ala Thr Val Leu Ala Ala Val His Glu Trp Pro Pro Ile Val Val His						
	100		105			110
Phe Leu Val Ala Glu Leu Ser Gln Asp Arg Pro Gly Gln His Pro Phe						
	115		120			125
Asp Lys Asp Val Val Leu Gln Arg His Trp Leu Ala Leu Arg Arg Ser						
	130		135		140	
Glu Thr Leu Glu His Thr Pro His Gly Arg Arg Pro Val Arg Pro Arg						
145		150		155		160
His Arg Gly Asp Asp Arg Phe His Glu Arg Asp Pro Leu His Ser Val						
	165		170			175
Ala Met Leu Val Ser Pro Val Glu Ala Glu Arg Arg Ala Pro Val Val						
	180		185			190
Gln His Gln Tyr His Val Val Ala Glu Val Glu Arg Ile Pro Glu Arg						
	195		200			205
Glu Gln Lys Val Ser Leu Leu Ala Ile Ala Ile Ala Val Gly Ser Arg						
	210		215		220	
Trp Ala Glu Leu Val Arg Arg Ala His Pro Asp Gln Ile Ala Gly His						
225		230		235		240
Gln Pro Ala Gln Pro Phe Gln Val Arg His Asp Val Ala Pro Gln Val						
	245		250			255
Arg Arg Arg Gly Val Ala Val Leu Lys Asp Asp Gly Val Thr Leu Ala						
	260		265			270
Phe Val Asp Ile Arg His Ala Leu Pro Gly Asp Phe						
	275		280			

(2) INFORMATION FOR SEQ ID NO:158:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 264 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:158:

ATGAACATGT	CGTCGGTGGT	GGGTCGCAAG	GCCTTTGCGC	GATTCGCCGG	CTACTCCTCC	60
GCCATGCACG	CGATCGCCGG	TTTCTCCGAT	GCGTTGCGCC	AAGAGCTGCG	GGGTAGCGGA	120
ATCGCCGTCT	CGGTGATCCA	CCCGGCGCTG	ACCCAGACAC	CGCTGTTGGC	CAACGTCGAC	180
CCCGCCGACA	TGCCGCCGCC	GTTTCGCAGC	CTCAGCCCCA	TTCCCGTTCA	CTGGGTCGCG	240
GCAGCGGTGC	TTGACGGTGT	GGCG				264

(2) INFORMATION FOR SEQ ID NO:159:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 1171 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:159:

TAGTCGGCGA	CGATGACGTC	GCGGTCCAGG	CCGACCGCTT	CAAGCACCAG	CGCGACCACG	60
AAGCCGGTGC	GATCCTTACC	CGCGAAGCAG	TGGGTGAGCA	CCGGGCGTCC	GGCGGCAAGC	120
AGTGTGACGA	CACGATGTAG	CGCGCGCTGT	GCTCCATTGC	GCGTTGGGAA	TTGGCGATAC	180
TCGTCGGTCA	TGTAGCGGGT	GGCCGCGTCA	TTTATCGACT	GGCTGGATTC	GCCGGACTCG	240
CCGTTGGACC	CGTCATTGGT	TAGCAGCCTC	TTGAATGCGG	TTTCGTGCGG	CGCTGAGTCG	300
TCGGCGTCAT	CATCGGCGAG	GTCGGGGAAC	GGCAGCAGGT	GGACGTCGAT	GCCGTCCGGA	360
ACCCGTCCCTG	GACCGCGGCG	GGCAACCTCC	CGGGACGACC	GCAGGTCGGC	AACGTCGGTG	420
ATCCCCAGCC	GGCGCAGCGT	TGCCCCCTCGT	GCCGAATTCTG	GCACGAGGCT	GGCGAGCCAC	480
CGGGCATCAC	CAAGCAACGC	TTGCCAGTA	CGGATCGTCA	CTTCCGCATC	CGGCAGACCA	540
ATCTCCTCGC	CGCCCATCGT	CAGATCCCGC	TCGTGCGTTG	ACAAGAACGG	CCGCAGATGT	600
GCCAGCGGGT	ATCGGAGATT	GAACCGCGCA	CGCAGTTCTT	CAATCGCTGC	GCGCTGCCGC	660
ACTATTGGCA	CTTCCGGCG	GTCGCGGTAT	TCAGCAAGCA	TGCGAGTCTC	GACGAACTCG	720
CCCCACGTAA	CCCACGGCGT	AGTCCCGGC	GTGACGCGGA	GGATCGGCGG	GTGATCTTTG	780
CCGCCACGCT	CGTAGCCGTT	GATCCACCGC	TTCGCGGTGC	CGGCGGGGAG	GCCGATCAGC	840
TTATCGACCT	CGGCGTATGC	CGACGGCAAG	CTGGGCGCGT	TCGTGAGGT	CAAGAACTCC	900
ACCATCGGCA	CCGGCACCAA	GGTGCCGCAC	CTGACCTACG	TCGGCGACGC	CGACATCGGC	960

GAGTACAGCA ACATCGGCGC CTCCAGCGTG TTCGTCAACT ACGACGGTAC GTCCAAACGG 1020
CGCACCACCG TCGGTTCGCA CGTACGGACC GGGTCCGACA CCATGTTTCGT GGCCCCAGTA 1080
ACCATCGGCG ACGGCGCGTA TACCGGGGCC GGCACAGTGG TGCGGGAGGA TGTCCCGCCG 1140
GGGGCGCTGG CAGTGTCGGC GGGTCCGCAA C 1171

(2) INFORMATION FOR SEQ ID NO:160:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 227 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:160:

GCAAAGGCGG CACCGGCGGG GCCGGCATGA ACAGCCTCGA CCCGCTGCTA GCCGCCCAAG 60
ACGGCGGCCA AGGCGGCACC GCGGCACCG GCGGCAACGC CGGCGCCGGC GGCACCAGCT 120
TCACCCAAGG CGCCGACGGC AACGCCGGCA ACGGCGGTGA CGGCGGGGTC GCGGGAACG 180
GCGGAAACGG CGGAAACGGC GCAGACAACA CCACCACCGC CGCCGCC 227

(2) INFORMATION FOR SEQ ID NO:161:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 304 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:161:

CCTCGCCACC ATGGGCGGGC AGGGCGGTAG CGGTGGCGCC GGCTCTACCC CAGGCGCCAA 60
GGGCGCCCAC GGCTTCACTC CAACCAGCGG CGGCGACGGC GCGACGGCG GCAACGGCGG 120
CAACTCCCAA GTGGTCGGCG GCAACGGCGG CGACGGCGGC AATGGCGGCA ACGGCGGCAG 180
CGCCGGCACG GCGGCAACG GCGGCCGCGG CGGCGACGGC GCGTTTGGTG GCATGAGTGC 240
CAACGCCACC AACCTGGTG AAAACGGGCC AAACGGTAAC CCCGGCGGCA ACGGTGGCGC 300

CGGC

304

(2) INFORMATION FOR SEQ ID NO:162:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 1439 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:162:

GTGGGACGCT GCCGAGGCTG TATAACAAGG ACAACATCGA CCAGCGCCGG CTCGGTGAGC	60
TGATCGACCT ATTTAACAGT GCGCGCTTCA GCCGGCAGGG CGAGCACCGC GCCCGGGATC	120
TGATGGGTGA GGTCTACGAA TACTTCCTCG GCAATTTTCG TCGCGCGGAA GGGAAGCGGG	180
GTGGCGAGTT CTTTACCCCG CCCAGCGTGG TCAAGGTGAT CGTGGAGGTG CTGGAGCCGT	240
CGAGTGGGCG GGTGTATGAC CCGTGCTGCG GTTCCGGAGG CATGTTTGTG CAGACCGAGA	300
AGTTCATCTA CGAACACGAC GGCGATCCGA AGGATGTCTC GATCTATGGC CAGGAAAGCA	360
TTGAGGAGAC CTGGCGGATG GCGAAGATGA ACCTCGCCAT CCACGGCATC GACAACAAGG	420
GGCTCGGCGC CCGATGGAGT GATACCTTCG CCCGCGACCA GCACCCGGAC GTGCAGATGG	480
ACTACGTGAT GGCCAATCCG CCGTTCAACA TCAAAGACTG GGCCCGCAAC GAGGAAGACC	540
CACGCTGGCG CTTGCGGTGTT CCGCCCGCCA ATAACGCCAA CTACGCATGG ATTCAGCACA	600
TCCTGTACAA CTTGGCGCCG GGAGGTCGGG CGGGCGTGGT GATGGCCAAC GGGTCGATGT	660
CGTCGAACTC CAACGGCAAG GGGGATATTC GCGCGCAAAT CGTGGAGGCG GATTTGGTTT	720
CCTGCATGGT CGCGTTACCC ACCCAGCTGT TCCGCAGCAC CGGAATCCCG GTGTGCCTGT	780
GGTTTTTTCG CAAAACAAG GCGGCAGGTA AGCAAGGGTC TATCAACCGG TCGGGGCAGG	840
TGCTGTTTCAT CGACGCTCGT GAACTGGGCG ACCTAGTGGA CCGGGCCGAG CGGGCGCTGA	900
CCAACGAGGA GATCGTCCGC ATCGGGGATA CCTTCCACGC GAGCACGACC ACCGGAACG	960
CCGGCTCCGG TGGTGCCGGC GGTAATGGGG GCACTGGCCT CAACGGCGCG GGCGGTGCTG	1020
GCGGGGCCGG CGGCAACGCG GGTGTCGCG GCGTGTCTT CGGCAACGCT GTGGGCGGCG	1080
ACGGCGGCAA CGGCGGCAAC GGCGGCCACG GCGGCGACGG CACGACGGGC GGCGCCGGCG	1140
GCAAGGGCGG CAACGGCAGC AGCGGTGCCG CCAGCGGCTC AGGCGTCGTC AACGTCACCG	1200

CCGGCCACGG CGGCAACGGC GGCAATGGCG GCAACGGCGG CAACGGCTCC GCGGGCGCCG	1260
GCGGCCAGGG CGGTGCCGGC GGCAGCGCCG GCAACGGCGG CCACGGCGGC GGTGCCACCG	1320
GCGGCGCCAG CGGCAAGGGC GGCAACGGCA CCAGCGGTGC CGCCAGCGGC TCAGGCGTCA	1380
TCAACGTCAC CGCCGGCCAC GCGGCAACG GCGGCAATGG CCGCAACGGC GGCAACGGC	1439

(2) INFORMATION FOR SEQ ID NO:163:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 329 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:163:

GGGCCGGCGG GGCCGATTT TCTCGTGCCT TGATTGTCGC TGGGGATAAC GGCGGTGATG	60
GTGGTAACGG CGGGATGGGC GGGGCTGGCG GGGCTGGCGG CCCC GGCGGG GCCGGCGGCC	120
TGATCAGCCT GCTGGGCGGC CAAGGCGCCG GCGGGGCGCG CCGGACCGGC GGGGCCGGCG	180
GTGTTGGCGG TGACGGCGGG GCCGGCGGCC CCGGCAACCA GGCCTTCAAC GCAGGTGCCG	240
GCGGGGCGCG CGGCCTGATC AGCCTGCTGG GCGGCCAAGG CGCCGGCGGG GCCGGCGGGA	300
CCGGCGGGGC CGGCGGTGTT GCGGGTGAC	329

(2) INFORMATION FOR SEQ ID NO:164:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 80 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:164:

GCAACGGTGG CAACGGCGGC ACCAGCACGA CCGTGGGGAT GGCCGGAGGT AACTGTGGTG	60
CCGCCGGGCT GATCGGCAAC	80

(2) INFORMATION FOR SEQ ID NO:165:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 392 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:165:

```

GGGCTGTGTC GCACTCACAC CGCCGCATTC GGCGACGTTG GCCGCCCAAT ATCCAGCTCA      60
AGGCCTACTA CTTACCGTCG GAGGACCGCC GCATCAAGGT GCGGGTCAGC GCCCAAGGAA      120
TCAAGGTCAT CGACCGCGAC GGGCATCGAG GCCGTCGTCG CGCGGCTCGG GCAGGATCCG      180
CCCCGGCGCA CTTCGCGCGC CAAGCGGGCT CATCGCTCCG AACGGCGGCG ATCCTGTGAG      240
CACAAGTGAT GGC GCGCAAC GAGATTCGTC CAATTGTCAA GCCGTGTTTC ACCGCAGGGA      300
CCGGTTATAC GTATGTCAAC CTATGTCACT CGCAAGAACC GGCATAACGA TCCCGTGATC      360
CGCCGACAGC CCACGAGTGC AAGACCGTTA CA      392

```

(2) INFORMATION FOR SEQ ID NO:166:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 535 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:166:

```

ACCGGCGCCA CCGGCGGCAC CGGGTTCGCC GGTGGCGCCG GCGGGGCCGG CGGGCAGGGC      60
GGTATCAGCG GTGCCGGCGG CACCAACGGC TCTGGTGGCG CTGGCGGCAC CGGCGGACAA      120
GGCGGCGCCG GGGGCGCTGG CGGGGCCGGC GCCGATAACC CCACCGGCAT CGGCGGCGCC      180
GGCGGCACCG GCGGCACCGG CGGAGCGGCC GGAGCCGGCG GGGCCGGTGG CGCCATCGGT      240
ACCGGCGGCA CCGGCGGCGC GGTGGGCAGC GTCGGTAACG CCGGGATCGG CGGTACCGGC      300
GGTACGGGTG GTGTCGGTGG TGCTGGTGGT GCAGGTGCGG CTGCGGCCGC TGGCAGCAGC      360
GCTACCGGTG GCGCCGGGTT CGCCGGCGGC GCCGGCGGAG AAGGCGGACC GGGCGGCAAC      420
AGCGGTGTGG GCGGCACCAA CGGCTCCGGC GGCGCCGGCG GTGCAGGCGG CAAGGGCGGC      480

```

ACCGGAGGTG CCGGCGGGTC CGGCGCGGAC AACCCACCG GTGCTGGTTT CGCCG

535

(2) INFORMATION FOR SEQ ID NO:167:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 690 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:167:

CCGACGTCGC CGGGGCGATA CGGGGGTCAC CGACTACTAC ATCATCCGCA CCGAGAATCG	60
GCCGCTGCTG CAACCGCTGC GGGCGGTGCC GGTTCATCGGA GATCCGCTGG CCGACCTGAT	120
CCAGCCGAAC CTGAAGGTGA TCGTCAACCT GGGCTACGGC GACCCGAAC AC GGCTACTC	180
GACGAGCTAC GCCGATGTGC GAACGCCGTT CGGGCTGTGG CCGAACGTGC CGCCTCAGGT	240
CATCGCCGAT GCCCTGGCCG CCGGAACACA AGAAGGCATC CTTGACTTCA CGGCCGACCT	300
GCAGGCGCTG TCCGCGCAAC CGCTCACGCT CCCGCAGATC CAGCTGCCGC AACCCGCCGA	360
TCTGGTGGCC GCGGTGGCCG CCGCACCGAC GCCGGCCGAG GTGGTGAACA CGCTCGCCAG	420
GATCATCTCA ACCAACTACG CGCTCCTGCT GCCACCGTG GACATCGCCC TCGCCTGGTC	480
ACCACCCTGC CGCTGTACAC CACCCAACTG TTCGTCAGGC AACTCGCTGC GGGCAATCTG	540
ATCAACGCGA TCGGCTATCC CCTGGCGGCC ACCGTAGGTT TAGGCACGAT CGATAGCGGG	600
CGGCGTGGAA TTGCTACCC TCCTCGCGGC GGCCTCGGAC ACCGTTTCGAA ACATCGAGGG	660
CCTCGTCACC TAACGGATTC CCGACGGCAT	690

(2) INFORMATION FOR SEQ ID NO:168:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 407 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:168:

ACGGTGACGG CGGTACTGGC GCGGCCACG GCGGCAACGG CCGGAATCCC GGGTGGCTCT	60
TGGGCACAGC CGGGGGTGGC GGCAACGGTG GCGCCGGCAG CACCGGTACT GCAGGTGGCG	120
GCTCTGGGGG CACCGGCGGC GACGGCGGGA CCGGCGGGCG TGGCGGCCTG TTAATGGGCG	180
CCGGCGCCGG CGGGCACGGT GGCCTGGCG GCGCGGGCGG TGCCGGTGTC GACGGTGGCG	240
GCGCCGGCGG GGCCGGCGGG GCCGGCGGCA ACGGCGGCGC CGGGGGTCAA GCCGCCCTGC	300
TGTTCTGGGCG CGGCGGCACC GGCGGAGCCG GCGGCTACGG CGGCGATGGC GGTGGCGGCG	360
GTGACGGCTT CGACGGCACG ATGGCCGGCC TGGGTGGTAC CGGTGGC	407

(2) INFORMATION FOR SEQ ID NO:169:

- (i) SEQUENCE CHARACTERISTICS:
- (A) LENGTH: 468 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:169:

GATCGGTGTCAG CGCATCGCCC TCGGCGGCAA GCGATTCCGC GGTCTCACCG AAGAACATCG	60
TGCACGCGGC GGCGCGGACC AGCCCGCTGC GCTGCGGCGC GTCGAACGCC TCCAGCAGGC	120
ACAGCCAGTC CTTGGCGGCC TGCAGGCGA ACACGTCGGT GTCACCGGTG TAGATCGCCG	180
GGATGCCCCG CTCCGCCAAC GCATTCCGGC ACGCCCGCGC GTCTTTGTGA TGCTCGACGA	240
TCACCGCGAT GTCTGCGGCC ACCACGGGCC GCCCGGCGAA GGTGGCCCCG CTGGCCAGTA	300
GCGCCGCGAC GTCGGCGGCC AGGTCGTCGG GGATGTGCCG GCGCAGCGCT CCGGCGCGAC	360
GCCCGAAAAA CGACCCCTCA CCCAGCTGGG TCCCGCTGGC ATATCCCTTG CCGTCCTGGG	420
CGATATTGGA CGCGCATGCC CCGACCGCGT ACAGGCCGGC CACCACCG	468

(2) INFORMATION FOR SEQ ID NO:170:

- (i) SEQUENCE CHARACTERISTICS:
- (A) LENGTH: 219 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:170:

GGTGGTAACG GCGGCCAGGG TGGCATCGGC GGCGCCGGCG AGAGAGGCGC CGACGGCGCC	60
GGCCCCAATG CTAACGGCGC AACGGCGAG AACGGCGGTA GCGGTGGTAA CGGTGGCGAC	120
GGCGGCGCCG GCGGCAATGG CGGCGCGGGC GGCAACGCGC AGGCGGCCGG GTACACCGAC	180
GGCGCCACGG GCACCGGCGG CGACGGCGGC AACGGCGGC	219

(2) INFORMATION FOR SEQ ID NO:171:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 494 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:171:

TAGCTCCGGC GAGGGCGGCA AGGGCGGCGA CGGTGGCCAC GGCGGTGACG GCGTCGGCGG	60
CAACAGTTCC GTCACCCAAG GCGGCAGCGG CGGTGGCGGC GGCGCCGGCG GCGCCGGCGG	120
CAGCGGCTTT TTCGGCGGCA AGGGCGGCTT CGGCGGCGAC GGCGGTCAGG GCGGCCCCAA	180
CGGCGGCGGT ACCGTCGGCA CCGTGGCCGG TGGCGGCGGC AACGGCGGTG TCGGCGGCCG	240
GGGCGGCGAC GGCGTCTTTG CCGGTGCCGG CGGCCAGGGC GGCCTCGGTG GGCAGGGCGG	300
CAATGGCGGC GGCTCCACCG GCGGCAACGG CGGCCTTGGC GGCGCGGGCG GTGGCGGAGG	360
CAACGCCCCG GCTCGTGCCG AATCCGGGCT GACCATGGAC AGCGCGGCCA AGTTCGCTGC	420
CATCGCATCA GGCGCGTACT GCCCCGAACA CCTGGAACAT CACCCGAGTT AGCGGGGCGC	480
ATTTCCTGAT CACC	494

(2) INFORMATION FOR SEQ ID NO:172:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 220 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:172:

GGGCCGGTGG TGCCGCGGGC CAGCTCTTCA GCGCCGGAGG CGCGGCGGGT GCCGTTGGGG	60
TTGGCGGCAC CGGCGGCCAG GGTGGGGCTG GCGGTGCCGG AGCGGCCGGC GCCGACGCCC	120
CCGCCAGCAC AGGTCTAACC GGTGGTACCG GGTTCGCTGG CGGGGCCGGC GGCGTCGGCG	180
GCCAGAGCGG CAACGCCATT GCCGGCGGCA TCAACGGCTC	220

(2) INFORMATION FOR SEQ ID NO:173:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 388 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:173:

ATGGCGGCAA CGGGGGCCCC GGCGGTGCTG GCGGGGCCGG CGACTACAAT TTCCAACGGC	60
GGGCAGGGTG GTGCCGGCGG CCAAGGCGGC CAAGGCGGCC TGGGCGGGGC AAGCACCACC	120
TGATCGGCCT AGCCGCACCC GGGAAAGCCG ATCCAACAGG CGACGATGCC GCCTTCCTTG	180
CCGCGTTGGA CCAGGCCGGC ATCACCTACG CTGACCCAGG CCACGCCATA ACGGCCGCCA	240
AGGCGATGTG TGGGCTGTGT GCTAACGGCG TAACAGGTCT ACAGCTGGTC GCGGACCTGC	300
GGGACTACAA TCCCGGGCTG ACCATGGACA GCGCGGCCAA GTTCGCTGCC ATCGCATCAG	360
GCGCGTACTG CCCCGAACAC CTGGAACA	388

(2) INFORMATION FOR SEQ ID NO:174:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 400 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:174:

GCAAAGGCGG CACCGGCGGG GCCGGCATGA ACAGCCTCGA CCCGCTGCTA GCCGCCCAAG	60
ACGGCGGCCA AGGCGGCACC GGCGGCACCG GCGGCAACGC CGGCGCCGGC GGCACCAGCT	120

TCACCCAAGG CGCCGACGGC AACGCCGGCA ACGGCGGTGA CGGCGGGGTC GGCGGCAACG	180
GCGGAAACGG CGGAAACGGC GCAGACAACA CCACCACCGC CGCCGCCGGC ACCACAGGCG	240
GCGACGGCGG GGCCGGCGGG GCCGGCGGAA CCGGCGGAAC CGGCGGAGCC GCCGGCACCG	300
GCACCGGCGG CCAACAAGGC AACGGCGGCA ACGGCGGCAC CGGCGGCAAA GGCGGCACCG	360
GCGGCGACGG TGCACTCTCA GGCAGCACCG GTGGTGCCGG	400

(2) INFORMATION FOR SEQ ID NO:175:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 538 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:175:

GGCAACGGCG GCAACGGCGG CATCGCCGGC ATTGGGCGGC AACGGCGTTC CGGGACGGGC	60
AGCGGCAACG GCGGCCAACG GCGGCAGCGG CGGCAACGGC GGCAACGCCG GCATGGGCGG	120
CAACAGCGGC ACCGGCAGCG GCGACGGCGG TGCCGGCGGG AACGGCGGCG CGGCGGGCAC	180
GGGCGGCACC GGCGGCGACG GCGGCCTCAC CGGTACTGGC GGCACCGGCG GCAGCGGTGG	240
CACCGGCGGT GACGGCGGTA ACGGCGGCAA CGGAGCAGAT AACACCGCAA ACATGACTGC	300
GCAGGCGGGC GGTGACGGTG GCAACGGCGG CGACGGTGGC TTCGGCGGCG GGGCCGGGGC	360
CGGCGGCGGT GGCTTGACCG CTGGCGCCAA CGGCACCGGC GGGCAAGGCG GCGCCGGCGG	420
CGATGGCGGC AACGGGGCCA TCGGCGGCCA CGGCCCACTC ACTGACGACC CCGGCGGCAA	480
CGGGGGCACC GGCGGCAACG GCGGCACCGG CGGCACCGGC GGCGCGGGCA TCGGCAGC	538

(2) INFORMATION FOR SEQ ID NO:176:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 239 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:176:

GGGCCGGTGG TGCCGCGGGC CAGCTCTTCA GCGCCGAGG CGCGGCGGGT GCCGTTGGGG	60
TTGGCGGCAC CGGCGGCCAG GGTGGGGCTG GCGGTGCCGG AGCGGCCGGC GCCGACGCCC	120
CCGCCAGCAC AGGTCTAACC GGTGGTACCG GGTTCGCTGG CGGGGCCGGC GGCGTCGGCG	180
GCCACGGCGG CAACGCCATT GCCGGCGGCA TCAACGGCTC CGGTGGTGCC GGCGGCACC	239

(2) INFORMATION FOR SEQ ID NO:177:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 985 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:177:

AGCAGCGCTA CCGGTGGCGC CGGGTTCGCC GGCGGCGCCG GCGGAGAAGG CGGAGCGGGC	60
GGCAACAGCG GTGTGGGCGG CACCAACGGC TCCGGCGGCG CCGGCGGTGC AGGCGGCAAG	120
GGCGGCACCG GAGGTGCCGG CGGGTCCGGC GCGGACAACC CCACCGGTGC TGGTTTCGCC	180
GGTGGCGCCG GCGGCACAGG TGGCGCGGCC GGCGCCGGCG GGGCCGGCGG GGCGACCGGT	240
ACCGGCGGCA CCGGCGGCGT TGTCGGCGCC ACCGGTAGTG CAGGCATCGG CGGGGCCGGC	300
GGCCGCGGCG GTGACGGCGG CGATGGGGCC AGCGGTCTCG GCCTGGGCCT CTCCGGCTTT	360
GACGGCGGCC AAGGCGGCCA AGGCGGGGCC GGCGGCAGCG CCGGCGCCGG CGGCATCAAC	420
GGGGCCGGCG GGGCCGGCGG CAACGGCGGC GACGGCGGGG ACGGCGCAAC CGGTGCCGCA	480
GGTCTCGGCG ACAACGGCGG GGTGCGCGGT GACGGTGGGG CCGGTGGCGC CGCCGGCAAC	540
GGCGGCAACG CGGGCGTCGG CCTGACAGCC AAGGCCGGCG ACGGCGGCGC CGCGGGCAAT	600
GGCGGCAACG GGGGCGCCGG CGGTGCTGGC GGGGCGGCG ACAACAATTT CAACGGCGGC	660
CAGGGTGGTG CCGGCGGCCA AGGCGGCCAA GGCGGCTTGG GCGGGCAAG CACCACCTGA	720
TCGGCCTAGC CGCACCCTGG AAAGCCGATC CAACAGGCGA CGATGCCGCC TTCCTTGCCG	780
CGTTGGACCA GGCCGGCATC ACCTACGCTG ACCCAGGCCA CGCCATAACG GCCGCCAAGG	840
CGATGTGTGG GCTGTGTGCT AACGGCGTAA CAGGTCTACA GCTGGTCGCG GACCTGCGGG	900
AATACAATCC CGGGCTGACC ATGGACAGCG CGGCCAAGTT CGCTGCCATC GCATCAGGCG	960

CGTACTGCCC CGAACACCTG GAACA

985

(2) INFORMATION FOR SEQ ID NO:178:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 2138 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:178:

CGGCACGAGG ATCGGTACCC CGCGGCATCG GCAGCTGCCG ATTCGCCGGG TTTCCCCACC	60
CGAGGAAAGC CGCTACCAGA TGGCGCTGCC GAAGTAGGGC GATCCGTTTC CGATGCCGGC	120
ATGAACGGGC GGCATCAAAT TAGTGCAGGA ACCTTTCAGT TTAGCGACGA TAATGGCTAT	180
AGCACTAAGG AGGATGATCC GATATGACGC AGTCGCAGAC CGTGACGGTG GATCAGCAAG	240
AGATTTTGAA CAGGGCCAAC GAGGTGGAGG CCCCATGGC GGACCCACCG ACTGATGTCC	300
CCATCACACC GTGCGAACTC ACGGCGGCTA AAAACGCCGC CCAACAGCTG GTATTGTCCG	360
CCGACAACAT GCGGGAATAC CTGGCGGCCG GTGCCAAAGA GCGGCAGCGT CTGGCGACCT	420
CGCTGCGCAA CGCGCCAAG GCGTATGGCG AGGTTGATGA GGAGGCTGCG ACCGCGCTGG	480
ACAACGACGG CGAAGGAACT GTGCAGGCAG AATCGGCCGG GGCCGTCGGA GGGGACAGTT	540
CGGCCGAACT AACCGATACG CCGAGGGTGG CCACGGCCGG TGAACCCAAC TTCATGGATC	600
TCAAAGAAGC GGCAAGGAAG CTCGAAACGG GCGACCAAGG CGCATCGCTC GCGCACTTTG	660
CGGATGGGTG GAACACTTTC AACCTGACGC TGCAAGGCGA CGTCAAGCGG TTCCGGGGGT	720
TTGACAACTG GGAAGGCGAT GCGGCTACCG CTTGCGAGGC TTCGCTCGAT CAACAACGGC	780
AATGGATACT CCACATGGCC AAATTGAGCG CTGCGATGGC CAAGCAGGCT CAATATGTCG	840
CGCAGCTGCA CGTGTGGGCT AGGCGGGAAC ATCCGACTTA TGAAGACATA GTCGGGCTCG	900
AACGGCTTTA CGCGGAAAAC CCTTCGGCCC GCGACCAAAT TCTCCCGGTG TACGCGGAGT	960
ATCAGCAGAG GTCGGAGAAG GTGCTGACCG AATACAACAA CAAGGCAGCC CTGGAACCGG	1020
TAAACCCGCC GAAGCCTCCC CCCGCCATCA AGATCGACCC GCCCCCGCCT CCGCAAGAGC	1080
AGGGATTGAT CCCTGGCTTC CTGATGCCGC CGTCTGACGG CTCCGGTGTG ACTCCCGGTA	1140

```

CCGGGATGCC AGCCGCACCG ATGGTTCCGC CTACCGGATC GCCGGGTGGT GGCCTCCCGG      1200
CTGACACGGC GGCGCAGCTG ACGTCGGCTG GGCGGGAAGC CGCAGCGCTG TCGGGCGACG      1260
TGGCGGTCAA AGCGGCATCG CTCGGTGGCG GTGGAGGCGG CGGGGTGCCG TCGGCGCCGT      1320
TGGGATCCGC GATCGGGGGC GCCGAATCGG TCGGGCCCGC TGGCGCTGGT GACATTGCCG      1380
GCTTAGGCCA GGGAAGGGCC GGCGGCGGCG CCGCGCTGGG CGGCGGTGGC ATGGGAATGC      1440
CGATGGGTGC CGCGCATCAG GGACAAGGGG GCGCCAAGTC CAAGGGTTCT CAGCAGGAAG      1500
ACGAGGCGCT CTACACCGAG GATCGGGCAT GGACCGAGGC CGTCATTGGT AACCGTCGGC      1560
GCCAGGACAG TAAGGAGTCG AAGTGAGCAT GGACGAATTG GACCCGCATG TCGCCCGGGC      1620
GTTGACGCTG GCGGCGCGGT TTCAGTCGGC CCTAGACGGG ACGCTCAATC AGATGAACAA      1680
CGGATCCTTC CGCGCCACCG ACGAAGCCGA GACCGTCGAA GTGACGATCA ATGGGCACCA      1740
GTGGCTCACC GGCTTGCACA TCGAAGATGG TTTGCTGAAG AAGCTGGGTG CCGAGGCGGT      1800
GGCTCAGCGG GTCAACGAGG CGCTGCACAA TGCGCAGGCC GCGGCGTCCG CGTATAACGA      1860
CGCGGCGGGC GAGCAGCTGA CCGCTGCGTT ATCGGCCATG TCCCGCGCGA TGAACGAAGG      1920
AATGGCCTAA GCCCATTGTT GCGGTGGTAG CGACTACGCA CCGAATGAGC GCCGCAATGC      1980
GGTCATTAG CGCGCCCGAC ACGGCGTGAG TACGCATTGT CAATGTTTTG ACATGGATCG      2040
GCCGGGTTCG GAGGGCGCCA TAGTCCTGGT CGCCAATATT GCCGCAGCTA GCTGGTCTTA      2100
GGTTCGGTTA CGCTGGTTAA TTATGACGTC CGTTACCA                                2138

```

(2) INFORMATION FOR SEQ ID NO:179:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 460 amino acids
- (B) TYPE: amino acid
- (C) STRANDEDNESS:
- (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:179:

```

Met Thr Gln Ser Gln Thr Val Thr Val Asp Gln Gln Glu Ile Leu Asn
1           5           10           15

Arg Ala Asn Glu Val Glu Ala Pro Met Ala Asp Pro Pro Thr Asp Val
          20           25           30

Pro Ile Thr Pro Cys Glu Leu Thr Ala Ala Lys Asn Ala Ala Gln Gln

```

190

35	40	45
Leu Val Leu Ser Ala Asp Asn Met Arg Glu Tyr Leu Ala Ala Gly Ala		
50	55	60
Lys Glu Arg Gln Arg Leu Ala Thr Ser Leu Arg Asn Ala Ala Lys Ala		
65	70	75
		80
Tyr Gly Glu Val Asp Glu Glu Ala Ala Thr Ala Leu Asp Asn Asp Gly		
	85	90
		95
Glu Gly Thr Val Gln Ala Glu Ser Ala Gly Ala Val Gly Gly Asp Ser		
	100	105
		110
Ser Ala Glu Leu Thr Asp Thr Pro Arg Val Ala Thr Ala Gly Glu Pro		
	115	120
		125
Asn Phe Met Asp Leu Lys Glu Ala Ala Arg Lys Leu Glu Thr Gly Asp		
	130	135
		140
Gln Gly Ala Ser Leu Ala His Phe Ala Asp Gly Trp Asn Thr Phe Asn		
	145	150
		155
		160
Leu Thr Leu Gln Gly Asp Val Lys Arg Phe Arg Gly Phe Asp Asn Trp		
	165	170
		175
Glu Gly Asp Ala Ala Thr Ala Cys Glu Ala Ser Leu Asp Gln Gln Arg		
	180	185
		190
Gln Trp Ile Leu His Met Ala Lys Leu Ser Ala Ala Met Ala Lys Gln		
	195	200
		205
Ala Gln Tyr Val Ala Gln Leu His Val Trp Ala Arg Arg Glu His Pro		
	210	215
		220
Thr Tyr Glu Asp Ile Val Gly Leu Glu Arg Leu Tyr Ala Glu Asn Pro		
	225	230
		235
		240
Ser Ala Arg Asp Gln Ile Leu Pro Val Tyr Ala Glu Tyr Gln Gln Arg		
	245	250
		255
Ser Glu Lys Val Leu Thr Glu Tyr Asn Asn Lys Ala Ala Leu Glu Pro		
	260	265
		270
Val Asn Pro Pro Lys Pro Pro Pro Ala Ile Lys Ile Asp Pro Pro Pro		
	275	280
		285
Pro Pro Gln Glu Gln Gly Leu Ile Pro Gly Phe Leu Met Pro Pro Ser		
	290	295
		300
Asp Gly Ser Gly Val Thr Pro Gly Thr Gly Met Pro Ala Ala Pro Met		
	305	310
		315
		320
Val Pro Pro Thr Gly Ser Pro Gly Gly Gly Leu Pro Ala Asp Thr Ala		
	325	330
		335

Ala Gln Leu Thr Ser Ala Gly Arg Glu Ala Ala Ala Leu Ser Gly Asp
 340 345 350

Val Ala Val Lys Ala Ala Ser Leu Gly Gly Gly Gly Gly Gly Gly Val
 355 360 365

Pro Ser Ala Pro Leu Gly Ser Ala Ile Gly Gly Ala Glu Ser Val Arg
 370 375 380

Pro Ala Gly Ala Gly Asp Ile Ala Gly Leu Gly Gln Gly Arg Ala Gly
 385 390 395 400

Gly Gly Ala Ala Leu Gly Gly Gly Gly Met Gly Met Pro Met Gly Ala
 405 410 415

Ala His Gln Gly Gln Gly Gly Ala Lys Ser Lys Gly Ser Gln Gln Glu
 420 425 430

Asp Glu Ala Leu Tyr Thr Glu Asp Arg Ala Trp Thr Glu Ala Val Ile
 435 440 445

Gly Asn Arg Arg Arg Gln Asp Ser Lys Glu Ser Lys
 450 455 460

(2) INFORMATION FOR SEQ ID NO:180:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 277 amino acids
- (B) TYPE: amino acid
- (C) STRANDEDNESS:
- (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:180:

Ala Gly Asn Val Thr Ser Ala Ser Gly Pro His Arg Phe Gly Ala Pro
 1 5 10 15

Asp Arg Gly Ser Gln Arg Arg Arg Arg His Pro Ala Ala Ser Thr Ala
 20 25 30

Thr Glu Arg Cys Arg Phe Asp Arg His Val Ala Arg Gln Arg Cys Gly
 35 40 45

Phe Pro Pro Ser Arg Arg Gln Leu Arg Arg Arg Val Ser Arg Glu Ala
 50 55 60

Thr Thr Arg Arg Ser Gly Arg Arg Asn His Arg Cys Gly Trp His Pro
 65 70 75 80

Gly Thr Gly Ser His Thr Gly Ala Val Arg Arg Arg His Gln Glu Ala

85

90

95

Arg	Asp	Gln	Ser	Leu	Leu	Leu	Arg	Arg	Arg	Gly	Arg	Val	Asp	Leu	Asp
			100				105						110		
Gly	Gly	Gly	Arg	Leu	Arg	Arg	Val	Tyr	Arg	Phe	Gln	Gly	Cys	Leu	Val
			115				120						125		
Val	Val	Phe	Gly	Gln	His	Leu	Leu	Arg	Pro	Leu	Leu	Ile	Leu	Arg	Val
			130				135						140		
His	Arg	Glu	Asn	Leu	Val	Ala	Gly	Arg	Arg	Val	Phe	Arg	Val	Lys	Pro
			145				150						155		
Phe	Glu	Pro	Asp	Tyr	Val	Phe	Ile	Ser	Arg	Met	Phe	Pro	Pro	Ser	Pro
			165				170						175		
His	Val	Gln	Leu	Arg	Asp	Ile	Leu	Ser	Leu	Leu	Gly	His	Arg	Ser	Ala
			180				185						190		
Gln	Phe	Gly	His	Val	Glu	Tyr	Pro	Leu	Pro	Leu	Leu	Ile	Glu	Arg	Ser
			195				200						205		
Leu	Ala	Ser	Gly	Ser	Arg	Ile	Ala	Phe	Pro	Val	Val	Lys	Pro	Pro	Glu
			210				215						220		
Pro	Leu	Asp	Val	Ala	Leu	Gln	Arg	Gln	Val	Glu	Ser	Val	Pro	Pro	Ile
			225				230						235		
Arg	Lys	Val	Arg	Glu	Arg	Cys	Ala	Leu	Val	Ala	Arg	Phe	Glu	Leu	Pro
			245				250						255		
Cys	Arg	Phe	Phe	Glu	Ile	His	Glu	Val	Gly	Phe	Thr	Gly	Arg	Gly	His
			260				265						270		
Pro	Arg	Arg	Ile	Gly											
			275												

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 192 amino acids
(B) TYPE: amino acid
(C) STRANDEDNESS:
(D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:181:

Arg Val Ala Ala Ser Phe Ile Asp Trp Leu Asp Ser Pro Asp Ser Pro
1 5 10 15

193

Leu Asp Pro Ser Leu Val Ser Ser Leu Leu Asn Ala Val Ser Cys Gly
 20 25 30
 Ala Glu Ser Ser Ala Ser Ser Ser Ala Arg Ser Gly Asn Gly Ser Arg
 35 40 45
 Trp Thr Ser Met Pro Ser Gly Thr Arg Pro Gly Pro Arg Arg Ala Thr
 50 55 60
 Ser Arg Asp Asp Arg Arg Ser Ala Thr Ser Val Ile Pro Ser Arg Arg
 65 70 75 80
 Ser Val Ala Pro Arg Ala Glu Phe Gly Thr Arg Leu Ala Ser His Arg
 85 90 95
 Ala Ser Pro Ser Asn Ala Cys Pro Val Arg Ile Val Thr Ser Ala Ser
 100 105 110
 Gly Arg Pro Ile Ser Ser Pro Pro Ile Val Arg Ser Arg Ser Cys Val
 115 120 125
 Asp Lys Asn Gly Arg Arg Cys Ala Ser Gly Tyr Arg Arg Leu Asn Arg
 130 135 140
 Ala Arg Ser Ser Ser Ile Ala Ala Arg Cys Arg Thr Ile Gly Thr Phe
 145 150 155 160
 Arg Arg Ser Arg Tyr Ser Ala Ser Met Arg Val Ser Thr Asn Ser Pro
 165 170 175
 His Val Thr His Gly Val Ala Pro Gly Val Thr Arg Arg Ile Gly Gly
 180 185 190

(2) INFORMATION FOR SEQ ID NO:182:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 196 amino acids
- (B) TYPE: amino acid
- (C) STRANDEDNESS:
- (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:182:

Gln Glu Arg Pro Gln Met Cys Gln Arg Val Ser Glu Ile Glu Pro Arg
 1 5 10 15
 Thr Gln Phe Phe Asn Arg Cys Ala Leu Pro His Tyr Trp His Phe Pro
 20 25 30
 Ala Val Ala Val Phe Ser Lys His Ala Ser Leu Asp Glu Leu Ala Pro

194

35	40	45
Arg Asn Pro Arg Arg Ser Ser Arg Arg Asp Ala Glu Asp Arg Arg Val		
50	55	60
Ile Phe Ala Ala Thr Leu Val Ala Val Asp Pro Pro Leu Arg Gly Ala		
65	70	75 80
Gly Gly Glu Ala Asp Gln Leu Ile Asp Leu Gly Val Cys Arg Arg Gln		
	85	90 95
Ala Gly Arg Val Arg Arg Gly Gln Glu Leu His His Arg His Arg His		
	100	105 110
Gln Gly Ala Ala Pro Asp Leu Arg Arg Arg Arg Arg His Arg Arg Val		
	115	120 125
Gln Gln His Arg Arg Leu Gln Arg Val Arg Gln Leu Arg Arg Tyr Val		
	130	135 140
Gln Thr Ala His His Arg Arg Phe Ala Arg Thr Asp Arg Val Arg His		
	145	150 155 160
His Val Arg Gly Pro Ser Asn His Arg Arg Arg Arg Val Tyr Arg Gly		
	165	170 175
Arg His Ser Gly Ala Gly Gly Cys Pro Ala Gly Gly Ala Gly Ser Val		
	180	185 190
Gly Gly Ser Ala		
195		

(2) INFORMATION FOR SEQ ID NO:183:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 311 amino acids
- (B) TYPE: amino acid
- (C) STRANDEDNESS:
- (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:183:

Val Arg Cys Gly Thr Leu Val Pro Val Pro Met Val Glu Phe Leu Thr
1 5 10 15
Ser Thr Asn Ala Pro Ser Leu Pro Ser Ala Tyr Ala Glu Val Asp Lys
20 25 30
Leu Ile Gly Leu Pro Ala Gly Thr Ala Lys Arg Trp Ile Asn Gly Tyr
35 40 45

195

Glu Arg Gly Gly Lys Asp His Pro Pro Ile Leu Arg Val Thr Pro Gly
 50 55 60

Ala Thr Pro Trp Val Thr Trp Gly Glu Phe Val Glu Thr Arg Met Leu
 65 70 75 80

Ala Glu Tyr Arg Asp Arg Arg Lys Val Pro Ile Val Arg Gln Arg Ala
 85 90 95

Ala Ile Glu Glu Leu Arg Ala Arg Phe Asn Leu Arg Tyr Pro Leu Ala
 100 105 110

His Leu Arg Pro Phe Leu Ser Thr His Glu Arg Asp Leu Thr Met Gly
 115 120 125

Gly Glu Glu Ile Gly Leu Pro Asp Ala Glu Val Thr Ile Arg Thr Gly
 130 135 140

Gln Ala Leu Leu Gly Asp Ala Arg Trp Leu Ala Ser Leu Val Pro Asn
 145 150 155 160

Ser Ala Arg Gly Ala Thr Leu Arg Arg Leu Gly Ile Thr Asp Val Ala
 165 170 175

Asp Leu Arg Ser Ser Arg Glu Val Ala Arg Arg Gly Pro Gly Arg Val
 180 185 190

Pro Asp Gly Ile Asp Val His Leu Leu Pro Phe Pro Asp Leu Ala Asp
 195 200 205

Asp Asp Ala Asp Asp Ser Ala Pro His Glu Thr Ala Phe Lys Arg Leu
 210 215 220

Leu Thr Asn Asp Gly Ser Asn Gly Glu Ser Gly Glu Ser Ser Gln Ser
 225 230 235 240

Ile Asn Asp Ala Ala Thr Arg Tyr Met Thr Asp Glu Tyr Arg Gln Phe
 245 250 255

Pro Thr Arg Asn Gly Ala Gln Arg Ala Leu His Arg Val Val Thr Leu
 260 265 270

Leu Ala Ala Gly Arg Pro Val Leu Thr His Cys Phe Ala Gly Lys Asp
 275 280 285

Arg Thr Gly Phe Val Val Ala Leu Val Leu Glu Ala Val Gly Leu Asp
 290 295 300

Arg Asp Val Ile Val Ala Asp
 305 310

(2) INFORMATION FOR SEQ ID NO:184:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 2072 base pairs

(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:184:

CTCGTGCCGA TTCGGCACGA GCTGAGCAGC CCAAGGGGCC GTTCGGCGAA GTCATCGAGG	60
CATTCGCCGA CGGGCTGGCC GGCAAGGGTA AGCAAATCAA CACCACGCTG AACAGCCTGT	120
CGCAGGCGTT GAACGCCTTG AATGAGGGCC GCGGCGACTT CTTCGCGGTG GTACGCAGCC	180
TGGCGCTATT CGTCAACGCG CTACATCAGG ACGACCAACA GTTCGTCGCG TTGAACAAGA	240
ACCTTGCGGA GTTCACCGAC AGGTTGACCC ACTCCGATGC GGACCTGTCT AACGCCATCC	300
AGCAATTCGA CAGCTTGCTC GCCGTCGCGC GCCCGTTCTT CGCCAAGAAC CGCGAGGTGC	360
TGACGCATGA CGTCAATAAT CTCGCGACCG TGACCACCAC GTTGCTGCAG CCCGATCCGT	420
TGGATGGGTT GGAGACCGTC CTGCACATCT TCCCGACGCT GCGGCGGAAC ATTAACCAGC	480
TTTACCATCC GACACACGGT GCGTGGTGT CGCTTTCGCG GTTCACGAAT TTCGCCAACC	540
CGATGGAGTT CATCTGCAGC TCGATTGAGG CGGGTAGCCG GCTCGGTTAT CAAGAGTCGG	600
CCGAAGTCTG TGCGCAGTAT CTGGCGCCAG TCCTCGATGC GATCAAGTTC AACTACTTTC	660
CGTTCGGCCT GAACGTGGCC AGCACCGCCT CGACACTGCC TAAAGAGATC GCGTACTCCG	720
AGCCCCGCTT GCAGCCGCCC AACGGGTACA AGGACACCAC GGTGCCCCGGC ATCTGGGTGC	780
CGGATACGCC GTTGTCACAC CGCAACACGC AGCCCGGTTG GGTGGTGGCA CCCGGGATGC	840
AAGGGGTTCA GGTGGGACCG ATCACGCAGG GTTTGCTGAC GCCGGAGTCC CTGGCCGAAC	900
TCATGGGTGG TCCCGATATC GCCCCTCCGT CGTCAGGGCT GCAAACCCCG CCCGGACCCC	960
CGAATGCGTA CGACGAGTAC CCCGTGCTGC CGCCGATCGG TTTACAGGCC CCACAGGTGC	1020
CGATACCACC GCCGCCTCCT GGGCCCGACG TAATCCCGGG TCCGGTGCCA CCGGTCTTGG	1080
CGGCGATCGT GTTCCCAAGA GATCGCCCGG CAGCGTCGGA AACTTCGAC TACATGGGCC	1140
TCTTGTTGCT GTCGCCGGGC CTGGCGACCT TCCTGTTGCG GGTGTCATCT AGCCCCGCCC	1200
GTGGAACGAT GGCCGATCGG CACGTGTTGA TACCGGCGAT CACCGGCCTG GCGTTGATCG	1260
CGGCATTCGT CGCACATTCG TGGTACCGCA CAGAACATCC GCTCATAGAC ATGCGCTTGT	1320
TCCAGAACCG AGCGGTCGCG CAGGCCAACA TGACGATGAC GGTGCTCTCC CTCGGGCTGT	1380

TTGGCTCCTT	CTTGCTGCTC	CCGAGCTACC	TCCAGCAAGT	GTTGCACCAA	TCACCGATGC	1440
AATCGGGGGT	GCATATCATC	CCACAGGGCC	TCGGTGCCAT	GCTGGCGATG	CCGATCGCCG	1500
GAGCGATGAT	GGACCGACGG	GGACCGGCCA	AGATCGTGCT	GGTTGGGATC	ATGCTGATCG	1560
CTGCGGGGTT	GGGCACCTTC	GCCTTTGGTG	TCGCGCGGCA	AGCGGACTAC	TTACCCATTC	1620
TGCCGACCGG	GCTGGCAATC	ATGGGCATGG	GCATGGGCTG	CTCCATGATG	CCACTGTCCG	1680
GGGCGGCAGT	GCAGACCCTG	GCCCCACATC	AGATCGCTCG	CGGTTCGACG	CTGATCAGCG	1740
TCAACCAGCA	GGTGGGCGGT	TCGATAGGGA	CCGCACTGAT	GTCGGTGCTG	CTCACCTACC	1800
AGTTCAATCA	CAGCGAAATC	ATCGCTACTG	CAAAGAAAGT	CGCACTGACC	CCAGAGAGTG	1860
GCGCCGGGCG	GGGGGCGGCG	GTTGACCCTT	CCTCGCTACC	GCGCCAAACC	AACTTCGCGG	1920
CCCAACTGCT	GCATGACCTT	TCGCACGCCT	ACGCGGTGGT	ATTCTGTATA	GCGACCGCGC	1980
TAGTGGTCTC	GACGCTGATC	CCCGCGGCAT	TCCTGCCGAA	ACAGCAGGCT	AGTCATCGAA	2040
GAGCACCGTT	GCTATCCGCA	TGACGTCTGC	TT			2072

(2) INFORMATION FOR SEQ ID NO:185:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 1923 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:185:

TCACCCCGGA	GAAGTCGTTT	GTCGACGACC	TGGACATCGA	CTCGCTGTCTG	ATGGTCGAGA	60
TCGCCGTGCA	GACCGAGGAC	AAGTACGGCG	TCAAGATCCC	CGACGAGGAC	CTCGCCGGTC	120
TGCGTACCGT	CGGTGACGTT	GTCGCCTACA	TCCAGAAGCT	CGAGGAAGAA	AACCCGGAGG	180
CGGCTCAGGC	GTTGCGCGCG	AAGATTGAGT	CGGAGAACCC	CGATGCGGCA	CGAGCAGATC	240
GGTGCGTTTC	ACCCACATCG	CAAGCTCGAG	ACGCCCCTCG	TCCTCTTGCA	CGCTCAGCCA	300
GGTTGGCGTG	TCGCCGCCTT	CCAGCAAGTG	TTCCCACCAC	ACGAAGGGAC	CCTCGCGAAA	360
GGTGACTGAT	CCGCGGACCA	CATAGTCGAT	GCCACCGTGG	CTGACAATTG	CGCCGGGTCC	420
GAGTTGGCGG	GGGCCGAATT	GCGGCATTGC	GTCGAAGGCC	AGCGGATCCC	GGCGCCCGCC	480

CGGCGTGGCT	GGTGT TTTGG	GCCGCCGGAT	GGCCACGACG	AGAACGACGA	TGGCGGCGAT	540
GAACAGCGCC	ACGGCAATCA	CGACCAGCAG	ATTTCCCACG	CATACCCTCT	CGTACCGCTG	600
CGCCGCGGTT	GGTCGATCGG	TCGCATATCG	ATGGCGCCGT	TTAACGTAAC	AGCTTTTCGG	660
GGACCGGGGG	TCACAACGGG	CGAGTTGTCC	GGCCGGGAAC	CCGGCAGGTC	TCGGCCGCGG	720
TCACCCAGC	TCACTGGTGC	ACCATCCGGG	TGTCGGTGAG	CGTGCAACTC	AAACACACTC	780
AACGGCAACG	GTTTCTCAGG	TCACCAGCTC	AACCTCGACC	CGCAATCGCT	CGTACGTTTC	840
GACCGCGCGC	AGGTCGCGAG	TCAGCAGCTT	TGCGCCGGCA	GCTTTCGCCG	TGAAGCCGAC	900
CAGGGCATCG	TAGGTTGCGC	CACCGGTGAC	ATCGTGCTCG	GCGAGGTGGT	CGGTCAAGCC	960
GCGATATGAG	CAGGCATCCA	GTGCCAGGTA	GTTGCTGGAG	GTGATGTCCG	CCAAGTAGGC	1020
GTGGACGGCA	ACAGGGGCAA	TACGATGCGG	CGGTGGTAGC	CGGGTCAAGA	CCGAATAGGT	1080
TTCCACAGCC	GCGTGCGCGA	TCAGATGGAC	GCCACGGTTG	AGCGCGCGCA	CGGCGGCCTC	1140
GTGCCCTTCG	TGCCAGGTCG	CGAATCCGGC	AACCAGCACG	CTGGTGTCTG	GTGCGATCAC	1200
CGCCGTGTGC	GATCGAGCGT	TTCCCGAACG	ATTCGTCGG	TCAACGGGGG	CAGGGGACGT	1260
TCTGGCCGTG	CGACGAGAAC	CGAGCCTTCC	CGAACGAGTT	CGACACCGGT	CGGGGCCGGC	1320
TCAATCTCGA	TGCGCCCATC	GCGCTCGGTG	ATCTCCACCT	GGTCGTTCCC	GCGCAAGCCA	1380
AGGCGCTCGC	GAATCCGCTT	GGGAATCACC	AGACGTCCTG	CGACATCGAT	GGTTGTTCGC	1440
ATGGTAGGAA	ATTTACCATC	GCACGTTCCTA	TAGGCGTGTC	CTGCGCGGGA	TGTCGGGACG	1500
ATCCGCTAGC	GTATCGAACG	ATTGTTTCGG	AAATGGCTGA	GGGAGCGTGC	GGTGCGGGTG	1560
ATGGGTGTGC	ATCCCGGGTT	GACCCGATGC	GGGCTGTCGC	TCATCGAGAG	TGGGCGTGGT	1620
CGGCAGCTCA	CCGCGCTGGA	TGTCGACGTG	GTGCGCACAC	CGTCGGATGC	GGCCTTGCGG	1680
CAGCGCCTGT	TGGCCATCAG	CGATGCCGTC	GAGCACTGGC	TGGACACCCA	TCATCCGGAG	1740
GTGGTGGCTA	TCGAACGGGT	GTTCTCTCAG	CTCAACGTGA	CCACGGTGAT	GGGCACCGCG	1800
CAGGCCGGCG	GCGTGATCGC	CCTGGCGGCG	GCCAAACGTG	GTGTCGACGT	GCATTTCCAT	1860
ACCCCCAGCG	AGGTCAAGGC	GGCGGTCACT	GGCAACGGTT	CCGCAGACAA	GGCTCAGGTC	1920
ACC						1923

(2) INFORMATION FOR SEQ ID NO:186:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 1055 base pairs

(B) TYPE: nucleic acid

- (C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:186:

CTGGCGTGCC AGTGTCACCG GCGATATGAC GTCGGCATTG AATTTCGCGG CCCC GCCGGA	60
CCCGTCGCCA CCCAATCTGG ACCACCCGGT CCGTCAATTG CCGAAGGTG CCAAGTGCGT	120
GCCCAATGTG GTGCTGGGTT TCTTGAACGA AGGCCTGCCG TATCGGGTGC CCTACCCCCA	180
AACAACGCCA GTCCAGGAAT CCGGTCCCGC GCGGCCGATT CCCAGCGGCA TCTGCTAGCC	240
GGGGATGGTT CAGACGTAAC GGTGGGCTAG GTCGAAACCC GCGCCAGGGC CGCTGGACGG	300
GCTCATGGCA GCGAAATTAG AAAACCCGGG ATATTGTCCG CGGATTGTCA TACGATGCTG	360
AGTGCTTGGT GGTTCGTGTT TAGCCATTGA GTGTGGATGT GTTGAGACCC TGGCCTGGAA	420
GGGGACAACG TGCTTTTGCC TCTTGGTCCG CCTTTGCCGC CCGACGCGGT GGTGGCGAAA	480
CGGGCTGAGT CGGGAATGCT CGGCGGGTTG TCGGTTCCGC TCAGCTGGGG AGTGGCTGTG	540
CCACCCGATG ATTATGACCA CTGGGCGCCT GCGCCGGAGG ACGGCGCCGA TGTCGATGTC	600
CAGGCGGCCG AAGGGGCGGA CGCAGAGGCC GCGGCCATGG ACGAGTGGGA TGAGTGGCAG	660
GCGTGGAACG AGTGGGTGGC GGAGAACGCT GAACCCCGCT TTGAGGTGCC ACGGAGTAGC	720
AGCAGCGTGA TTCCGCATTC TCCGGCGGCC GGCTAGGAGA GGGGGCGCAG ACTGTCGTTA	780
TTTGACCACT GATCGGCGGT CTCGGTGTTT CCGCGGCCGG CTATGACAAC AGTCAATGTG	840
CATGACAAGT TACAGGTATT AGGTCCAGGT TCAACAAGGA GACAGGCAAC ATGGCAACAC	900
GTTTTATGAC GGATCCGCAC GCGATGCGGG ACATGGCGGG CCGTTTTGAG GTGCACGCCC	960
AGACGGTGGA GGACGAGGCT CGCCGGATGT GGGCGTCCGC GCAAAACATC TCGGGNGCGG	1020
GCTGGAGTGG CATGGCCGAG GCGACCTCGC TAGAC	1055

(2) INFORMATION FOR SEQ ID NO:187:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 359 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:187:

CCGCCTCGTT GTTGGCATAC TCCGCCGCGG CCGCCTCGAC CGCACTGGCC GTGGCGTGTG	60
TCCGGGCTGA CCACCGGGAT CGCCGAACCA TCCGAGATCA CCTCGCAATG ATCCACCTCG	120
CGCAGCTGGT CACCCAGCCA CCGGGCGGTG TGCGACAGCG CCTGCATCAC CTTGGTATAG	180
CCGTCGCGCC CCAGCCGCAG GAAGTTGTAG TACTGGCCCA CCACCTGGTT ACCGGGACGG	240
GAGAAGTTCA GGGTGAAGGT CGGCATGTCG CCGCCGAGGT AGTTGACCCG GAAAACCAGA	300
TCCTCCGGCA GGTGCTCGGG CCCGCGCCAC ACGACAAACC CGACGCCGGG ATAGGTCAG	359

(2) INFORMATION FOR SEQ ID NO:188:

- (i) SEQUENCE CHARACTERISTICS:
- (A) LENGTH: 350 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:188:

AACGGGCCCCG TGGGCACCGC TCCTCTAAGG GCTCTCGTTG GTCGCATGAA GTGCTGGAAG	60
GATGCATCTT GGCAGATTCC CGCCAGAGCA AAACAGCCGC TAGTCCTAGT CCGAGTCGCC	120
CGCAAAGTTC CTCGAATAAC TCCGTACCCG GAGCGCCAAA CCGGGTCTCC TTCGCTAAGC	180
TGCGCGAACC ACTTGAGGTT CCGGGACTCC TTGACGTCCA GACCGATTCTG TTCGAGTGGC	240
TGATCGGTTT GCCGCGCTGG CGCGAATCCG CCGCCGAGCG GGGTGATGTC AACCCAGTGG	300
GTGGCCTGGA AGAGGTGCTC TACGAGCTGT CTCCGATCGA GGACTTCTCC	350

(2) INFORMATION FOR SEQ ID NO:189:

- (i) SEQUENCE CHARACTERISTICS:
- (A) LENGTH: 679 amino acids
 - (B) TYPE: amino acid
 - (C) STRANDEDNESS:
 - (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:189:

201

Glu Gln Pro Lys Gly Pro Phe Gly Glu Val Ile Glu Ala Phe Ala Asp
 1 5 10 15
 Gly Leu Ala Gly Lys Gly Lys Gln Ile Asn Thr Thr Leu Asn Ser Leu
 20 25 30
 Ser Gln Ala Leu Asn Ala Leu Asn Glu Gly Arg Gly Asp Phe Phe Ala
 35 40 45
 Val Val Arg Ser Leu Ala Leu Phe Val Asn Ala Leu His Gln Asp Asp
 50 55 60
 Gln Gln Phe Val Ala Leu Asn Lys Asn Leu Ala Glu Phe Thr Asp Arg
 65 70 75 80
 Leu Thr His Ser Asp Ala Asp Leu Ser Asn Ala Ile Gln Gln Phe Asp
 85 90 95
 Ser Leu Leu Ala Val Ala Arg Pro Phe Phe Ala Lys Asn Arg Glu Val
 100 105 110
 Leu Thr His Asp Val Asn Asn Leu Ala Thr Val Thr Thr Thr Leu Leu
 115 120 125
 Gln Pro Asp Pro Leu Asp Gly Leu Glu Thr Val Leu His Ile Phe Pro
 130 135 140
 Thr Leu Ala Ala Asn Ile Asn Gln Leu Tyr His Pro Thr His Gly Gly
 145 150 155 160
 Val Val Ser Leu Ser Ala Phe Thr Asn Phe Ala Asn Pro Met Glu Phe
 165 170 175
 Ile Cys Ser Ser Ile Gln Ala Gly Ser Arg Leu Gly Tyr Gln Glu Ser
 180 185 190
 Ala Glu Leu Cys Ala Gln Tyr Leu Ala Pro Val Leu Asp Ala Ile Lys
 195 200 205
 Phe Asn Tyr Phe Pro Phe Gly Leu Asn Val Ala Ser Thr Ala Ser Thr
 210 215 220
 Leu Pro Lys Glu Ile Ala Tyr Ser Glu Pro Arg Leu Gln Pro Pro Asn
 225 230 235 240
 Gly Tyr Lys Asp Thr Thr Val Pro Gly Ile Trp Val Pro Asp Thr Pro
 245 250 255
 Leu Ser His Arg Asn Thr Gln Pro Gly Trp Val Val Ala Pro Gly Met
 260 265 270
 Gln Gly Val Gln Val Gly Pro Ile Thr Gln Gly Leu Leu Thr Pro Glu
 275 280 285

Ser Leu Ala Glu Leu Met Gly Gly Pro Asp Ile Ala Pro Pro Ser Ser
 290 295 300
 Gly Leu Gln Thr Pro Pro Gly Pro Pro Asn Ala Tyr Asp Glu Tyr Pro
 305 310 315 320
 Val Leu Pro Pro Ile Gly Leu Gln Ala Pro Gln Val Pro Ile Pro Pro
 325 330 335
 Pro Pro Pro Gly Pro Asp Val Ile Pro Gly Pro Val Pro Pro Val Leu
 340 345 350
 Ala Ala Ile Val Phe Pro Arg Asp Arg Pro Ala Ala Ser Glu Asn Phe
 355 360 365
 Asp Tyr Met Gly Leu Leu Leu Leu Ser Pro Gly Leu Ala Thr Phe Leu
 370 375 380
 Phe Gly Val Ser Ser Ser Pro Ala Arg Gly Thr Met Ala Asp Arg His
 385 390 395 400
 Val Leu Ile Pro Ala Ile Thr Gly Leu Ala Leu Ile Ala Ala Phe Val
 405 410 415
 Ala His Ser Trp Tyr Arg Thr Glu His Pro Leu Ile Asp Met Arg Leu
 420 425 430
 Phe Gln Asn Arg Ala Val Ala Gln Ala Asn Met Thr Met Thr Val Leu
 435 440 445
 Ser Leu Gly Leu Phe Gly Ser Phe Leu Leu Leu Pro Ser Tyr Leu Gln
 450 455 460
 Gln Val Leu His Gln Ser Pro Met Gln Ser Gly Val His Ile Ile Pro
 465 470 475 480
 Gln Gly Leu Gly Ala Met Leu Ala Met Pro Ile Ala Gly Ala Met Met
 485 490 495
 Asp Arg Arg Gly Pro Ala Lys Ile Val Leu Val Gly Ile Met Leu Ile
 500 505 510
 Ala Ala Gly Leu Gly Thr Phe Ala Phe Gly Val Ala Arg Gln Ala Asp
 515 520 525
 Tyr Leu Pro Ile Leu Pro Thr Gly Leu Ala Ile Met Gly Met Gly Met
 530 535 540
 Gly Cys Ser Met Met Pro Leu Ser Gly Ala Ala Val Gln Thr Leu Ala
 545 550 555 560
 Pro His Gln Ile Ala Arg Gly Ser Thr Leu Ile Ser Val Asn Gln Gln
 565 570 575
 Val Gly Gly Ser Ile Gly Thr Ala Leu Met Ser Val Leu Leu Thr Tyr

(2) INFORMATION FOR SEQ ID NO:190:

(A) LENGTH: 120 amino acids
(B) TYPE: amino acid
(C) STRANDEDNESS:
(D) TOPOLOGY: linear

Thr 1	Pro	Glu	Lys 5	Ser 5	Phe	Val	Asp	Asp	Leu 10	Asp	Ile	Asp	Ser	Leu 15	Ser
Met	Val	Glu	Ile 20	Ala	Val	Gln	Thr	Glu 25	Asp	Lys	Tyr	Gly	Val 30	Lys	Ile
Pro	Asp	Glu 35	Asp	Leu	Ala	Gly	Leu 40	Arg	Thr	Val	Gly	Asp 45	Val	Val	Ala
Tyr	Ile 50	Gln	Lys	Leu	Glu	Glu 55	Glu	Asn	Pro	Glu	Ala 60	Ala	Gln	Ala	Leu
Arg 65	Ala	Lys	Ile	Glu	Ser 70	Glu	Asn	Pro	Asp	Ala 75	Ala	Arg	Ala	Asp	Arg 80
Cys	Val	Ser	Pro	Thr 85	Ser	Gln	Ala	Arg	Asp 90	Ala	Arg	Arg	Pro	Leu 95	Ala
Arg	Ser	Ala	Arg 100	Leu	Ala	Cys	Arg	Arg 105	Leu	Pro	Ala	Ser	Val 110	Pro	Thr

Thr Arg Arg Asp Pro Arg Glu Arg
 115 120

(2) INFORMATION FOR SEQ ID NO:191:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 89 amino acids
- (B) TYPE: amino acid
- (C) STRANDEDNESS:
- (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:191:

Leu Ala Cys Gln Cys His Arg Arg Tyr Asp Val Gly Ile Gln Phe Arg
 1 5 10 15
 Gly Pro Ala Gly Pro Val Ala Thr Gln Ser Gly Pro Pro Gly Pro Ser
 20 25 30
 Ile Ala Glu Gly Arg Gln Val Arg Ala Gln Cys Gly Ala Gly Phe Leu
 35 40 45
 Glu Arg Arg Pro Ala Val Ser Gly Ala Leu Pro Pro Asn Asn Ala Ser
 50 55 60
 Pro Gly Ile Arg Ser Arg Ala Ala Asp Ser Gln Arg His Leu Leu Ala
 65 70 75 80
 Gly Asp Gly Ser Asp Val Thr Val Gly
 85

(2) INFORMATION FOR SEQ ID NO:192:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 119 amino acids
- (B) TYPE: amino acid
- (C) STRANDEDNESS:
- (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:192:

Ala Ser Leu Leu Ala Tyr Ser Ala Ala Ala Ser Thr Ala Leu Ala
 1 5 10 15
 Val Ala Cys Val Arg Ala Asp His Arg Asp Arg Arg Thr Ile Arg Asp
 20 25 30

His Leu Ala Met Ile His Leu Ala Gln Leu Val Thr Gln Pro Pro Gly
 35 40 45
 Gly Val Arg Gln Arg Leu His His Leu Gly Ile Ala Val Ala Pro Gln
 50 55 60
 Pro Gln Glu Val Val Val Leu Ala His His Leu Val Thr Gly Thr Gly
 65 70 75 80
 Glu Val Gln Gly Glu Gly Arg His Val Ala Ala Glu Val Val Asp Pro
 85 90 95
 Glu Asn Gln Ile Leu Arg Gln Val Leu Gly Pro Ala Pro His Asp Lys
 100 105 110
 Pro Asp Ala Gly Ile Gly Gln
 115

(2) INFORMATION FOR SEQ ID NO:193:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 116 amino acids
- (B) TYPE: amino acid
- (C) STRANDEDNESS:
- (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:193:

Arg Ala Arg Gly His Arg Ser Ser Lys Gly Ser Arg Trp Ser His Glu
 1 5 10 15
 Val Leu Glu Gly Cys Ile Leu Ala Asp Ser Arg Gln Ser Lys Thr Ala
 20 25 30
 Ala Ser Pro Ser Pro Ser Arg Pro Gln Ser Ser Ser Asn Asn Ser Val
 35 40 45
 Pro Gly Ala Pro Asn Arg Val Ser Phe Ala Lys Leu Arg Glu Pro Leu
 50 55 60
 Glu Val Pro Gly Leu Leu Asp Val Gln Thr Asp Ser Phe Glu Trp Leu
 65 70 75 80
 Ile Gly Ser Pro Arg Trp Arg Glu Ser Ala Ala Glu Arg Gly Asp Val
 85 90 95
 Asn Pro Val Gly Gly Leu Glu Glu Val Leu Tyr Glu Leu Ser Pro Ile
 100 105 110
 Glu Asp Phe Ser
 115

(2) INFORMATION FOR SEQ ID NO:194:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 811 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:194:

```

TGCTACGCAG CAATCGCTTT GGTGACAGAT GTGGATGCCG GCGTCGCTGC TGGCGATGGC      60
GTGAAAGCCG CCGACGTGTT CGCCGCATTC GGGGAGAACA TCGAACTGCT CAAAAGGCTG      120
GTGCGGGCCG CCATCGATCG GGTGCGCCGAC GAGCGCACGT GCACGCACTG TCAACACCAC      180
GCCGGTGTTT CGTTGCCGTT CGAGCTGCCA TGAGGGTGCT GCTGACCGGC GCGGCCGGCT      240
TCATCGGGTC GCGCGTGGAT GCGGCGTTAC GGGCTGCGGG TCACGACGTG GTGGGCGTCG      300
ACGCGCTGCT GCCCCGCCG CACGGGCCAA ACCCGGTGCT GCCACCGGGC TGCCAGCGGG      360
TCGACGTGCG CGACGCCAGC GCGCTGGCCC CGTTGTTGGC CGGTGTCGAT CTGGTGTGTC      420
ACCAGGCCGC CATGGTGGGT GCCGGCGTCA ACGCCGCCGA CGCACCCGCC TATGGCGGCC      480
ACAACGATTT CGCCACCACG GTGCTGCTGG CGCAGATGTT CGCCGCCGGG GTCCGCCGTT      540
TGGTGCTGGC GTCGTCGATG GTGGTTTACG GGCAGGGGCG CTATGACTGT CCCCAGCATG      600
GACCGGTCTGA CCCGCTGCCG CGGCGGCGAG CCGACCTGGA CAATGGGGTC TTCGAGCACC      660
GTTGCCCCGGG GTGCGGCGAG CCAGTCATCT GGCAATTGGT CGACGAAGAT GCCCCGTTGC      720
GCCCCGCGCAG CCTGTACGCG GCAGCAAGAC CGCGCAGGAG CACTACGCGC TGGCGTGGTC      780
GGAAACGAAT GGCGGTTCCG TGGTGGCGTT G                                     811

```

(2) INFORMATION FOR SEQ ID NO:195:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 966 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:195:

GTCCCGCGAT GTGGCCGAGC ATGACTTTCG GCAACACCGG CGTAGTAGTC GAAGATATCG	60
GACTTTGTGG TCCCGGTGGC GGGATAGAGC ACCTGTCGGC GTTGGTCAGC GTCACCCGTT	120
GCTCGGACGC CGAACCCATG CTTTCAACGT AGCCTGTCGG TCACACAAGT CGCGAGCGTA	180
ACGTCACGGT CAAATATCGC GTGGAATTTT GCCGTGACGT TCCGCTCGCG GACAATCAAG	240
GCATACTCAC TTACATGCGA GCCATTTGGA CGGGTTCGAT CGCCTTCGGG CTGGTGAACG	300
TGCCGGTCAA GGTGTACAGC GCTACCGCAG ACCACGACAT CAGGTTCCAC CAGGTGCACG	360
CCAAGGACAA CGGACGCATC CGGTACAAGC GCGTCTGCGA GGCGTGTGGC GAGGTGGTGC	420
ACTACCGCGA TCTTGCCCCG GCCTACGAGT CCGGCGACGG CCAAATGGTG GCGATCACCG	480
ACGACGACAT CGCCAGCTTG CCTGAAGAAC GCAGCCGGGA GATCGAGGTG TTGGAGTTTCG	540
TCCCGCCGCG CGACGTGGAC CCGATGATGT TCGACCGCAG CTACTTTTTG GAGCCTGATT	600
CGAAGTCGTC GAAATCGTAT GTGCTGCTGG CTAAGACACT CGCCGAGACC GACCGGATGG	660
CGATCGTGGA TCGCCCCACC GGCCGTGAAT GCAGGAAAAA TAAGAGCCGC TATCCACAAT	720
TCGGCGTCGA GCTCGGCTAC CACAAACGGT AGAACGATCG AGACATTCCC GAGCTGAAGT	780
GCGGCGCTAT AGAAGCCGCT CTGCGCGATT ATCAAACGCA AAATACGCTT ACTCATGCCA	840
TCGGCGCTGC TCACCCGATG CGACGTTTTT GCCACGCTCC ACCGCCTGCC GCGCGACCTC	900
AAGTGGGCAT GCATCCCACC CGTTCCCGGA AACCGGTTCC GGCGGGTCGG CTCATCGCTT	960
CATCCT	966

(2) INFORMATION FOR SEQ ID NO:196:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 2367 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:196:

CCGCACCGCC GGCAATACCG CCAGCGCCAC CGTTACCGCC GTTTCGCGCC TTGCCCCCGT	60
TGCCGCCCCGT CCCGCCGGCC CCGCCGATGG AGTTCTCATC GCCAAAAGTA CTGGCGTTGC	120
CACCGGAGCC GCCGTTGCCG CCGTCACCGC CAGCCCCGCC GACTCCACCG GCCCCACCGA	180


```

CGGCCGCATT AGCCGCGCTC ACCTGCGAAC CAATAGTCGA TAAATCCAAA GCCGCAGTTG      1860
CCAGCAGCTG CGGCGTCGCG ATCACCAAGG ACACCTCGCA CCTCCGATA CCCCATATCG      1920
CCGCACCGTG TCCCCAGCGG CCACGTGACC TTTGGTCGCT GGCTGGCGGC CCTGACTATG      1980
GCCGCGACGG CCCTCGTTCT GATTGCCCCC GGCGCGCAGC TTGTTGCGCG AGTTGAAGAC      2040
GGGAGGACAG GCCGAGCTTG GTGTAGACGT GGGTCAAGTG GGAATGCACG GTCCGCGGCG      2100
AGATGAATAG GCGGACGCCG ATCTCCTTGT TGCTGAGTCC CTCACCGACC AGTAGAGCCA      2160
CCTCAAGCTC TGTCGGTGTC AACGCGCCCC AGCCACTTGT CGGGCGTTTC CGTGCACCGC      2220
GGCCTCGTTG CGCGTACGCG ATCGCCTCAT CGATCGATAA CGCAGTTCCT TCGGCCCAGG      2280
CATCGTCGAA CTCGCTGTCA CCCATGGATT TTCGAAGGGT GGCTAGCGAC GAGTTACAGC      2340
CCGCCTGGTA GATCCCGAAG CGGACCG                                     2367

```

(2) INFORMATION FOR SEQ ID NO:197:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 376 amino acids
- (B) TYPE: amino acid
- (C) STRANDEDNESS:
- (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:197:

```

Gln Pro Ala Gly Ala Thr Ile Ala Ala Ser Ser Pro Cys Ala Thr Val
1           5           10           15
Gly Ala Gly Gly Gly Thr Gly Ser Pro Val Thr Thr Glu Thr Ala Ala
20           25           30
Thr Thr Gly Arg Gly Gly Ser Gly Asp Val Tyr Glu Ser Ala Ala Ser
35           40           45
Gly Ala Ala Ala Thr Thr Pro Thr Ala Gly Gly Tyr Thr Val Gly Pro
50           55           60
Val Ala Thr Ile Thr Ala Lys Gly Ala Arg Asn Val Ala Leu Arg Asp
65           70           75           80
Ser Ala Val Ala Ala Val Ala Ala Ala Ala Thr Gly Ser Gly Gly Thr
85           90           95
Ala Val Thr Thr Gly Thr Ala Gly Gly Leu Ala Arg Ala Cys Arg Arg
100          105          110

```

210

Gly Gly Thr Val Ala Ala Gly Ala Thr Gly Arg Arg Ala Gly Ser Ala
 115 120 125
 Met Ala Ala Arg Ala Ala Val Ala Ala Gly Leu Ile Thr Asp Ala Gly
 130 135 140
 His Ile Cys Arg Ala Val Pro Gly Ala Gly Arg Gly Ala Gly Arg Gly
 145 150 155 160
 Ile Asp Pro Val Cys Pro Gly Glu Ala Gly Ala Ala Gly Thr Thr Gly
 165 170 175
 Ala Ala Met Ala Glu Gln Pro Gly Val Ala Ala Val Thr Ala Arg Thr
 180 185 190
 Pro Asp Ala Cys Gly His Ala Gly Ala Ala Asp Thr Ala Val Ala Ala
 195 200 205
 Val Ala Pro Gln Pro Pro Pro Val Pro Thr Gly Thr Ala Gly Arg Ala
 210 215 220
 Gly Thr Thr Gly Pro Ala Val Ala Ala Val Ala Asp Gln Pro Gly Arg
 225 230 235 240
 Ala Ser Ala Ala Ala Gly Leu Thr Glu Pro Ala Ser Arg Ala Val Ala
 245 250 255
 Thr Val Ala Lys Gln Gln Pro Ala Gly Arg Ala Arg Leu Pro Gly Cys
 260 265 270
 Arg Pro Val Gly Ala Val Ser Asp Gln Arg Ala Pro Gln Lys Arg Leu
 275 280 285
 Gly Gly Arg Ile His Arg Thr Gln Gln Thr Pro Leu Asn Ser Gly Phe
 290 295 300
 Ser Ala Gly Ile Pro Thr Arg Gly Arg Ser Gln Arg Leu His Lys Leu
 305 310 315 320
 Leu Val Lys Arg Cys His Leu Tyr Ala Glu Arg Leu Ile Leu Pro Ser
 325 330 335
 Met Gly Pro Glu Gln Pro Arg Asn Arg Arg Arg His Phe Ile Gly Ser
 340 345 350
 Arg Ser His His Phe Arg Arg Arg Asp Arg Arg Gly Arg Ile Ser Arg
 355 360 365
 Ala His Leu Arg Thr Asn Ser Arg
 370 375

(2) INFORMATION FOR SEQ ID NO:198:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 2852 base pairs

(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:198:

GGCCAAACG CCCC GGCGAT CGCGGCCACC GAGGCCGCCT ACGACCAGAT GTGGGCCCAG	60
GACGTGGCGG CGATGTTTGG CTACCATGCC GGGGCTTCGG CGGCCGTCTC GCGTTGACA	120
CCGTTCGGCC AGGCGTGCC GACCGTGGCG GGC GGCGGTG CGCTGGTCAG CGCGCCGCG	180
GCTCAGGTGA CCACGCGGGT CTTCCGCAAC CTGGGCTTGG CGAACGTCCG CGAGGGCAAC	240
GTCCGCAACG GTAATGTCCG GAACTTCAAT CTCGGCTCGG CCAACATCGG CAACGGCAAC	300
ATCGGCAGCG GCAACATCGG CAGCTCCAAC ATCGGGTTTG GCAACGTGGG TCCTGGGTTG	360
ACCGCAGCGC TGAACAACAT CGGTTTCGGC AACACCGGCA GCAACAACAT CGGGTTTGGC	420
AACACCGGCA GCAACAACAT CGGGTTTCGGC AATACCGGAG ACGGCAACCG AGGTATCGGG	480
CTCACGGGTA GCGGTTTGTG GGGGTTTCGGC GGCCTGAACT CGGGCACCGG CAACATCGGT	540
CTGTTCAACT CGGGCACCGG AAACGTTCGGC ATCGGCAACT CGGGTACCGG GAACTGGGGC	600
ATTGGCAACT CGGGCAACAG CTACAACACC GGTTTTGGCA ACTCCGGCGA CGCCAACACG	660
GGCTTCTTCA ACTCCGGAAT AGCCAACACC GGCGTCGGCA ACGCCGGCAA CTACAACACC	720
GGTAGCTACA ACCCGGGCAA CAGCAATACC GGCGGCTTCA ACATGGGCCA GTACAACACG	780
GGCTACCTGA ACAGCGGCAA CTACAACACC GGCTTGCAA ACTCCGGCAA TGTCAACACC	840
GGCGCCTTCA TTA CTGGCAA CTTCAACAAC GGCTTCTTGT GGCGCGGCGA CCACCAAGGC	900
CTGATTTTCG GGAGCCCCGG CTTCTTCAAC TCGACCAGTG CGCCGTCGTC GGGATTCTTC	960
AACAGCGGTG CCGGTAGCGC GTCCGGCTTC CTGAACTCCG GTGCCAACAA TTCTGGCTTC	1020
TTCAACTCTT CGTCGGGGGC CATCGGTAAC TCCGGCCTGG CAAACGCGGG CGTGCTGGTA	1080
TCGGGCGTGA TCAACTCGGG CAACACCGTA TCGGGTTTGT TCAACATGAG CCTGGTGGCC	1140
ATCACAACGC CGGCCTTGAT CTCGGGCTTC TTCAACACCG GAAGCAACAT GTCGGGATTT	1200
TTCGGTGGCC CACCGGTCTT CAATCTCGGC CTGGCAAACC GGGGCGTCGT GAACATTCTC	1260
GGCAACGCCA ACATCGGCAA TTACAACATT CTCGGCAGCG GAAACGTCCG TGA CTTCAAC	1320
ATCCTTGGCA GCGGCAACCT CGGCAGCCAA AACATCTTGG GCAGCGGCAA CGTCGGCAGC	1380

TTCAATATCG GCAGTGGAAA CATCGGAGTA TTCAATGTCG GTTCCGGAAG CCTGGGAAAC	1440
TACAACATCG GATCCGGAAA CCTCGGGATC TACAACATCG GTTTTGAAA CGTCGGCGAC	1500
TACAACGTCG GCTTCGGGAA CGCGGGCGAC TTCAACCAAG GCTTTGCCAA CACCGGCAAC	1560
AACAACATCG GGTTCGCCAA CACCGGCAAC AACAACATCG GCATCGGGCT GTCCGGCGAC	1620
AACCAGCAGG GCTTCAATAT TGCTAGCGGC TGGAACTCGG GCACCGGCAA CAGCGGCCTG	1680
TTCAATTCGG GCACCAATAA CGTTGGCATC TTCAACGCGG GCACCGGAAA CGTCGGCATC	1740
GCAAACTCGG GCACCGGGAA CTGGGGTATC GGAACCCGG GTACCGACAA TACCGGCATC	1800
CTCAATGCTG GCAGCTACAA CACGGGCATC CTCAACGCCG GCGACTTCAA CACGGGCTTC	1860
TACAACACGG GCAGCTACAA CACCGGCGGC TTCAACGTCG GTAACACCAA CACCGGCAAC	1920
TTCAACGTGG GTGACACCAA TACCGGCAGC TATAACCCGG GTGACACCAA CACCGGCTTC	1980
TTCAATCCCG GCAACGTCAA TACCGGCGCT TTCGACACGG GCGACTTCAA CAATGGCTTC	2040
TTGGTGGCGG GCGATAACCA GGGCCAGATT GCCATCGATC TCTCGGTCAC CACTCCATTC	2100
ATCCCCATAA ACGAGCAGAT GGTCAATTGAC GTACACAACG TAATGACCTT CGGCGGCAAC	2160
ATGATCACGG TCACCGAGGC CTCGACCGTT TTCCCCAAA CCTTCTATCT GAGCGGTTTG	2220
TTCTTCTTCG GCCCGGTCAA TCTCAGCGCA TCCACGCTGA CCGTTCCGAC GATCACCCCTC	2280
ACCATCGGCG GACCGACGGT GACCGTCCCC ATCAGCATTG TCGGTGCTCT GGAGAGCCGC	2340
ACGATTACCT TCCTCAAGAT CGATCCGGCG CCGGGCATCG GAAATTCGAC CACCAACCCC	2400
TCGTCCGGCT TCTTCAACTC GGGCACCGGT GGCACATCTG GCTTCCAAA CGTCGGCGGC	2460
GGCAGTTCAG GCGTCTGGAA CAGTGGTTTG AGCAGCGCGA TAGGGAATTC GGGTTTCCAG	2520
AACCTCGGCT CGCTGCAGTC AGGCTGGGCG AACCTGGGCA ACTCCGTATC GGGCTTTTTTC	2580
AACACCAGTA CGGTGAACCT CTCCACGCCG GCCAATGTCT CGGGCCTGAA CAACATCGGC	2640
ACCAACCTGT CCGGCGTGTT CCGCGGTCCG ACCGGGACGA TTTTCAACGC GGGCCTTGCC	2700
AACCTGGGCC AGTTGAACAT CGGCAGCGCC TCGTGCCGAA TTCGGCACGA GTTAGATACG	2760
GTTTCAACAA TCATATCCGC GTTTTGCGGC AGTGCATCAG ACGAATCGAA CCCGGGAAGC	2820
GTAAGCGAAT AAACCGAATG GCGGCCTGTC AT	2852

(2) INFORMATION FOR SEQ ID NO:199:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 943 amino acids

(B) TYPE: amino acid
 (C) STRANDEDNESS:
 (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:199:

Gly	Gln	Asn	Ala	Pro	Ala	Ile	Ala	Ala	Thr	Glu	Ala	Ala	Tyr	Asp	Gln	1	5	10	15
Met	Trp	Ala	Gln	Asp	Val	Ala	Ala	Met	Phe	Gly	Tyr	His	Ala	Gly	Ala	20	25	30	
Ser	Ala	Ala	Val	Ser	Ala	Leu	Thr	Pro	Phe	Gly	Gln	Ala	Leu	Pro	Thr	35	40	45	
Val	Ala	Gly	Gly	Gly	Ala	Leu	Val	Ser	Ala	Ala	Ala	Ala	Gln	Val	Thr	50	55	60	
Thr	Arg	Val	Phe	Arg	Asn	Leu	Gly	Leu	Ala	Asn	Val	Arg	Glu	Gly	Asn	65	70	75	80
Val	Arg	Asn	Gly	Asn	Val	Arg	Asn	Phe	Asn	Leu	Gly	Ser	Ala	Asn	Ile	85	90	95	
Gly	Asn	Gly	Asn	Ile	Gly	Ser	Gly	Asn	Ile	Gly	Ser	Ser	Asn	Ile	Gly	100	105	110	
Phe	Gly	Asn	Val	Gly	Pro	Gly	Leu	Thr	Ala	Ala	Leu	Asn	Asn	Ile	Gly	115	120	125	
Phe	Gly	Asn	Thr	Gly	Ser	Asn	Asn	Ile	Gly	Phe	Gly	Asn	Thr	Gly	Ser	130	135	140	
Asn	Asn	Ile	Gly	Phe	Gly	Asn	Thr	Gly	Asp	Gly	Asn	Arg	Gly	Ile	Gly	145	150	155	160
Leu	Thr	Gly	Ser	Gly	Leu	Leu	Gly	Phe	Gly	Gly	Leu	Asn	Ser	Gly	Thr	165	170	175	
Gly	Asn	Ile	Gly	Leu	Phe	Asn	Ser	Gly	Thr	Gly	Asn	Val	Gly	Ile	Gly	180	185	190	
Asn	Ser	Gly	Thr	Gly	Asn	Trp	Gly	Ile	Gly	Asn	Ser	Gly	Asn	Ser	Tyr	195	200	205	
Asn	Thr	Gly	Phe	Gly	Asn	Ser	Gly	Asp	Ala	Asn	Thr	Gly	Phe	Phe	Asn	210	215	220	
Ser	Gly	Ile	Ala	Asn	Thr	Gly	Val	Gly	Asn	Ala	Gly	Asn	Tyr	Asn	Thr	225	230	235	240

214

Gly Ser Tyr Asn Pro Gly Asn Ser Asn Thr Gly Gly Phe Asn Met Gly
 245 250 255
 Gln Tyr Asn Thr Gly Tyr Leu Asn Ser Gly Asn Tyr Asn Thr Gly Leu
 260 265 270
 Ala Asn Ser Gly Asn Val Asn Thr Gly Ala Phe Ile Thr Gly Asn Phe
 275 280 285
 Asn Asn Gly Phe Leu Trp Arg Gly Asp His Gln Gly Leu Ile Phe Gly
 290 295 300
 Ser Pro Gly Phe Phe Asn Ser Thr Ser Ala Pro Ser Ser Gly Phe Phe
 305 310 315 320
 Asn Ser Gly Ala Gly Ser Ala Ser Gly Phe Leu Asn Ser Gly Ala Asn
 325 330 335
 Asn Ser Gly Phe Phe Asn Ser Ser Ser Gly Ala Ile Gly Asn Ser Gly
 340 345 350
 Leu Ala Asn Ala Gly Val Leu Val Ser Gly Val Ile Asn Ser Gly Asn
 355 360 365
 Thr Val Ser Gly Leu Phe Asn Met Ser Leu Val Ala Ile Thr Thr Pro
 370 375 380
 Ala Leu Ile Ser Gly Phe Phe Asn Thr Gly Ser Asn Met Ser Gly Phe
 385 390 395 400
 Phe Gly Gly Pro Pro Val Phe Asn Leu Gly Leu Ala Asn Arg Gly Val
 405 410 415
 Val Asn Ile Leu Gly Asn Ala Asn Ile Gly Asn Tyr Asn Ile Leu Gly
 420 425 430
 Ser Gly Asn Val Gly Asp Phe Asn Ile Leu Gly Ser Gly Asn Leu Gly
 435 440 445
 Ser Gln Asn Ile Leu Gly Ser Gly Asn Val Gly Ser Phe Asn Ile Gly
 450 455 460
 Ser Gly Asn Ile Gly Val Phe Asn Val Gly Ser Gly Ser Leu Gly Asn
 465 470 475 480
 Tyr Asn Ile Gly Ser Gly Asn Leu Gly Ile Tyr Asn Ile Gly Phe Gly
 485 490 495
 Asn Val Gly Asp Tyr Asn Val Gly Phe Gly Asn Ala Gly Asp Phe Asn
 500 505 510
 Gln Gly Phe Ala Asn Thr Gly Asn Asn Asn Ile Gly Phe Ala Asn Thr
 515 520 525
 Gly Asn Asn Asn Ile Gly Ile Gly Leu Ser Gly Asp Asn Gln Gln Gly

530	535	540
Phe Asn Ile Ala Ser Gly	Trp Asn Ser Gly Thr Gly	Asn Ser Gly Leu
545	550	555 560
Phe Asn Ser Gly Thr	Asn Asn Val Gly Ile	Phe Asn Ala Gly Thr Gly
565	570	575
Asn Val Gly Ile Ala Asn Ser	Gly Thr Gly Asn Trp Gly	Ile Gly Asn
580	585	590
Pro Gly Thr Asp Asn Thr	Gly Ile Leu Asn Ala Gly	Ser Tyr Asn Thr
595	600	605
Gly Ile Leu Asn Ala Gly	Asp Phe Asn Thr Gly	Phe Tyr Asn Thr Gly
610	615	620
Ser Tyr Asn Thr Gly	Gly Phe Asn Val Gly	Asn Thr Asn Thr Gly
625	630	635 640
Phe Asn Val Gly Asp Thr	Asn Thr Gly Ser Tyr Asn	Pro Gly Asp Thr
645	650	655
Asn Thr Gly Phe Phe Asn	Pro Gly Asn Val Asn Thr	Gly Ala Phe Asp
660	665	670
Thr Gly Asp Phe Asn Asn	Gly Phe Leu Val Ala Gly	Asp Asn Gln Gly
675	680	685
Gln Ile Ala Ile Asp Leu	Ser Val Thr Thr Pro	Phe Ile Pro Ile Asn
690	695	700
Glu Gln Met Val Ile Asp	Val His Asn Val Met	Thr Phe Gly Gly Asn
705	710	715 720
Met Ile Thr Val Thr	Glu Ala Ser Thr Val	Phe Pro Gln Thr Phe Tyr
725	730	735
Leu Ser Gly Leu Phe Phe	Phe Gly Pro Val Asn	Leu Ser Ala Ser Thr
740	745	750
Leu Thr Val Pro Thr Ile	Thr Leu Thr Ile Gly	Gly Pro Thr Val Thr
755	760	765
Val Pro Ile Ser Ile Val	Gly Ala Leu Glu Ser	Arg Thr Ile Thr Phe
770	775	780
Leu Lys Ile Asp Pro Ala	Pro Gly Ile Gly Asn	Ser Thr Thr Asn Pro
785	790	795 800
Ser Ser Gly Phe Phe Asn	Ser Gly Thr Gly Gly	Thr Ser Gly Phe Gln
805	810	815
Asn Val Gly Gly Gly Ser	Ser Gly Val Trp Asn	Ser Gly Leu Ser Ser
820	825	830

216

Ala Ile Gly Asn Ser Gly Phe Gln Asn Leu Gly Ser Leu Gln Ser Gly
 835 840 845

Trp Ala Asn Leu Gly Asn Ser Val Ser Gly Phe Phe Asn Thr Ser Thr
 850 855 860

Val Asn Leu Ser Thr Pro Ala Asn Val Ser Gly Leu Asn Asn Ile Gly
 865 870 875 880

Thr Asn Leu Ser Gly Val Phe Arg Gly Pro Thr Gly Thr Ile Phe Asn
 885 890 895

Ala Gly Leu Ala Asn Leu Gly Gln Leu Asn Ile Gly Ser Ala Ser Cys
 900 905 910

Arg Ile Arg His Glu Leu Asp Thr Val Ser Thr Ile Ile Ser Ala Phe
 915 920 925

Cys Gly Ser Ala Ser Asp Glu Ser Asn Pro Gly Ser Val Ser Glu
 930 935 940

(2) INFORMATION FOR SEQ ID NO:200:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 53 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:200:

GGATCCATAT GGGCCATCAT CATCATCATC ACGTGATCGA CATCATCGGG ACC

53

(2) INFORMATION FOR SEQ ID NO:201:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 42 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:201:

CCTGAATTCA GGCCTCGGTT GCGCCGGCCT CATCTTGAAC GA

42

(2) INFORMATION FOR SEQ ID NO:202:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 31 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:202:

GGATCCTGCA GGCTCGAAAC CACCGAGCGG T

31

(2) INFORMATION FOR SEQ ID NO:203:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 31 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:203:

CTCTGAATTC AGCGCTGGAA ATCGTCGCGA T

31

(2) INFORMATION FOR SEQ ID NO:204:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 33 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:204:

GGATCCAGCG CTGAGATGAA GACCGATGCC GCT

33

(2) INFORMATION FOR SEQ ID NO:205:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 38 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:205:

GGATATCTGC AGAATTCAGG TTAAAGCCC ATTTGCGA

38

(2) INFORMATION FOR SEQ ID NO:206:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 30 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:206:

CCGCATGCGA GCCACGTGCC CACAACGGCC

30

(2) INFORMATION FOR SEQ ID NO:207:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 37 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:207:

CTTCATGGAA TTCTCAGGCC GGTAAGGTCC GCTGCGG

37

(2) INFORMATION FOR SEQ ID NO:208:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 7676 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:208:

TGGCGAATGG GACGCGCCCT GTAGCGGCGC ATTAAGCGCG GCGGGTGTGG TGGTTACGCG

60

CAGCGTGACC GCTACACTTG CCAGCGCCCT AGCGCCCGCT CCTTTCGCTT TCTTCCCTTC	120
CTTTCTCGCC ACGTTCGCCG GCTTTCCCCG TCAAGCTCTA AATCGGGGGC TCCCTTTAGG	180
GTTCCGATTT AGTGCTTTAC GGCACCTCGA CCCCCAAAAA CTTGATTAGG GTGATGGTTC	240
ACGTAGTGGG CCATCGCCCT GATAGACGGT TTTTCGCCCT TTGACGTTGG AGTCCACGTT	300
CTTTAATAGT GGACTCTTGT TCCAACTGG AACAACTC AACCCCTATCT CGGTCTATTC	360
TTTTGATTTA TAAGGGATTT TGCCGATTTT GGCCTATTGG TTAAAAAATG AGCTGATTTA	420
ACAAAAATTT AACGCGAATT TTAACAAAAT ATTAACGTTT ACAATTTTCA GTGGCACTTT	480
TCGGGGAAAT GTGCGCGGAA CCCCTATTTG TTTATTTTTC TAAATACATT CAAATATGTA	540
TCCGCTCATG AATTAATTCT TAGAAAACT CATCGAGCAT CAAATGAAAC TGCAATTTAT	600
TCATATCAGG ATTATCAATA CCATATTTTT GAAAAAGCCG TTTCTGTAAT GAAGGAGAAA	660
ACTCACCGAG GCAGTTCCAT AGGATGGCAA GATCCTGGTA TCGGTCTGCG ATTCCGACTC	720
GTCCAACATC AATACAACCT ATTAATTTCC CCTCGTCAAA AATAAGGTTA TCAAGTGAGA	780
AATCACCATG AGTGACGACT GAATCCGGTG AGAATGGCAA AAGTTTATGC ATTTCTTTCC	840
AGACTTGTTT AACAGGCCAG CCATTACGCT CGTCATCAAA ATCACTCGCA TCAACCAAAC	900
CGTTATTCAT TCGTGATTGC GCCTGAGCGA GACGAAATAC GCGATCGCTG TTAAAAGGAC	960
AATTACAAAC AGGAATCGAA TGCAACCGGC GCAGGAACAC TGCCAGCGCA TCAACAATAT	1020
TTTACCTGA ATCAGGATAT TCTTCTAATA CCTGGAATGC TGTTTTCCCG GGGATCGCAG	1080
TGGTGAGTAA CCATGCATCA TCAGGAGTAC GGATAAAATG CTTGATGGTC GGAAGAGGCA	1140
TAAATTCCGT CAGCCAGTTT AGTCTGACCA TCTCATCTGT AACATCATTG GCAACGCTAC	1200
CTTTGCCATG TTTGAGAAAC AACTCTGGCG CATCGGGCTT CCCATACAAT CGATAGATTG	1260
TCGCACCTGA TTGCCCAGCA TTATCGCGAG CCCATTTATA CCCATATAAA TCAGCATCCA	1320
TGTTGGAATT TAATCGCGGC CTAGAGCAAG ACGTTTCCCG TTGAATATGG CTCATAACAC	1380
CCCTTGATT ACTGTTTATG TAAGCAGACA GTTTTATTGT TCATGACCAA AATCCCTTAA	1440
CGTGAGTTTT CGTTCCACTG AGCGTCAGAC CCCGTAGAAA AGATCAAAGG ATCTTCTTGA	1500
GATCCTTTTT TTCTGCGCGT AATCTGCTGC TTGCAAACAA AAAAACCACC GCTACCAGCG	1560
GTGGTTTGTT TGCCGGATCA AGAGCTACCA ACTCTTTTTT CGAAGGTAAC TGGCTTCAGC	1620
AGAGCGCAGA TACCAAATAC TGTCTTCTA GTGTAGCCGT AGTTAGGCCA CCACTTCAAG	1680

AACTCTGTAG CACCGCCTAC ATACCTCGCT CTGCTAATCC TGTTACCAGT GGCTGCTGCC	1740
AGTGGCGATA AGTCGTGTCT TACCGGGTTG GACTCAAGAC GATAGTTACC GGATAAGGCG	1800
CAGCGGTCGG GCTGAACGGG GGGTTCGTGC ACACAGCCCA GCTTGGAGCG AACGACCTAC	1860
ACCGAACTGA GATACCTACA GCGTGAGCTA TGAGAAAGCG CCACGCTTCC CGAAGGGAGA	1920
AAGGCGGACA GGTATCCGGT AAGCGGCAGG GTCGGAACAG GAGAGCGCAC GAGGGAGCTT	1980
CCAGGGGGAA ACGCCTGGTA TCTTTATAGT CCTGTCGGGT TTCGCCACCT CTGACTTGAG	2040
CGTCGATTTT TGTGATGCTC GTCAGGGGGG CGGAGCCTAT GGAAAAACGC CAGCAACGCG	2100
GCCTTTTAC GGTTCCCTGGC CTTTTGCTGG CCTTTTGCTC ACATGTTCTT TCCTGCGTTA	2160
TCCCCTGATT CTGTGGATAA CCGTATTACC GCCTTTGAGT GAGCTGATAC CGCTCGCCGC	2220
AGCCGAACGA CCGAGCGCAG CGAGTCAGTG AGCGAGGAAG CGGAAGAGCG CCTGATGCGG	2280
TATTTTCTCC TTACGCATCT GTGCGGTATT TCACACCGCA TATATGGTGC ACTCTCAGTA	2340
CAATCTGCTC TGATGCCGCA TAGTTAAGCC AGTATACACT CCGCTATCGC TACGTGACTG	2400
GGTCATGGCT GCGCCCCGAC ACCCGCCAAC ACCCGCTGAC GCGCCCTGAC GGGCTTGTCT	2460
GCTCCCGGCA TCCGCTTACA GACAAGCTGT GACCGTCTCC GGGAGCTGCA TGTGTCAGAG	2520
GTTTTACCG TCATCACCGA AACGCGCGAG GCAGCTGCGG TAAAGCTCAT CAGCGTGGTC	2580
GTGAAGCGAT TCACAGATGT CTGCCTGTTC ATCCGCGTCC AGCTCGTTGA GTTCTCCAG	2640
AAGCGTTAAT GTCTGGCTTC TGATAAAGCG GGCCATGTTA AGGGCGGTTT TTTCCTGTTT	2700
GGTCACTGAT GCCTCCGTGT AAGGGGGATT TCTGTTTCATG GGGGTAATGA TACCGATGAA	2760
ACGAGAGAGG ATGCTCACGA TACGGGTTAC TGATGATGAA CATGCCCGGT TACTGGAACG	2820
TTGTGAGGGT AAACAACCTGG CGGTATGGAT GCGGCGGGAC CAGAGAAAAA TCACTCAGGG	2880
TCAATGCCAG CGCTTCGTTA ATACAGATGT AGGTGTTCCA CAGGGTAGCC AGCAGCATCC	2940
TGCGATGCAG ATCCGGAACA TAATGGTGCA GGGCGCTGAC TTCCGCGTTT CCAGACTTTA	3000
CGAAACACGG AAACCGAAGA CCATTCATGT TGTGCTCAG GTCGAGACG TTTTGACGCA	3060
GCAGTCGCTT CACGTTTCGCT CGCGTATCGG TGATTCATTC TGCTAACCAG TAAGGCAACC	3120
CCGCCAGCCT AGCCGGGTCC TCAACGACAG GAGCACGATC ATGCGCACCC GTGGGGCCGC	3180
CATGCCGGCG ATAATGGCCT GCTTCTCGCC GAAACGTTTG GTGGCGGGAC CAGTGACGAA	3240
GGCTTGAGCG AGGGCGTGCA AGATTCCGAA TACCGCAAGC GACAGGCCGA TCATCGTCGC	3300
GCTCCAGCGA AAGCGGTCCT CGCCGAAAAT GACCCAGAGC GCTGCCGGCA CCTGTCCTAC	3360

GAGTTGCATG ATAAAGAAGA CAGTCATAAG TCGGGCGACG ATAGTCATGC CCCGCGCCCA	3420
CCGGAAGGAG CTGACTGGGT TGAAGGCTCT CAAGGGCATC GGTCGAGATC CCGGTGCCTA	3480
ATGAGTGAGC TAACTTACAT TAATTGCGTT GCGCTCACTG CCCGCTTCC AGTCGGGAAA	3540
CCTGTCGTGC CAGCTGCATT AATGAATCGG CCAACGCGCG GGGAGAGGCG GTTTGCGTAT	3600
TGGGCGCCAG GGTGGTTTTT CTTTTCACCA GTGAGACGGG CAACAGCTGA TTGCCCTTCA	3660
CCGCCTGGCC CTGAGAGAGT TGCAGCAAGC GGTCCACGCT GGTTTGCCCC AGCAGGCGAA	3720
AATCCTGTTT GATGGTGGTT AACGGCGGGA TATAACATGA GCTGTCTTCG GTATCGTCGT	3780
ATCCCACTAC CGAGATATCC GCACCAACGC GCAGCCCGGA CTCGGTAATG GCGCGCATTG	3840
CGCCCAGCGC CATCTGATCG TTGGCAACCA GCATCGCAGT GGAACGATG CCCTCATTCA	3900
GCATTTGCAT GGTTTGTGA AAACCGGACA TGGCACTCCA GTCGCCTTCC CGTTCCGCTA	3960
TCGGCTGAAT TTGATTGCGA GTGAGATATT TATGCCAGCC AGCCAGACGC AGACGCGCCG	4020
AGACAGAACT TAATGGGCCC GCTAACAGCG CGATTTGCTG GTGACCCAAT GCGACCAGAT	4080
GCTCCACGCC CAGTCGCGTA CCGTCTTCAT GGGAGAAAAT AATACTGTTG ATGGGTGTCT	4140
GGTCAGAGAC ATCAAGAAAT AACGCCGGA CATTAGTGCA GGCAGCTTCC ACAGCAATGG	4200
CATCCTGGTC ATCCAGCGGA TAGTTAATGA TCAGCCCACT GACGCGTTGC GCGAGAAGAT	4260
TGTGCACCGC CGCTTTACAG GCTTCGACGC CGCTTCGTTT TACCATCGAC ACCACCACGC	4320
TGGCACCCAG TTGATCGGCG CGAGATTTAA TCGCCGCGAC AATTTGCGAC GGCGCGTGCA	4380
GGGCCAGACT GGAGGTGGCA ACGCCAATCA GCAACGACTG TTTGCCCGCC AGTTGTTGTG	4440
CCACGCGGTT GGAATGTAA TTCAGCTCCG CCATCGCCGC TTCCACTTTT TCCCGCGTTT	4500
TCGCAGAAAC GTGGCTGGCC TGGTTCACCA CGCGGGAAAC GGTCTGATAA GAGACACCGG	4560
CATACTCTGC GACATCGTAT AACGTTACTG GTTTCACATT CACCACCCTG AATTGACTCT	4620
CTTCCGGGCG CTATCATGCC ATACCGCGAA AGGTTTTGCG CCATTCGATG GTGTCCGGGA	4680
TCTCGACGCT CTCCCTTATG CGACTCCTGC ATTAGGAAGC AGCCAGTAG TAGGTTGAGG	4740
CCGTTGAGCA CCGCCGCCGC AAGGAATGGT GCATGCAAGG AGATGGCGCC CAACAGTCCC	4800
CCGGCCACGG GGCCTGCCAC CATACCACG CCGAAACAAG CGCTCATGAG CCCGAAGTGG	4860
CGAGCCCGAT CTTCCCCATC GGTGATGTG GCGATATAGG CGCCAGCAAC CGCACCTGTG	4920
GCGCCGGTGA TGCCGGCCAC GATGCGTCCG GCGTAGAGGA TCGAGATCTC GATCCCGCGA	4980

AATTAATACG	ACTCACTATA	GGGGAATTGT	GAGCGGATAA	CAATTCCCCT	CTAGAAATAA	5040
TTTTGTTTAA	CTTTAAGAAG	GAGATATACA	TATGGGCCAT	CATCATCATC	ATCACGTGAT	5100
CGACATCATC	GGGACCAGCC	CCACATCCTG	GGAACAGGCG	GCGGCGGAGG	CGGTCCAGCG	5160
GGCGCGGGAT	AGCGTCGATG	ACATCCGCGT	CGCTCGGGTC	ATTGAGCAGG	ACATGGCCGT	5220
GGACAGCGCC	GGCAAGATCA	CCTACCGCAT	CAAGCTCGAA	GTGTCGTTCA	AGATGAGGCC	5280
GGCGCAACCG	AGGGGCTCGA	AACCACCGAG	CGGTTCGCCT	GAAACGGGCG	CCGGCGCCGG	5340
TACTGTCGCG	ACTACCCCCG	CGTCGTCGCC	GGTGACGTTG	GCGGAGACCG	GTAGCACGCT	5400
GCTCTACCCG	CTGTTCAACC	TGTGGGGTCC	GGCCTTTTAC	GAGAGGTATC	CGAACGTCAC	5460
GATCACCGCT	CAGGGCACCG	GTTCTGGTGC	CGGGATCGCG	CAGGCCGCCG	CCGGGACGGT	5520
CAACATTGGG	GCCTCCGACG	CCTATCTGTC	GGAAGGTGAT	ATGGCCGCGC	ACAAGGGGCT	5580
GATGAACATC	GCGCTAGCCA	TCTCCGCTCA	GCAGGTCAAC	TACAACCTGC	CCGGAGTGAG	5640
CGAGCACCTC	AAGCTGAACG	GAAAAGTCCT	GGCGGCCATG	TACCAGGGCA	CCATCAAAAC	5700
CTGGGACGAC	CCGCAGATCG	CTGCGCTCAA	CCCCGGCGTG	AACCTGCCCC	GCACCGCGGT	5760
AGTTCCGCTG	CACCGCTCCG	ACGGGTCCGG	TGACACCTTC	TTGTTACCCC	AGTACCTGTC	5820
CAAGCAAGAT	CCCGAGGGCT	GGGGCAAGTC	GCCCGGCTTC	GGCACCACCG	TCGACTTCCC	5880
GGCGGTGCCG	GGTGCGCTGG	GTGAGAACGG	CAACGGCGGC	ATGGTGACCG	GTTGCGCCGA	5940
GACACCGGGC	TGCGTGCCCT	ATATCGGCAT	CAGCTTCCTC	GACCAGGCCA	GTCAACGGGG	6000
ACTCGGCGAG	GCCCAACTAG	GCAATAGCTC	TGGCAATTTT	TTGTTGCCCC	ACGCGCAAAG	6060
CATTACAGCC	GCGGCGGCTG	GCTTCGCATC	GAAAACCCCG	GCGAACCAGG	CGATTTTCGAT	6120
GATCGACGGG	CCCGCCCCGG	ACGGCTACCC	GATCATCAAC	TACGAGTACG	CCATCGTCAA	6180
CAACCGGCAA	AAGGACGCCG	CCACCGCGCA	GACCTTGCAG	GCATTTCTGC	ACTGGGCGAT	6240
CACCGACGGC	AACAAGGCCT	CGTTCCTCGA	CCAGGTTTCAT	TTCCAGCCGC	TGCCGCCCGC	6300
GGTGGTGAAG	TTGTCTGACG	CGTTGATCGC	GACGATTTCC	AGCGCTGAGA	TGAAGACCGA	6360
TGCCGCTACC	CTCGCGCAGG	AGGCAGGTAA	TTTCGAGCGG	ATCTCCGGCG	ACCTGAAAAC	6420
CCAGATCGAC	CAGGTGGAGT	CGACGGCAGG	TTCGTTGCAG	GGCCAGTGGC	GCGGCGCGGC	6480
GGGGACGGCC	GCCCAGGCCG	CGGTGGTGCG	CTTCCAAGAA	GCAGCCAATA	AGCAGAAGCA	6540
GGAACCTCGAC	GAGATCTCGA	CGAATATTCG	TCAGGCCGGC	GTCCAATACT	CGAGGGCCGA	6600
CGAGGAGCAG	CAGCAGGCGC	TGTCCTCGCA	AATGGGCTTT	GTGCCACAA	CGGCCGCCTC	6660

GCCGCCGTCG ACCGCTGCAG CGCCACCCGC ACCGGCGACA CCTGTTGCCC CCCCACCACC 6720
 GGCCGCCGCC AACACGCCGA ATGCCAGCC GGGCGATCCC AACGCAGCAC CTCCGCCGGC 6780
 CGACCCGAAC GCACCGCCGC CACCTGTCAT TGCCCCAAC GCACCCCAAC CTGTCCGGAT 6840
 CGACAACCCG GTTGGAGGAT TCAGCTTCGC GCTGCCTGCT GGCTGGGTGG AGTCTGACGC 6900
 CGCCCACTTC GACTACGGTT CAGCACTCCT CAGCAAAACC ACCGGGGACC CGCCATTTCC 6960
 CGGACAGCCG CCGCCGGTGG CCAATGACAC CCGTATCGTG CTCGGCCGGC TAGACCAAAA 7020
 GCTTTACGCC AGCGCCGAAG CCACCGACTC CAAGGCCGCG GCCCGGTTGG GCTCGGACAT 7080
 GGGTGAGTTC TATATGCCCT ACCCGGGCAC CCGGATCAAC CAGGAAACCG TCTCGCTTGA 7140
 CGCCAACGGG GTGTCTGGAA GCGCGTCGTA TTACGAAGTC AAGTTCAGCG ATCCGAGTAA 7200
 GCCGAACGGC CAGATCTGGA CGGGCGTAAT CGGCTCGCCC GCGGCGAACG CACCGGACGC 7260
 CGGGCCCCCT CAGCGCTGGT TTGTGGTATG GCTCGGGACC GCCAACAACC CGGTGGACAA 7320
 GGGCGCGGCC AAGGCGCTGG CCGAATCGAT CCGGCCTTTG GTCGCCCCGC CGCCGGCGCC 7380
 GGCACCGGCT CCTGCAGAGC CCGCTCCGGC GCCGGCGCCG GCCGGGGAAG TCGCTCCTAC 7440
 CCCGACGACA CCGACACCGC AGCGGACCTT ACCGGCCTGA GAATTCTGCA GATATCCATC 7500
 AACTGGCGG CCGCTCGAGC ACCACCACCA CCACCACTGA GATCCGGCTG CTAACAAAGC 7560
 CCGAAAGGAA GCTGAGTTGG CTGCTGCCAC CGCTGAGCAA TAACTAGCAT AACCCCTTGG 7620
 GGCCTCTAAA CGGGTCTTGA GGGGTTTTTT GCTGAAAGGA GGAACCTATAT CCGGAT 7676

(2) INFORMATION FOR SEQ ID NO:209:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 802 amino acids
- (B) TYPE: amino acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:209:

Met Gly His His His His His His Val Ile Asp Ile Ile Gly Thr Ser
 1 5 10 15
 Pro Thr Ser Trp Glu Gln Ala Ala Ala Glu Ala Val Gln Arg Ala Arg
 20 25 30

224

Asp Ser Val Asp Asp Ile Arg Val Ala Arg Val Ile Glu Gln Asp Met
 35 40 45
 Ala Val Asp Ser Ala Gly Lys Ile Thr Tyr Arg Ile Lys Leu Glu Val
 50 55 60
 Ser Phe Lys Met Arg Pro Ala Gln Pro Arg Gly Ser Lys Pro Pro Ser
 65 70 75 80
 Gly Ser Pro Glu Thr Gly Ala Gly Ala Gly Thr Val Ala Thr Thr Pro
 85 90 95
 Ala Ser Ser Pro Val Thr Leu Ala Glu Thr Gly Ser Thr Leu Leu Tyr
 100 105 110
 Pro Leu Phe Asn Leu Trp Gly Pro Ala Phe His Glu Arg Tyr Pro Asn
 115 120 125
 Val Thr Ile Thr Ala Gln Gly Thr Gly Ser Gly Ala Gly Ile Ala Gln
 130 135 140
 Ala Ala Ala Gly Thr Val Asn Ile Gly Ala Ser Asp Ala Tyr Leu Ser
 145 150 155 160
 Glu Gly Asp Met Ala Ala His Lys Gly Leu Met Asn Ile Ala Leu Ala
 165 170 175
 Ile Ser Ala Gln Gln Val Asn Tyr Asn Leu Pro Gly Val Ser Glu His
 180 185 190
 Leu Lys Leu Asn Gly Lys Val Leu Ala Ala Met Tyr Gln Gly Thr Ile
 195 200 205
 Lys Thr Trp Asp Asp Pro Gln Ile Ala Ala Leu Asn Pro Gly Val Asn
 210 215 220
 Leu Pro Gly Thr Ala Val Val Pro Leu His Arg Ser Asp Gly Ser Gly
 225 230 235 240
 Asp Thr Phe Leu Phe Thr Gln Tyr Leu Ser Lys Gln Asp Pro Glu Gly
 245 250 255
 Trp Gly Lys Ser Pro Gly Phe Gly Thr Thr Val Asp Phe Pro Ala Val
 260 265 270
 Pro Gly Ala Leu Gly Glu Asn Gly Asn Gly Gly Met Val Thr Gly Cys
 275 280 285
 Ala Glu Thr Pro Gly Cys Val Ala Tyr Ile Gly Ile Ser Phe Leu Asp
 290 295 300
 Gln Ala Ser Gln Arg Gly Leu Gly Glu Ala Gln Leu Gly Asn Ser Ser
 305 310 315 320
 Gly Asn Phe Leu Leu Pro Asp Ala Gln Ser Ile Gln Ala Ala Ala Ala

[illegible]

Gly Asp Pro Pro Phe Pro Gly Gln Pro Pro Pro Val Ala Asn Asp Thr
625 630 635 640

Arg Ile Val Leu Gly Arg Leu Asp Gln Lys Leu Tyr Ala Ser Ala Glu
645 650 655

Ala Thr Asp Ser Lys Ala Ala Ala Arg Leu Gly Ser Asp Met Gly Glu
660 665 670

Phe Tyr Met Pro Tyr Pro Gly Thr Arg Ile Asn Gln Glu Thr Val Ser
675 680 685

Leu Asp Ala Asn Gly Val Ser Gly Ser Ala Ser Tyr Tyr Glu Val Lys
690 695 700

Phe Ser Asp Pro Ser Lys Pro Asn Gly Gln Ile Trp Thr Gly Val Ile
705 710 715 720

Gly Ser Pro Ala Ala Asn Ala Pro Asp Ala Gly Pro Pro Gln Arg Trp
725 730 735

Phe Val Val Trp Leu Gly Thr Ala Asn Asn Pro Val Asp Lys Gly Ala
740 745 750

Ala Lys Ala Leu Ala Glu Ser Ile Arg Pro Leu Val Ala Pro Pro Pro
755 760 765

Ala Pro Ala Pro Ala Pro Ala Glu Pro Ala Pro Ala Pro Ala Pro Ala
770 775 780

Gly Glu Val Ala Pro Thr Pro Thr Thr Pro Thr Pro Gln Arg Thr Leu
785 790 795 800

Pro Ala

CLAIMS

We claim:

1. A polypeptide comprising an antigenic portion of a soluble *M. tuberculosis* antigen, or a variant of said antigen that differs only in conservative substitutions and/or modifications, wherein said antigen has an N-terminal sequence selected from the group consisting of:

- (a) Asp-Pro-Val-Asp-Ala-Val-Ile-Asn-Thr-Thr-Cys-Asn-Tyr-Gly-Gln-Val-Val-Ala-Ala-Leu (SEQ ID NO: 115);
- (b) Ala-Val-Glu-Ser-Gly-Met-Leu-Ala-Leu-Gly-Thr-Pro-Ala-Pro-Ser (SEQ ID NO: 116);
- (c) Ala-Ala-Met-Lys-Pro-Arg-Thr-Gly-Asp-Gly-Pro-Leu-Glu-Ala-Ala-Lys-Glu-Gly-Arg (SEQ ID NO: 17);
- (d) Tyr-Tyr-Trp-Cys-Pro-Gly-Gln-Pro-Phe-Asp-Pro-Ala-Trp-Gly-Pro (SEQ ID NO: 118);
- (e) Asp-Ile-Gly-Ser-Glu-Ser-Thr-Glu-Asp-Gln-Gln-Xaa-Ala-Val (SEQ ID NO: 119);
- (f) Ala-Glu-Glu-Ser-Ile-Ser-Thr-Xaa-Glu-Xaa-Ile-Val-Pro (SEQ ID NO: 120);
- (g) Asp-Pro-Glu-Pro-Ala-Pro-Pro-Val-Pro-Thr-Thr-Ala-Ala-Ser-Pro-Pro-Ser (SEQ ID NO: 121);
- (h) Ala-Pro-Lys-Thr-Tyr-Xaa-Glu-Glu-Leu-Lys-Gly-Thr-Asp-Thr-Gly (SEQ ID NO: 122);
- (i) Asp-Pro-Ala-Ser-Ala-Pro-Asp-Val-Pro-Thr-Ala-Ala-Gln-Leu-Thr-Ser-Leu-Leu-Asn-Ser-Leu-Ala-Asp-Pro-Asn-Val-Ser-Phe-Ala-Asn (SEQ ID NO: 123); and
- (j) Ala-Pro-Glu-Ser-Gly-Ala-Gly-Leu-Gly-Gly-Thr-Val-Gln-Ala-Gly; (SEQ ID NO: 131)

wherein Xaa may be any amino acid.

2. A polypeptide comprising an immunogenic portion of an *M. tuberculosis* antigen, or a variant of said antigen that differs only in conservative substitutions and/or modifications, wherein said antigen has an N-terminal sequence selected from the group consisting of:

- (a) Asp-Pro-Pro-Asp-Pro-His-Gln-Xaa-Asp-Met-Thr-Lys-Gly-Tyr-Tyr-Pro-Gly-Gly-Arg-Arg-Xaa-Phe; (SEQ ID NO: 124) and
- (b) Xaa-Tyr-Ile-Ala-Tyr-Xaa-Thr-Thr-Ala-Gly-Ile-Val-Pro-Gly-Lys-Ile-Asn-Val-His-Leu-Val; (SEQ ID NO: 132), wherein Xaa may be any amino acid.

3. A polypeptide comprising an antigenic portion of a soluble *M. tuberculosis* antigen, or a variant of said antigen that differs only in conservative substitutions and/or modifications, wherein said antigen comprises an amino acid sequence encoded by a DNA sequence selected from the group consisting of the sequences recited in SEQ ID NOS: 1, 2, 4-10, 13-25, 52, 94 and 96, the complements of said sequences, and DNA sequences that hybridize to a sequence recited in SEQ ID NOS: 1, 2, 4-10, 13-25, 52, 94 and 96 or a complement thereof under moderately stringent conditions.

4. A polypeptide comprising an antigenic portion of a *M. tuberculosis* antigen, or a variant of said antigen that differs only in conservative substitutions and/or modifications, wherein said antigen comprises an amino acid sequence encoded by a DNA sequence selected from the group consisting of the sequences recited in SEQ ID NOS: 26-51, 133, 134, 158-178 and 196, the complements of said sequences, and DNA sequences that hybridize to a sequence recited in SEQ ID NOS: 26-51, 133, 134, 158-178 and 196 or a complement thereof under moderately stringent conditions.

5. A DNA molecule comprising a nucleotide sequence encoding a polypeptide according to any one of claims 1-4.

6. A recombinant expression vector comprising a DNA molecule according to claim 5.

7. A host cell transformed with an expression vector according to claim 6.

8. The host cell of claim 7 wherein the host cell is selected from the group consisting of *E. coli*, yeast and mammalian cells.

9. A method for detecting *M. tuberculosis* infection in a biological sample, comprising:

(a) contacting a biological sample with one or more polypeptides according to any of claims 1-4; and

(b) detecting in the sample the presence of antibodies that bind to at least one of the polypeptides, thereby detecting *M. tuberculosis* infection in the biological sample.

10. A method for detecting *M. tuberculosis* infection in a biological sample, comprising:

(a) contacting a biological sample with a polypeptide having an N-terminal sequence selected from the group consisting of sequences provided in SEQ ID NO: 129 and 130; and

(b) detecting in the sample the presence of antibodies that bind to at least one of the polypeptides, thereby detecting *M. tuberculosis* infection in the biological sample.

11. A method for detecting *M. tuberculosis* infection in a biological sample, comprising:

(a) contacting a biological sample with one or more polypeptides encoded by a DNA sequence selected from the group consisting of SEQ ID NOS: 3, 11, 12, 135, 136, 151-155, 184-188, 194-195 and 198, the complements of said sequences, and DNA sequences that hybridize to a sequence recited in SEQ ID NOS: 3, 11, 12, 135, 136, 151-155, 184-188, 194-195 and 198; and

(b) detecting in the sample the presence of antibodies that bind to at least one of the polypeptides, thereby detecting *M. tuberculosis* infection in the biological sample.

12. The method of any one of claims 9-11 wherein step (a) additionally comprises contacting the biological sample with a 38 kD *M. tuberculosis* antigen and step (b) additionally comprises detecting in the sample the presence of antibodies that bind to the 38 kD *M. tuberculosis* antigen.

13. The method of any one of claims 9-11 wherein the polypeptide(s) are bound to a solid support.

14. The method of claim 13 wherein the solid support comprises nitrocellulose, latex or a plastic material.

15. The method of any one of claims 9-11 wherein the biological sample is selected from the group consisting of whole blood, serum, plasma, saliva, cerebrospinal fluid and urine.

16. The method of claim 15 wherein the biological sample is whole blood or serum.

17. A method for detecting *M. tuberculosis* infection in a biological sample, comprising:

(a) contacting the sample with at least two oligonucleotide primers in a polymerase chain reaction, wherein at least one of the oligonucleotide primers is specific for a DNA molecule according to claim 5; and

(b) detecting in the sample a DNA sequence that amplifies in the presence of the oligonucleotide primers, thereby detecting *M. tuberculosis* infection.

18. The method of claim 17, wherein at least one of the oligonucleotide primers comprises at least about 10 contiguous nucleotides of a DNA molecule according to claim 5.

19. A method for detecting *M. tuberculosis* infection in a biological sample, comprising:

(a) contacting the sample with at least two oligonucleotide primers in a polymerase chain reaction, wherein at least one of the oligonucleotide primers is specific for a DNA sequence selected from the group consisting of SEQ ID NOS: 3, 11, 12, 135, 136, 151-155, 184-188, 194-195 and 198; and

(b) detecting in the sample a DNA sequence that amplifies in the presence of the first and second oligonucleotide primers, thereby detecting *M. tuberculosis* infection.

20. The method of claim 19, wherein at least one of the oligonucleotide primers comprises at least about 10 contiguous nucleotides of a DNA sequence selected from the group consisting of SEQ ID NOS: 3, 11, 12, 135, 136, 151-155, 184-188, 194-195 and 198.

21. The method of claims 17 or 19 wherein the biological sample is selected from the group consisting of whole blood, sputum, serum, plasma, saliva, cerebrospinal fluid and urine.

22. A method for detecting *M. tuberculosis* infection in a biological sample, comprising:

(a) contacting the sample with one or more oligonucleotide probes specific for a DNA molecule according to claim 5; and

(b) detecting in the sample a DNA sequence that hybridizes to the oligonucleotide probe, thereby detecting *M. tuberculosis* infection.

23. The method of claim 22 wherein the probe comprises at least about 15 contiguous nucleotides of a DNA molecule according to claim 5.

24. A method for detecting *M. tuberculosis* infection in a biological sample, comprising:

(a) contacting the sample with one or more oligonucleotide probes specific for a DNA sequence selected from the group consisting of SEQ ID NOS: 3, 11, 12, 135, 136, 151-155, 184-188, 194-195 and 198; and

(b) detecting in the sample a DNA sequence that hybridizes to the oligonucleotide probe, thereby detecting *M. tuberculosis* infection.

25. The method of claim 24 wherein the oligonucleotide probe comprises at least about 15 contiguous nucleotides of a DNA sequence selected from the group consisting of SEQ ID NOS: 3, 11, 12, 135, 136, 151-155, 184-188, 194-195 and 198.

26. The method of claims 22 or 24 wherein the biological sample is selected from the group consisting of whole blood, sputum, serum, plasma, saliva, cerebrospinal fluid and urine.

27. A method for detecting *M. tuberculosis* infection in a biological sample, comprising:

(a) contacting the biological sample with a binding agent which is capable of binding to a polypeptide according to any one of claims 1-4; and

(b) detecting in the sample a protein or polypeptide that binds to the binding agent, thereby detecting *M. tuberculosis* infection in the biological sample.

28. A method for detecting *M. tuberculosis* infection in a biological sample, comprising:

(a) contacting the biological sample with a binding agent which is capable of binding to a polypeptide having an N-terminal sequence selected from the group consisting of sequences provided in SEQ ID NO: 129 and 130; and

(b) detecting in the sample a protein or polypeptide that binds to the binding agent, thereby detecting *M. tuberculosis* infection in the biological sample.

29. A method for detecting *M. tuberculosis* infection in a biological sample, comprising:

(a) contacting the biological sample with a binding agent which is capable of binding to a polypeptide encoded by a DNA sequence selected from the group consisting of SEQ ID NOS: 3, 11, 12, 135, 136, 151-155, 184-188, 194-195 and 198, the complements of said sequences, and DNA sequences that hybridize to a sequence recited in SEQ ID NOS: 3, 11, 12, 135, 136, 151-155, 184-188, 194-195 and 198; and

(b) detecting in the sample a protein or polypeptide that binds to the binding agent, thereby detecting *M. tuberculosis* infection in the biological sample.

30. The method of any one of claims 27-29 wherein the binding agent is a monoclonal antibody.

31. The method of any one of claims 27-29 wherein the binding agent is a polyclonal antibody.

32. A diagnostic kit comprising:

- (a) one or more polypeptides according to any of claims 1-4; and
- (b) a detection reagent.

33. A diagnostic kit comprising:

- (a) one or more polypeptides having an N-terminal sequence selected from the group consisting of sequences provided in SEQ ID NO: 129 and 130; and
- (b) a detection reagent.

34. A diagnostic kit comprising:
- (a) one or more polypeptides encoded by a DNA sequence selected from the group consisting of SEQ ID NOS: 3, 11, 12, 135, 136, 151-155, 184-188, 194-195 and 198, the complements of said sequences, and DNA sequences that hybridize to a sequence recited in SEQ ID NOS: 3, 11, 12, 135, 136, 151-155, 184-188, 194-195 and 198; and
 - (b) a detection reagent.
35. The kit of any one of claims 32-34 wherein the polypeptide(s) are immobilized on a solid support.
36. The kit of claim 35 wherein the solid support comprises nitrocellulose, latex or a plastic material.
37. The kit of any one of claims 32-34 wherein the detection reagent comprises a reporter group conjugated to a binding agent.
38. The kit of claim 37 wherein the binding agent is selected from the group consisting of anti-immunoglobulins, Protein G, Protein A and lectins.
39. The kit of claim 37 wherein the reporter group is selected from the group consisting of radioisotopes, fluorescent groups, luminescent groups, enzymes, biotin and dye particles.
40. A diagnostic kit comprising at least two oligonucleotide primers, at least one of the oligonucleotide primers being specific for a DNA molecule according to claim 5.

41. A diagnostic kit according to claim 40, wherein at least one of the oligonucleotide primers comprises at least about 10 contiguous nucleotide of a DNA molecule according to claim 5.

42. A diagnostic kit comprising a at least two oligonucleotide primers, at least one of the primers being specific for a DNA sequence selected from the group consisting of SEQ ID NOS: 3, 11, 12, 135, 136, 151-155, 184-188, 194-195 and 198.

43. A diagnostic kit according to claim 42, wherein at least one of the oligonucleotide primers comprises at least about 10 contiguous nucleotide of a DNA sequence selected from the group consisting of SEQ ID NOS: 3, 11, 12, 135, 136, 151-155, 184-188, 194-195 and 198.

44. A diagnostic kit comprising at least one oligonucleotide probe, the oligonucleotide probe being specific for a DNA molecule according to claim 5.

45. A kit according to claim 44, wherein the oligonucleotide probe comprises at least about 15 contiguous nucleotides of a DNA molecule according to claim 5.



46. A diagnostic kit comprising at least one oligonucleotide probe, the oligonucleotide probe being specific for a DNA sequence selected from the group consisting of SEQ ID NOS: 3, 11, 12, 135, 136, 151-155, 184-188, 194-195 and 198.

47. A kit according to claim 46, wherein the oligonucleotide probe comprises at least about 15 contiguous nucleotides of a DNA sequence selected from the group consisting of SEQ ID NOS: 3, 11, 12, 135, 136, 151-155, 184-188, 194-195 and 198.



48. A monoclonal antibody that binds to a polypeptide according to any of claims 1-4.

49. A polyclonal antibody that binds to a polypeptide according to any of claims 1-4.
50. A fusion protein comprising two or more polypeptides according to any one of claims 1-4.
51. A fusion protein comprising one or more polypeptides according to any one of claims 1-4 and ESAT-6 (SEQ ID NO: 99).
52. A fusion protein comprising a polypeptide having an N-terminal sequence selected from the group of sequences provided in SEQ ID NOS: 129 and 130.
53. A fusion protein comprising one or more polypeptides according to any one of claims 1-4 and the *M. tuberculosis* antigen 38 kD (SEQ ID NO: 150).
54. A diagnostic kit comprising:
- (a) one or more fusion proteins according to any one of claims 50-53; and
 - (b) a detection reagent.

1/13

*Fig. 1A-1**Fig. 1A-2*

2/13

*Fig. 1B-1**Fig. 1B-2*

M 1 2 3 4 5
68-
43-
29-
18-
14-

I
II
Fig. 2B

M 1 2 3 4 5
97-
68-
43-
29-
18-
14-

I
II
Fig. 2D

M 1 2 3 4 5
97-
68-
43-
29-
18-
14-

I
II
Fig. 2A

M 1 2 3 4 5
97-
68-
43-
29-
18-
14-

I
II
Fig. 2C

4/13





Fig. 3A

5/13

*Fig. 3B*

6/13




Fig. 4

7/13




Fig. 5

8/13




Fig. 6

9/13




Fig. 7

10/13




Fig. 8

SUBSTITUTE SHEET (RULE 26)

11/13




Fig. 9

12/13




Fig. 10





Fig. 11



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶ : C12N 15/31, C07K 14/35, 16/12, C12Q 1/68, C12N 15/62, G01N 33/53		A3	(11) International Publication Number: WO 98/16645
(21) International Application Number: PCT/US97/18214		(43) International Publication Date: 23 April 1998 (23.04.98)	
(22) International Filing Date: 7 October 1997 (07.10.97)		(74) Agents: MAKI, David, J. et al.; Seed and Berry LLP, 6300 Columbia Center, 701 Fifth Avenue, Seattle, WA 98104-7092 (US).	
(30) Priority Data: 08/729,622 11 October 1996 (11.10.96) US 08/818,111 13 March 1997 (13.03.97) US		(81) Designated States: AL, AM, AT, AU, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GE, GH, HU, ID, IL, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, UZ, VN, YU, ARIPO patent (GH, KE, LS, MW, SD, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG).	
(71) Applicant: CORIXA CORPORATION [US/US]; 1124 Columbia Street, Seattle, WA 98104 (US).		<p>Published <i>With international search report.</i> <i>Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.</i></p>	
(72) Inventors: REED, Steven, G.; 2843 - 122nd Place N.E., Bellevue, WA 98005 (US). SKEIKY, Yasir, A., W.; 8327 - 25th Avenue N.W., Seattle, WA 98107 (US). DILLON, Davin, C.; 21607 N.E. 24th Street, Redmond, WA 98053 (US). CAMPOS-NETO, Antonio; 9308 Midship Court N.E., Bainbridge Island, WA 98021 (US). HOUGHTON, Raymond; 2636 - 242nd Place S.E., Bothell, WA 98021 (US). VEDVICK, Thomas, S.; 124 South 300th Place, Federal Way, WA 98003 (US). TWARDZIK, Daniel, R.; 10195 South Beach Drive, Bainbridge Island, WA 98110 (US). LODES, Michael, J.; 9223 - 36th Avenue S.W., Seattle, WA 98126 (US).			
(88) Date of publication of the international search report: 6 August 1998 (06.08.98)			

(54) Title: COMPOUNDS AND METHODS FOR DIAGNOSIS OF TUBERCULOSIS



(57) Abstract

Compounds and methods for diagnosing tuberculosis are disclosed. The compounds provided include polypeptides that contain at least one antigenic portion of one or more *M. tuberculosis* proteins, and DNA sequences encoding such polypeptides. Diagnostic kits containing such polypeptides or DNA sequences and a suitable detection reagent may be used for the detection of *M. tuberculosis* infection in patients and biological samples. Antibodies directed against such polypeptides are also provided.

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece			TR	Turkey
BG	Bulgaria	HU	Hungary	ML	Mali	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MN	Mongolia	UA	Ukraine
BR	Brazil	IL	Israel	MR	Mauritania	UG	Uganda
BY	Belarus	IS	Iceland	MW	Malawi	US	United States of America
CA	Canada	IT	Italy	MX	Mexico	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NE	Niger	VN	Viet Nam
CG	Congo	KE	Kenya	NL	Netherlands	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NO	Norway	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	NZ	New Zealand		
CM	Cameroon			PL	Poland		
CN	China	KR	Republic of Korea	PT	Portugal		
CU	Cuba	KZ	Kazakhstan	RO	Romania		
CZ	Czech Republic	LC	Saint Lucia	RU	Russian Federation		
DE	Germany	LI	Liechtenstein	SD	Sudan		
DK	Denmark	LK	Sri Lanka	SE	Sweden		
EE	Estonia	LR	Liberia	SG	Singapore		

INTERNATIONAL SEARCH REPORT

International Application No
PCT/US 97/18214

A. CLASSIFICATION OF SUBJECT MATTER IPC 6 C12N15/31 C07K14/35 C07K16/12 C12Q1/68 C12N15/62 G01N33/53		
According to International Patent Classification (IPC) or to both national classification and IPC		
B. FIELDS SEARCHED Minimum documentation searched (classification system followed by classification symbols) IPC 6 C12N C07K C12Q G01N		
Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched		
Electronic data base consulted during the international search (name of data base and, where practical, search terms used)		
C. DOCUMENTS CONSIDERED TO BE RELEVANT		
Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	EP 0 419 355 A (INNOGENETICS NV) 27 March 1991 see abstract see page 24, line 45 - page 26, line 19 see page 56 - page 72; claims ---	1,3,5-9, 13-18, 21-23, 26,27, 30-32, 35-41, 44,45, 48,49
A	WO 95 01441 A (STATENS SERUMSINSTITUT ;ANDERSEN PETER (DK); ANDERSEN AASE BENGGAAR) 12 January 1995 see abstract see page 20, line 13 - page 25, line 16 see page 73; claim 30 ---	50,51,54
-/--		
<input checked="" type="checkbox"/> Further documents are listed in the continuation of box C.		
<input checked="" type="checkbox"/> Patent family members are listed in annex.		
* Special categories of cited documents : "A" document defining the general state of the art which is not considered to be of particular relevance "E" earlier document but published on or after the international filing date "L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified) "O" document referring to an oral disclosure, use, exhibition or other means "P" document published prior to the international filing date but later than the priority date claimed "T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention "X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone "Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art. "&" document member of the same patent family		
Date of the actual completion of the international search 5 March 1998		Date of mailing of the international search report 23.06.98
Name and mailing address of the ISA European Patent Office, P.B. 5818 Patentlaan 2 NL - 2280 HV Rijswijk Tel. (+31-70) 340-2040, Tx. 31 651 epo nl, Fax: (+31-70) 340-3016		Authorized officer Macchia, G

INTERNATIONAL SEARCH REPORT

Intern. Application No

PCT/US 97/18214

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	WO 95 01440 A (STATENS SERUMINSTITUT ;HASLOEV KAARE (DK); ANDERSEN AASE BENGAARD) 12 January 1995 ---	
A	ANDERSEN P. ET AL.: "Identification of immunodominant antigens during infection with Mycobacterium tuberculosis" SCANDINAVIAN JOURNAL OF IMMUNOLOGY, vol. 36, 1992, pages 823-831, XP002057751 ---	
A	ANDERSEN A B ET AL: "STRUCTURE AND MAPPING OF ANTIGENIC DOMAINS OF PROTEIN ANTIGEN B, A 38,000-MOLECULAR-WEIGHT PROTEIN OF MYCOBACTERIUM TUBERCULOSIS" INFECTION AND IMMUNITY, vol. 57, no. 8, August 1989, pages 2481-2488, XP002026677 cited in the application see the whole document ---	12,53
A	WO 96 23885 A (PASTEUR INSTITUT ;LAQUEYRERIE ANNE (FR); MARCHAL GILLES (FR); PESCE) 8 August 1996 ---	
A	WO 92 21758 A (PASTEUR INSTITUT) 10 December 1992 ---	
A	AUSUBEL ET AL: "ISOLATION OF PROTEINS FOR MICROSEQUENCE ANALYSIS" CURRENT PROTOCOLS IN MOLECULAR BIOLOGY, 1993, pages 10.19.01-10.19.12, XP002026411 cited in the application ---	
A	YOUNG D B ET AL: "SCREENING OF A RECOMBINANT MYCOBACTERIAL DNA LIBRARY WITH POLYCLONAL ANTISERUM AND MOLECULAR WEIGHT ANALYSIS OF EXPRESSED ANTIGENS" INFECTION AND IMMUNITY, vol. 55, no. 6, June 1987, pages 1421-1425, XP002026410 ---	
A	WO 94 00493 A (KAPOOR ARCHANA ;MUNSHI ANIL (US)) 6 January 1994 ---	
A	FR 2 265 402 A (MITSUI PHARMACEUTICALS) 24 October 1975 ---	
A	FR 2 244 539 A (MITSUI PHARMACEUTICALS) 18 April 1975 ---	
	--- -/--	

15

INTERNATIONAL SEARCH REPORT

International Application No
PCT/US 97/18214

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	ROMAIN ET AL: "PREPARATION OF TUBERCULIN ANTIGEN L" ANNALES DE L'INSTITUT PASTEUR / MICROBIOLOGIE, vol. 136B, 1985, pages 235-248, XP002026409 ---	
P,X	WO 97 09429 A (CORIXA CORP) 13 March 1997 see abstract see page 173-181; claims ---	1,3,5-9, 12-18, 21-23, 26,27, 30-32, 35-41, 44,45, 48-51
P,X	WO 97 09428 A (CORIXA CORP) 13 March 1997 see abstract see page 158 - page 163; claims -----	// 1,3,5-8

INTERNATIONAL SEARCH REPORT

International application No.

PCT/US 97/ 18214

Box I Observations where certain claims were found unsearchable (Continuation of item 1 of first sheet)

This International Search Report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☐ Claims Nos.:
because they relate to subject matter not required to be searched by this Authority, namely:
2. ☐ Claims Nos.:
because they relate to parts of the International Application that do not comply with the prescribed requirements to such an extent that no meaningful International Search can be carried out, specifically:
3. ☐ Claims Nos.:
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

Box II Observations where unity of invention is lacking (Continuation of item 2 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

see continuation-sheet

1. ☐ As all required additional search fees were timely paid by the applicant, this International Search Report covers all searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this International Search Report covers only those claims for which fees were paid, specifically claims Nos.:
4. ☒ No required additional search fees were timely paid by the applicant. Consequently, this International Search Report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

1, 3, 5-9, 12-18, 21-23, 26, 27, 30-32, 35-41, 44, 45, 48-51, 53, 54 all partially
(subject 1. on next sheet)

Remark on Protest

- ☐ The additional search fees were accompanied by the applicant's protest.
- ☐ No protest accompanied the payment of additional search fees.

FURTHER INFORMATION CONTINUED FROM PCT/ISA/ 210

1. Claims: 1, 3, 5-9, 12-18, 21-23, 26, 27, 30-32, 35-41, 44, 45, 48-51, 53, 54 all partially.

A polypeptide comprising an antigenic portion of a soluble M. tuberculosis antigen or a variant, having an N-terminal aminoacid sequence as in Seq.ID:115 and/or encoded by a DNA molecule as in Seq.ID:96, complements of said sequence or sequences hybridizing to it. A DNA molecule comprising a sequence encoding said polypeptide. An expression vector comprising said DNA molecule, a host cell transformed with said expression vector. A method for detecting M. tuberculosis infection in a biological sample by detection of antibodies binding to said polypeptide or by detection of said polypeptide. A method for detecting M. tuberculosis infection in a biological sample by detection of said DNA sequence. Diagnostic kits thereof. An antibody binding to said polypeptide. A fusion protein comprising said polypeptide. Diagnostic kit comprising said fusion protein.

2. Claims: 1, 5-9, 12-18, 21-23, 26, 27, 30-32, 35-41, 44, 45, 48-51, 53, 54 all partially.

Same as invention 1 but for Seq.ID:116.

3. Claims: 1, 3, 5-9, 12-18, 21-23, 26, 27, 30-32, 35-41, 44, 45, 48-51, 53, 54 all partially.

Same as invention 1 but for Seq.ID:(1)17 and 25.

4. Claims: 1, 3, 5-9, 12-18, 21-23, 26, 27, 30-32, 35-41, 44, 45, 48-51, 53, 54 all partially.

Same as invention 1 but for Seq.ID:118 and 24.

5. Claims: 1, 5-9, 12-18, 21-23, 26, 27, 30-32, 35-41, 44, 45, 48-51, 53, 54 all partially.

Same as invention 1 but for Seq.ID:119.

6. Claims: 1, 5-9, 12-18, 21-23, 26, 27, 30-32, 35-41, 44, 45, 48-51, 53, 54 all partially.

Same as invention 1 but for Seq.ID:120.

FURTHER INFORMATION CONTINUED FROM PCT/ISA/ 210

7. Claims: 1, 3, 5-9, 12-18, 21-23, 26, 27, 30-32, 35-41, 44,
45, 48-51, 53, 54 all partially.

Same as invention 1 but for Seq.ID:121 and 52.

8. Claims: 1, 5-9, 12-18, 21-23, 26, 27, 30-32, 35-41, 44, 45,
48-51, 53, 54 all partially.

Same as invention 1 but for Seq.ID:122.

9. Claims: 1, 3, 5-9, 12-18, 21-23, 26, 27, 30-32, 35-41, 44,
45, 48-51, 53, 54 all partially.

Same as invention 1 but for Seq.ID:123 and 94.

10. Claims: 1, 5-9, 12-18, 21-23, 26, 27, 30-32, 35-41, 44, 45,
48-51, 53, 54 all partially.

Same as invention 1 but for Seq.ID:131.

11. Claims: 2, 5-9, 12-18, 21-23, 26, 27, 30-32, 35-41, 44, 45,
48-51, 53, 54 all partially.

Same as invention 1 but for Seq.ID:124.

12. Claims: 2, 5-9, 12-18, 21-23, 26, 27, 30-32, 35-41, 44, 45,
48-51, 53, 54 all partially.

Same as invention 1 but for Seq.ID:132.

13. Claims: 3, 5-9, 12-18, 21-23, 26, 27, 30-32, 35-41, 44, 45,
48-51, 53, 54 all partially.

Same as invention 1 but for Seq.ID:1.

14. Claims: 3, 5-9, 12-18, 21-23, 26, 27, 30-32, 35-41, 44, 45,
48-51, 53, 54 all partially.

Same as invention 1 but for Seq.ID:2.

FURTHER INFORMATION CONTINUED FROM PCT/ISA/ 210

15. Claims: 3, 5-9, 12-18, 21-23, 26, 27, 30-32, 35-41, 44, 45,
48-51, 53, 54 all partially.

Same as invention 1 but for Seq.ID:4 and 17.

16. Claims: 3, 5-9, 12-18, 21-23, 26, 27, 30-32, 35-41, 44, 45,
48-51, 53, 54 all partially.

Same as invention 1 but for Seq.ID:5.

17. Claims: 3, 5-9, 12-18, 21-23, 26, 27, 30-32, 35-41, 44, 45,
48-51, 53, 54 all partially.

Same as invention 1 but for Seq.ID:6.

18. Claims: 3, 5-9, 12-18, 21-23, 26, 27, 30-32, 35-41, 44, 45,
48-51, 53, 54 all partially.

Same as invention 1 but for Seq.ID:7.

19. Claims: 3, 5-9, 12-18, 21-23, 26, 27, 30-32, 35-41, 44, 45,
48-51, 53, 54 all partially.

Same as invention 1 but for Seq.ID:8.

20. Claims: 3, 5-9, 12-18, 21-23, 26, 27, 30-32, 35-41, 44, 45,
48-51, 53, 54 all partially.

Same as invention 1 but for Seq.ID:9.

21. Claims: 3, 5-9, 12-18, 21-23, 26, 27, 30-32, 35-41, 44, 45,
48-51, 53, 54 all partially.

Same as invention 1 but for Seq.ID:10 and 13.

22. Claims: 3, 5-9, 12-18, 21-23, 26, 27, 30-32, 35-41, 44, 45,
48-51, 53, 54 all partially.

Same as invention 1 but for Seq.ID:14.

FURTHER INFORMATION CONTINUED FROM PCT/ISA/ 210

23. Claims: 3, 5-9, 12-18, 21-23, 26, 27, 30-32, 35-41, 44, 45,
48-51, 53, 54 all partially.

Same as invention 1 but for Seq.ID:15.

24. Claims: 3, 5-9, 12-18, 21-23, 26, 27, 30-32, 35-41, 44, 45,
48-51, 53, 54 all partially.

Same as invention 1 but for Seq.ID:16.

25. Claims: 3, 5-9, 12-18, 21-23, 26, 27, 30-32, 35-41, 44, 45,
48-51, 53, 54 all partially.

Same as invention 1 but for Seq.ID:18.

26. Claims: 3, 5-9, 12-18, 21-23, 26, 27, 30-32, 35-41, 44, 45,
48-51, 53, 54 all partially.

Same as invention 1 but for Seq.ID:19.

27. Claims: 3, 5-9, 12-18, 21-23, 26, 27, 30-32, 35-41, 44, 45,
48-51, 53, 54 all partially.

Same as invention 1 but for Seq.ID:20.

28. Claims: 3, 5-9, 12-18, 21-23, 26, 27, 30-32, 35-41, 44, 45,
48-51, 53, 54 all partially.

Same as invention 1 but for Seq.ID:21.

29. Claims: 3, 5-9, 12-18, 21-23, 26, 27, 30-32, 35-41, 44, 45,
48-51, 53, 54 all partially.

Same as invention 1 but for Seq.ID:22.

30. Claims: 3, 5-9, 12-18, 21-23, 26, 27, 30-32, 35-41, 44, 45,
48-51, 53, 54 all partially.

Same as invention 1 but for Seq.ID:23.

/.

FURTHER INFORMATION CONTINUED FROM PCT/ISA/ 210

31. Claims: 4-9, 12-18, 21-23, 26, 27, 30-32, 35-41, 44, 45,
48-51, 53, 54 all partially.

Same as invention 1 but for Seq.ID:26.

32. Claims: 4-9, 12-18, 21-23, 26, 27, 30-32, 35-41, 44, 45,
48-51, 53, 54 all partially.

Same as invention 1 but for Seq.ID:27.

33. Claims: 4-9, 12-18, 21-23, 26, 27, 30-32, 35-41, 44, 45,
48-51, 53, 54 all partially.

Same as invention 1 but for Seq.ID:28.

34. Claims: 4-9, 12-18, 21-23, 26, 27, 30-32, 35-41, 44, 45,
48-51, 53, 54 all partially.

Same as invention 1 but for Seq.ID:29.

35. Claims: 4-9, 12-18, 21-23, 26, 27, 30-32, 35-41, 44, 45,
48-51, 53, 54 all partially.

Same as invention 1 but for Seq.ID:30.

36. Claims: 4-9, 12-18, 21-23, 26, 27, 30-32, 35-41, 44, 45,
48-51, 53, 54 all partially.

Same as invention 1 but for Seq.ID:31.

37. Claims: 4-9, 12-18, 21-23, 26, 27, 30-32, 35-41, 44, 45,
48-51, 53, 54 all partially.

Same as invention 1 but for Seq.ID:32.

38. Claims: 4-9, 12-18, 21-23, 26, 27, 30-32, 35-41, 44, 45,
48-51, 53, 54 all partially.

Same as invention 1 but for Seq.ID:33.

FURTHER INFORMATION CONTINUED FROM PCT/ISA/ 210

39. Claims: 4-9, 12-18, 21-23, 26, 27, 30-32, 35-41, 44, 45,
48-51, 53, 54 all partially.

Same as invention 1 but for Seq.ID:34.

40. Claims: 4-9, 12-18, 21-23, 26, 27, 30-32, 35-41, 44, 45,
48-51, 53, 54 all partially.

Same as invention 1 but for Seq.ID:35.

41. Claims: 4-9, 12-18, 21-23, 26, 27, 30-32, 35-41, 44, 45,
48-51, 53, 54 all partially.

Same as invention 1 but for Seq.ID:36.

42. Claims: 4-9, 12-18, 21-23, 26, 27, 30-32, 35-41, 44, 45,
48-51, 53, 54 all partially.

Same as invention 1 but for Seq.ID:37.

43. Claims: 4-9, 12-18, 21-23, 26, 27, 30-32, 35-41, 44, 45,
48-51, 53, 54 all partially.

Same as invention 1 but for Seq.ID:38.

44. Claims: 4-9, 12-18, 21-23, 26, 27, 30-32, 35-41, 44, 45,
48-51, 53, 54 all partially.

Same as invention 1 but for Seq.ID:39.

45. Claims: 4-9, 12-18, 21-23, 26, 27, 30-32, 35-41, 44, 45,
48-51, 53, 54 all partially.

Same as invention 1 but for Seq.ID:40.

46. Claims: 4-9, 12-18, 21-23, 26, 27, 30-32, 35-41, 44, 45,
48-51, 53, 54 all partially.

Same as invention 1 but for Seq.ID:41.

/.

FURTHER INFORMATION CONTINUED FROM PCT/ISA/ 210

47. Claims: 4-9, 12-18, 21-23, 26, 27, 30-32, 35-41, 44, 45,
48-51, 53, 54 all partially.

Same as invention 1 but for Seq.ID:42.

48. Claims: 4-9, 12-18, 21-23, 26, 27, 30-32, 35-41, 44, 45,
48-51, 53, 54 all partially.

Same as invention 1 but for Seq.ID:43, 44 and 178.

49. Claims: 4-9, 12-18, 21-23, 26, 27, 30-32, 35-41, 44, 45,
48-51, 53, 54 all partially.

Same as invention 1 but for Seq.ID:45.

50. Claims: 4-9, 12-18, 21-23, 26, 27, 30-32, 35-41, 44, 45,
48-51, 53, 54 all partially.

Same as invention 1 but for Seq.ID:46.

51. Claims: 4-9, 12-18, 21-23, 26, 27, 30-32, 35-41, 44, 45,
48-51, 53, 54 all partially.

Same as invention 1 but for Seq.ID:47.

52. Claims: 4-9, 12-18, 21-23, 26, 27, 30-32, 35-41, 44, 45,
48-51, 53, 54 all partially.

Same as invention 1 but for Seq.ID:48.

53. Claims: 4-9, 12-18, 21-23, 26, 27, 30-32, 35-41, 44, 45,
48-51, 53, 54 all partially.

Same as invention 1 but for Seq.ID:49.

54. Claims: 4-9, 12-18, 21-23, 26, 27, 30-32, 35-41, 44, 45,
48-51, 53, 54 all partially.

Same as invention 1 but for Seq.ID:50.

FURTHER INFORMATION CONTINUED FROM PCT/ISA/ 210

55. Claims: 4-9, 12-18, 21-23, 26, 27, 30-32, 35-41, 44, 45,
48-51, 53, 54 all partially.

Same as invention 1 but for Seq.ID:51.

56. Claims: 4-9, 12-18, 21-23, 26, 27, 30-32, 35-41, 44, 45,
48-51, 53, 54 all partially.

Same as invention 1 but for Seq.ID:133.

57. Claims: 4-9, 12-18, 21-23, 26, 27, 30-32, 35-41, 44, 45,
48-51, 53, 54 all partially.

Same as invention 1 but for Seq.ID:134.

58. Claims: 4-9, 12-18, 21-23, 26, 27, 30-32, 35-41, 44, 45,
48-51, 53, 54 all partially.

Same as invention 1 but for Seq.ID:158.

59. Claims: 4-9, 12-18, 21-23, 26, 27, 30-32, 35-41, 44, 45,
48-51, 53, 54 all partially.

Same as invention 1 but for Seq.ID:159.

60. Claims: 4-9, 12-18, 21-23, 26, 27, 30-32, 35-41, 44, 45,
48-51, 53, 54 all partially.

Same as invention 1 but for Seq.ID:160.

61. Claims: 4-9, 12-18, 21-23, 26, 27, 30-32, 35-41, 44, 45,
48-51, 53, 54 all partially.

Same as invention 1 but for Seq.ID:161.

62. Claims: 4-9, 12-18, 21-23, 26, 27, 30-32, 35-41, 44, 45,
48-51, 53, 54 all partially.

Same as invention 1 but for Seq.ID:162.

FURTHER INFORMATION CONTINUED FROM PCT/ISA/ 210

63. Claims: 4-9, 12-18, 21-23, 26, 27, 30-32, 35-41, 44, 45,
48-51, 53, 54 all partially.

Same as invention 1 but for Seq.ID:163.

64. Claims: 4-9, 12-18, 21-23, 26, 27, 30-32, 35-41, 44, 45,
48-51, 53, 54 all partially.

Same as invention 1 but for Seq.ID:164 and 165.

65. Claims: 4-9, 12-18, 21-23, 26, 27, 30-32, 35-41, 44, 45,
48-51, 53, 54 all partially.

Same as invention 1 but for Seq.ID:166 and 167.

66. Claims: 4-9, 12-18, 21-23, 26, 27, 30-32, 35-41, 44, 45,
48-51, 53, 54 all partially.

Same as invention 1 but for Seq.ID:168 and 169.

67. Claims: 4-9, 12-18, 21-23, 26, 27, 30-32, 35-41, 44, 45,
48-51, 53, 54 all partially.

Same as invention 1 but for Seq.ID:170 and 171.

68. Claims: 4-9, 12-18, 21-23, 26, 27, 30-32, 35-41, 44, 45,
48-51, 53, 54 all partially.

Same as invention 1 but for Seq.ID:172 and 173.

69. Claims: 4-9, 12-18, 21-23, 26, 27, 30-32, 35-41, 44, 45,
48-51, 53, 54 all partially.

Same as invention 1 but for Seq.ID:174 and 175.

70. Claims: 4-9, 12-18, 21-23, 26, 27, 30-32, 35-41, 44, 45,
48-51, 53, 54 all partially.

Same as invention 1 but for Seq.ID:176 and 177.

/.

FURTHER INFORMATION CONTINUED FROM PCT/ISA/ 210

71. Claims: 4-9, 12-18, 21-23, 26, 27, 30-32, 35-41, 44, 45, 48-51, 53, 54 all partially.

Same as invention 1 but for Seq.ID:196.

72. Claims: 10, 12-16, 28, 30, 31, 33, 35-39, 52, 54 all partially.

A method for detecting M. tuberculosis infection in a biological sample by detection of antibodies binding to a polypeptide having an N-terminal sequence as in Seq.ID:129, or by detection of a protein or polypeptide that binds to an agent binding to a polypeptide having an N-terminal sequence as in Seq.ID:129. Diagnostic kits thereof. A fusion protein comprising said polypeptide. Diagnostic kit comprising said fusion protein.

73. Claims: 10, 12-16, 28, 30, 31, 33, 35-39, 52, 54 all partially.

Same as invention 72 but for Seq.ID:130.

74. Claims: 11-16, 19-21, 24-26, 29-31, 34-39, 42, 43, 46, 47 all partially.

A method for detecting M. tuberculosis infection in a biological sample by detection of antibodies binding to a polypeptide encoded by a DNA sequence consisting of Seq.ID:3, complements or hybridizing sequences. A method for detecting M. tuberculosis infection in a biological sample by detection of said DNA sequence. A method for detecting M. tuberculosis infection in a biological sample by detection of a protein or polypeptide that binds to an agent binding to a polypeptide encoded by Seq.ID:3, complements or hybridizing sequences. Diagnostic kits thereof.

75. Claims: 11-16, 19-21, 24-26, 29-31, 34-39, 42, 43, 46, 47 all partially.

Same as invention 74 but for Seq.ID:11.

76. Claims: 11-16, 19-21, 24-26, 29-31, 34-39, 42, 43, 46, 47 all partially.

Same as invention 74 but for Seq.ID:12.

FURTHER INFORMATION CONTINUED FROM PCT/ISA/ 210

77. Claims: 11-16, 19-21, 24-26, 29-31, 34-39, 42, 43, 46,
47 all partially.

Same as invention 74 but for Seq.ID:135.

78. Claims: 11-16, 19-21, 24-26, 29-31, 34-39, 42, 43, 46,
47 all partially.

Same as invention 74 but for Seq.ID:136.

79. Claims: 11-16, 19-21, 24-26, 29-31, 34-39, 42, 43, 46,
47 all partially.

Same as invention 74 but for Seq.ID:151.

80. Claims: 11-16, 19-21, 24-26, 29-31, 34-39, 42, 43, 46,
47 all partially.

Same as invention 74 but for Seq.ID:152.

81. Claims: 11-16, 19-21, 24-26, 29-31, 34-39, 42, 43, 46,
47 all partially.

Same as invention 74 but for Seq.ID:153.

82. Claims: 11-16, 19-21, 24-26, 29-31, 34-39, 42, 43, 46,
47 all partially.

Same as invention 74 but for Seq.ID:154 and 155.

83. Claims: 11-16, 19-21, 24-26, 29-31, 34-39, 42, 43, 46,
47 all partially.

Same as invention 74 but for Seq.ID:184.

84. Claims: 11-16, 19-21, 24-26, 29-31, 34-39, 42, 43, 46,
47 all partially.

Same as invention 74 but for Seq.ID:185.

FURTHER INFORMATION CONTINUED FROM PCT/ISA/ 210

85. Claims: 11-16, 19-21, 24-26, 29-31, 34-39, 42, 43, 46,
47 all partially.

Same as invention 74 but for Seq.ID:186.

86. Claims: 11-16, 19-21, 24-26, 29-31, 34-39, 42, 43, 46,
47 all partially.

Same as invention 74 but for Seq.ID:187.

87. Claims: 11-16, 19-21, 24-26, 29-31, 34-39, 42, 43, 46,
47 all partially.

Same as invention 74 but for Seq.ID:188.

88. Claims: 11-16, 19-21, 24-26, 29-31, 34-39, 42, 43, 46,
47 all partially.

Same as invention 74 but for Seq.ID:194 and 195.

89. Claims: 11-16, 19-21, 24-26, 29-31, 34-39, 42, 43, 46,
47 all partially.

Same as invention 74 but for Seq.ID:198.

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/US 97/18214

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
EP 0419355 A	27-03-91	AU 6414390 A CA 2042016 A JP 9234096 A JP 2756368 B JP 4501811 T	18-04-91 20-03-91 09-09-97 25-05-98 02-04-92
WO 9501441 A	12-01-95	AU 682879 B AU 7068894 A CA 2165949 A EP 0706571 A NZ 267984 A	23-10-97 24-01-95 12-01-95 17-04-96 22-09-97
WO 9501440 A	12-01-95	AU 685133 B AU 7068694 A EP 0749486 A	15-01-98 24-01-95 27-12-96
WO 9623885 A	08-08-96	US 5714593 A AU 4667596 A CA 2210928 A EP 0807178 A	03-02-98 21-08-96 08-08-96 19-11-97
WO 9221758 A	10-12-92	FR 2677365 A CA 2110389 A EP 0589943 A JP 6508513 T	11-12-92 10-12-92 06-04-94 29-09-94
WO 9400493 A	06-01-94	US 5330754 A AU 689075 B AU 4651193 A EP 0649435 A JP 7508649 T US 5559011 A	19-07-94 26-03-98 24-01-94 26-04-95 28-09-95 24-09-96
FR 2265402 A	24-10-75	NONE	
FR 2244539 A	18-04-75	NONE	
WO 9709429 A	13-03-97	AU 7158796 A	27-03-97
WO 9709428 A	13-03-97	AU 7158696 A	27-03-97

THIS PAGE BLANK (USPTO)